



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Volume: 7 Issue: VI Month of publication: June 2019

DOI: http://doi.org/10.22214/ijraset.2019.6175

www.ijraset.com

Call: © 08813907089 E-mail ID: ijraset@gmail.com

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com

Effect of Meat Consumption on Prevalence of Alzheimer's Disease

Akash Singh

Dept. of Biochemistry, Deshbandhu College, University of Delhi

Abstract: Meat has become a prominent portion in nutritional diet across continents. Just this year world has consumed more than 13 crore tons of meat, according to The World Counts and this number is increasing with every second passing by. The major meat consuming countries are United States of America(USA), European Region, Latin America and Australia. Same countries top the chart when it comes to Alzheimer's disease (AD) prevalence. AD is a neuro-degenerative condition that is caused by accumulation of β amyloid plaques and neurofibrillary tangles leading t progressive mental degradation causing dementia as its major symptom. Meat on cooking forms Heterocyclic Amines (HCA) that after consumption can generate Reactive Oxygen Species (ROS) that are known to cause oxidative stress and hyper-phosphorylation of tau proteins accelerating this neuro-degenerative condition.

Keywords: Meat consumption, Reactive Oxygen Species, Alzheimer's Disease, Heterocyclic Amines

I. INTRODUCTION

Meat has become a prominent portion of the meal. According to NGC, meat consumption of an average non vegetarian increased from 7% in 1961 to 9% in 2011 i.e. 93 grams in 1961 to 173 grams in 2011. While meat consumption increased from 30.7 kilograms per capita to 41.3 kilograms per capita and will tend to rise to 45.3% by 2030, statistically explains FAO. The major meat consuming countries are United States of America, Europe, Latin America and Australia.

The same countries top the charts when it comes to prevalence of Alzheimer's disease (AD). AD is a neuro-degenerative condition that is caused by accumulation of β amyloid plaques and neurofibrillary tangles, leading to progressive mental deterioration causing dementia as its major symptom. Consumption of meat can be a major factor inducing oxidative stress leading to AD. Keywords Meat consumption, Reactive Oxygen Species, Alzheimer's Disease, Heterocyclic Amines


II. BACKGROUND

According to The World Counts 130,873,729 tons of meat is already consumed globally this year and this number is increasing with every second passing by. It is observed that the countries with highest meat consumption rate also have high prevalence rate of Alzheimer's disease(AD).

III. METHODOLOGIES

Global Statistics of meat consumption was studied taking, prevalence of AD into consideration.

A. Meat Consumption



©NGC, 2011 data



International Journal for Research in Applied Science & Engineering Technology (IJRASET)


ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com

According to FAO meat consumption stats 2011, Japan was leading followed by Australia, Argentina and USA.

Trend as of 2018

According to FAO stats (2018) meat consumption is highest in North America followed by European Union, Latin America and Asia. [5]



©Source: OECD/FAO (2018), "OECD-FAO Agricultural Outlook"

B. Reactive Oxygen Species And Oxidative Stress

Meat contains fat, protein and iron, all of which takes part in oxidation processes. Post oxidation it generates a hydroxyl radical and a hydroperoxyl ion. These free radicals are often termed as Reactive Oxygen Species (ROS) that are highly unstable as they initiate chain of oxidative reactions. [1] Cooking of meat (frying, barbeque, baking) can also generate Heterocyclic amines (HCA) that can also lead to production of ROS resulting in oxidative stress. [2] This oxidative stress can damage brain membrane phospholipids which can be significant for AD as it can affect brain neurons and glial function causing neuro-degeneration. [3] Hence, formation of β amyloid plaques and neurofibrillary tangles. These β amyloid plaque meditated oxidative stress causes hyper-phosphorylation of tau proteins leading to AD. [4].

C. Alzheimer's Disease (AD)

AD characterized by accumulation of β amyloid and intracellular deposition of neurofibrillary tangles of hyper-phosphorylated tau protein. [6] Leading to progressive mental deterioration causing dementia as its major symptom.

D. AD Prevalence around the world

Estimated 46.8 million people worldwide lived with dementia in 2015 and this number is believed to be close to 50 million people in 2017. This number will almost double every 20 years, reaching 75 million in 2030 and 131.5 million in 2050, says Alzheimer's disease International. [7]


The global number of people living with dementia more than doubled from 1990 to 2016. [8]



International Journal for Research in Applied Science & Engineering Technology (IJRASET)


ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177

Volume 7 Issue VI, June 2019- Available at www.ijraset.com



- ©Global Burden of Disease Study 2016 [8]
- E. Alzheimer's Prevalence In Major Meat Consuming Countries
- *United States:* Alzheimer's disease affects an estimated 5.7 million Americans. It is the sixth leading cause of death among all adults and the fifth leading cause for those aged 65 or older. [9]
- Europe: Prevalence of AD in Europe was estimated to be 5.05% (men-3.31% and female-7.13%) [10]
- 3) Latin America and Caribbean: Alzheimer's affect more than 3.4 million people in the region of Latin America and Caribbean.
- 4) India: Rural India validates for lowest rates of Alzheimer's in the world because their diet has more of grains and beans than meat. [12]

AD Trends in different countries vs fat supply Alzheimer's disease prevalence (65+) vs. fat supply



[13] Grant, William. (2014). Grant Alzheimer 1997.

IV. **CONCLUSION**

Americans, Europeans and Latin Americans consume meat as a major portion in their diet while most of Asians (esp. Indians) have more grains and beans than meat in their diet thus accounting for low AD prevalence while US, Europe and Latin America's- 5.7 million, 5.05% and 3.4 million populations respectively suffer from some sort of dementia which accounts for one of the highest prevalence of AD in world concluding that consumption of meat can cause oxidative stress that can lead to Alzheimer's disease.



International Journal for Research in Applied Science & Engineering Technology (IJRASET)

ISSN: 2321-9653; IC Value: 45.98; SJ Impact Factor: 7.177 Volume 7 Issue VI, June 2019- Available at www.ijraset.com

REFERENCES

- [1] Hecke, T., Camp, J. and Smet, S. (2017), Oxidation During Digestion of Meat: Interactions with the Diet and Helicobacter pylori Gastritis, and Implications on Human Health. COMPREHENSIVE REVIEWS IN FOOD SCIENCE AND FOOD SAFETY, 16: 214-233. doi:10.1111/1541-4337.12248
- [2] Carvalho, A., Miranda, A., Santos, F., Loureiro, A., Fisberg, R., & Marchioni, D. (2015). High intake of heterocyclic amines from meat is associated with oxidative stress. British Journal of Nutrition, 113(8), 1301-1307. doi:10.1017/S0007114515000628
- [3] Huang WJ, Zhang X, Chen WW. Role of oxidative stress in Alzheimer's disease. Biomed Rep. 2016;4(5):519-522. doi:10.3892/br.2016.630
- [4] Liu Z, Li T, Li P, et al. The Ambiguous Relationship of Oxidative Stress, Tau Hyperphosphorylation, and Autophagy Dysfunction in Alzheimer's Disease. Oxid Med Cell Longev. 2015;2015:352723. doi:10.1155/2015/352723
- [5] Source: OECD/FAO (2018), "OECD-FAO Agricultural Outlook", OECD Agriculture statistics (database), http://dx.doi.org/10.1787/agr-outl-data-en. 12 http://dx.doi.org/10.1787/888933743328
- [6] Huang WJ, Zhang X, Chen WW. Role of oxidative stress in Alzheimer's disease. Biomed Rep. 2016;4(5):519-522. doi:10.3892/br.2016.63
- [7] Alzheimer's disease International https://www.alz.co.uk/research/statistics
- [8] Global, regional and national burden of AD and other dementia 1996-2016: a systematic analysis for Global Burden of Disease Study 2016 https://doi.org/10.1016/S1474-4422(18)30403-4
- [9] National Centre for Chronic Disease Prevention and Health Promotion https://www.cdc.gov/chronicdisease/resources/publications/aag/alzheimers.htm
- [10] NiuH,Álvarez-ÁlvarezI,Guillén-GrimaF,Aguinaga-OntosoI.PrevalenciaeincidenciadelaenfermedaddeAlzheimerenEuropa:metaanálisis.Neurología.2017;32:523—532
- [11] https://alzgerm.org/news/alzheimers-disease-important-latin-america/
- [12] https://nutritionfacts.org/2015/11/12/where-are-the-lowest-rates-of-alzheimers-in-the-world/









45.98



IMPACT FACTOR: 7.129



IMPACT FACTOR: 7.429



INTERNATIONAL JOURNAL FOR RESEARCH

IN APPLIED SCIENCE & ENGINEERING TECHNOLOGY

Call: 08813907089 🕓 (24*7 Support on Whatsapp)