



# ABO and Rh Blood Group Distribution in İstanbul Province (Turkey)

Zeyneb İrem Yüksel Salduz<sup>1</sup>, Güven Çetin<sup>2</sup>, Cumali Karatoprak<sup>3</sup>, Aclan Özder<sup>4</sup>, Mesut Bilginç<sup>4</sup>, İlhami Gültepe<sup>3</sup>, Ömer Gül<sup>1</sup>

## Abstract

**Objective:** An understanding of the distribution of blood groups in a city is useful for individuals in need. In this study, we aimed to determine the distribution rates of ABO and Rh blood groups in İstanbul Province.

**Methods:** This retrospective study was conducted by investigating the records of 6041 healthy blood donors who visited the Bezmialem Vakıf University Faculty of Medicine, Blood Bank, between 1st January, 2011 and 31<sup>st</sup> December, 2013. ABO and Rh blood groups were examined by lam agglutination, micro plate agglutination, or gel centrifugation methods.

**Results:** In this study, the frequency of the blood groups A, O, B, and AB were found to be 43.44%, 33.02%, 15%, and 8.54%, respectively. Rh (+) positivity was determined in 5192 cases (85.95%) and Rh (-) negativity was determined in 849 cases (14.05%). The frequency of intra-group Rh was found to be 87.84% of Rh (+) and 12.16% of Rh (-) in blood group A, 87.52% of Rh (+) and 14.48% of Rh (-) in blood group O, 82.12% of Rh (+) and 17.88% of Rh (-) in blood group B, and 76.94% of Rh (+) and 23.06% of Rh (-) in blood group AB. A majority of the blood donors were males (92.42%), and 69.28% were in the age group of 25–44 years.

**Conclusion:** ABO and Rh blood group distribution rates in our region were similar to the general rates in Turkey. There was a positive correlation between the frequency of ABO blood group type and Rh positivity rate in each group. We believe that the present study will contribute to the literature with respect to the database of blood groups.

**Keywords:** Blood groups, ABO, Rh, İstanbul

## Introduction

Blood groups are classified into different groups according to the antigens on the surface of erythrocytes (1). The International Society for Blood Transfusion (ISBT) Working Committee catalogue currently lists 339 blood group antigens, of which 297 are clustered within 33 blood group systems (2). The milestone in the modern medical practice of blood transfusion is the identification of blood groups as A, B, and O by Karl Landsteiner in 1901. One year later, the fourth group AB, in which both antigens are present on red blood cells and neither anti-A nor anti-B antibodies are present in serum, was described by Sturli and von Decastelo (3). The histo-blood group ABO that is the major human alloantigen system involves three carbohydrate antigens (4). Antigens of this system are present on the surface of erythrocytes, platelets, and various human tissues including vascular, intestinal, cervical, and glandular mammary epithelial cells; these antigens are also soluble in plasma, saliva, milk, urine, and feces. Additionally, serum contains antibodies to antigens that are not on the surface of erythrocytes. Therefore, ABO system plays the most important role in transfusion and transplantation (5).

Rh blood group system, another important system, is the most polymorphic of all human blood groups and is composed of at least 45 antigens. The discovery of the Rh factor by Landsteiner and Alexander Wiener in 1940 provided the pathophysiological basis for erythroblastosis.

The Rh system is clinically important because of its role in the hemolytic diseases of newborns and in transfusion incompatibility (3).

An understanding of the distribution of blood groups in a city would be helpful for blood bank employees and individuals in need. There are many studies in Turkey on blood group distribution. In this study, we aimed to determine the distribution rates of ABO and Rh blood groups in İstanbul Province and provide new data to the literature.

## Methods

This retrospective study was conducted by investigating the records of 6041 healthy blood donors who applied to the Bezmialem Vakıf University Faculty of Medicine, Blood Bank, between 1<sup>st</sup> January, 2011 and 31<sup>st</sup> December, 2013 after obtaining ethical approval from the ethics committee of the Bezmialem Vakıf University (05.02.2014/25). Written informed consent as "Blood

This study was presented at the 6<sup>th</sup> National Congress of Blood Bank and Transfusion Medicine, 27 November-1 December 2013, Antalya, Türkiye.

<sup>1</sup>Blood Bank, Bezmialem Vakıf University Faculty of Medicine, İstanbul, Türkiye

<sup>2</sup>Department of Internal Medicine and Hematology, Bezmialem Vakıf University Faculty of Medicine, İstanbul, Türkiye

<sup>3</sup>Department of Internal Medicine, Bezmialem Vakıf University Faculty of Medicine, İstanbul, Türkiye

<sup>4</sup>Department of Family Medicine, Bezmialem Vakıf University Faculty of Medicine, İstanbul, Türkiye

### Address for Correspondence:

Zeyneb İrem Yüksel Salduz  
Phone: +90 555 636 78 24  
E-mail: iremsalduz@hotmail.com

Received:  
29.12.2014

Accepted:  
21.04.2015

© Copyright 2015 by Available online at  
www.istanbulmedicaljournal.org

**Table 1. Distribution of ABO and Rh blood groups**

	A		O		B		AB	
	Rh+	Rh-	Rh +	Rh-	Rh +	Rh-	Rh +	Rh-
Intra-group Rh (%)	87.84	12.16	87.52	14.48	82.12	17.88	76.94	23.06
Rh (%)	38.16	5.28	28.90	4.12	12.32	2.68	6.57	1.97
ABO (%)	43.44		33.02		15		8.54	
Rh +/- Ratio	7.22		6.04		4.59		3.33	

donor questionnaire and consent forms" was obtained from the blood donors. ABO and Rh blood groups were determined by the lam agglutination method using Blood Grouping Reagent (Dia-Gast, Loos, France), microplate agglutination method using Galileo System (Stratec, Frankfurt, Germany), or gel centrifugation method using IH-1000 Fully Automated System (Diamed, Cressier, Switzerland).

### Statistical analysis

Statistical analyses were performed with descriptive statistics methods and Pearson's correlation test using Statistical Package for Social Sciences (SPSS) 20.0 (IBM, Armonk New York, USA). A probability value of less than 0.05 was considered to be statistically significant.

### Results

A total of 6041 blood donors were included in our study. As shown in Table 1, the frequencies of blood groups A, O, B, and AB were found to be 43.44% (2624), 33.02% (1995), 15% (906), and 8.54% (516), respectively. In total, 5192 (85.95%) blood donors were found to be Rh (+) positive and 849 (14.05%) were found to be Rh (-) negative. The frequency of intra-group Rh was found to be 87.84% (2305) of Rh (+) and 12.16% (319) of Rh (-) in blood group A, 87.52% (1746) of Rh (+) and 14.48% (249) of Rh (-) in blood group O, 82.12% (744) of Rh (+) and 17.88% (162) of Rh (-) in blood group B, and 76.94% (397) of Rh (+) and 23.06% (119) of Rh (-) in blood group AB. There was a positive correlation between the frequency of ABO blood group type and Rh positivity rate in each group ( $p=0.011$ ). In table 2, the age of the blood donors are categorized into four groups; the number of individuals in each group was as follows: 1071 (17.73%) blood donors were in the age group of 18–24 years, 4185 (69.28%) blood donors were in the age group of 25–44 years, 783 (12.96%) blood donors were in the age group of 45–64 years, and 2 (0.03%) blood donors were in the age group of over 65 years. A total of 5583 (92.42%) blood donors were males and 458 (7.58%) blood donors were females.

### Discussion

The inheritance of ABO and Rh blood groups follows the Mendelian rules. ABO blood group genes are located on the ninth chromosome and the Rh gene is located on the first chromosome (6). The frequency of ABO and Rh blood groups vary according to ethnic groups (7). The worldwide distribution rates of A, O, B, and AB blood group were reported to be 41%, 47%, 9%, and 3%, respectively; in the USA, the distribution rates were 37.10%, 46.70%, 12.10%, and 4.10%; in the UK, 41.78%, 46.63%, 8.56%, and 3.04%; in Greece, 48.19%, 34.21%, 12.04%, and 5.56%; and in Bulgaria, 39.96%, 35.80%, 16.84%, and 7.60%, respectively, (Table 3) (7-9). In table 4, we gave the distribution of ABO and Rh blood groups in some regions of Turkey (9-22). When we look at the distribu-

tion of blood groups in İstanbul Province, it is seen that the frequencies of A, O, B, and AB blood groups were 44.80%, 30.80%, 15.90%, and 8.10%, respectively, by Gül et al. (11) in 2005. The frequencies obtained in our study were lower than those in the study by Gül et al. (11) in terms of blood group A and were higher than that study in terms of blood group O. The differences between these two studies could be related with the recent increase in the emigration rate in İstanbul Province. When we look at the other cities of Turkey, the frequency of blood group A was reported between 36.38% in Şanlıurfa and 45.06% in Konya (12, 13). The distribution rate of blood group A in Eskişehir was found to be 43.52%, which was the rate closest to that found in our results (9). The frequency of blood group O in Turkey was reported between 30.80% in İstanbul and Van and 44.07% in Rize (11, 14, 15). The rate closest to that of our study for blood group O was 33.30% in Denizli and Kayseri (16, 17). The highest rate of blood group B in Turkey was found to be 21.25% in Şanlıurfa and the lowest rate was found to be 9.26% in Rize (12, 15). In this study, the frequency of blood group B was up to 15.00%, which is closest to the frequency of 15.45% in Ankara (18). The AB blood group rate was shown as 9.20% in Van and as 2.60% in Rize, which are the highest and lowest rates in the literature of Turkey (14, 15). In a recent study, the frequency of blood group AB was found to be 8.54% in İstanbul. For blood group AB, Eskişehir had the closest rate to of our city with 8.50% (9). The results of our study are similar to the general rates in Turkey (10).

The rate of Rh positivity was reported to be 85% in Caucasian population (7). In our study, the percentage of Rh positivity and negativity were 85.95% and 14.05%, respectively. In terms of Rh positivity, it was revealed that Eskişehir (86.65%) and Van (86.80%) had a percentage rate closest to that found in our study (9, 14). In addition, there was a positive correlation between the frequency of ABO blood group type and Rh positivity rate in each group ( $p=0.011$ ). The frequencies of Rh positivity were the lowest in the AB blood group (76.94%) and the highest in the A blood group (87.84%) (Table 1).

In our study, a majority of the blood donors were males (92.42%), and 69.28% were in the age group of 25–44 years (Table 2). The reason for this could be that the female individuals in this age group were not accepted as blood donors because of anemia. It was also shown that anemia prevalence was higher, particularly during the fertile period of females depending on the number of parity and gravity increases (23).

Our limitation is that the study was conducted with a relatively small number of volunteers. However, we can say that we included a considerably good number of volunteers in the study by considering the limitation of the study period and that the study was conducted at one center.

**Table 2. Distribution of the blood donors according to age and gender**

Age groups (%)	Male (%)	Female (%)	M/F Ratio
18–24 (17.73)	15.88	1.85	8.58
25–44 (69.28)	64.74	4.54	14.26
45–64 (12.96)	11.77	1.19	9.83
>65 (0.03)	0.03	-	-
Total	92.42	7.58	12.19

**Table 3. Distribution of the ABO blood group in some countries of the world**

Region	Reference	A (%)	O (%)	B (%)	AB (%)
Worldwide	7	41	47	9	3
USA	8	37.10	46.70	12.10	4.10
UK	9	41.78	46.63	8.56	3.04
Greece	9	48.19	34.21	12.04	5.56
Bulgaria	9	39.96	35.80	16.84	7.60

**Table 4. Distribution of the ABO and Rh blood groups in some regions of Turkey**

Region	Reference	A (%)	O (%)	B (%)	AB (%)	Rh+ (%)	Rh- (%)
Turkey (Total)	10	42.84	32.67	16.46	8.03	88.54	11.46
Istanbul	11	44.80	30.80	15.90	8.10	87.20	12.80
Istanbul	Our Study	43.44	33.02	15.00	8.54	85.95	14.05
Şanlıurfa	12	36.38	34.69	21.25	7.68	90.79	9.21
Konya	13	45.06	32.21	15.63	7.12	87.40	12.60
Eskişehir	9	43.52	31.10	16.84	8.50	86.65	13.35
Van	14	43.80	30.80	16.20	9.20	86.80	13.20
Rize	15	44.07	44.07	9.26	2.60	83.70	16.30
Denizli	16	42.60	33.30	16.80	7.40	89.90	10.10
Kayseri	17	44.00	33.30	16.20	6.50	88.20	11.80
Ankara	18	44.62	32.24	15.45	7.69	88.13	11.87
Yozgat	19	44.30	31.70	15.90	8.10	88.00	12.00
Gaziantep	20	40.01	35.09	18.10	6.80	90.83	9.17
Malatya	21	39.32	41.28	13.36	6.04	89.00	11.00
Diyarbakır	22	40.81	33.66	18.53	6.98	89.17	10.82

## Conclusion

ABO and Rh blood group distribution rates in Istanbul are similar to the general rates in Turkey. An understanding of the frequency of blood groups and high Rh positivity rate in common blood groups will be helpful to provide suitable blood storage for individuals in need. We think that the present study will be a valuable contribution to the literature.

**Ethics Committee Approval:** Ethics committee approval was received for this study from the ethics committee of Bezmialem Vakıf University.

**Informed Consent:** Written informed consent was obtained from patient who participated in this study.

**Peer-review:** Externally peer-reviewed.

**Author Contributions:** Concept - Z.İ.Y.S.; Design - Z.İ.Y.S.; Supervision - G.Ç., C.K., A.Ö.; Funding - Z.İ.Y.S., G.Ç.; Materials - M.B., İ.G., Ö.G.; Data Collection and/or Processing - Z.İ.Y.S., Ö.G.; Analysis and/or Interpretation - G.Ç., C.K., A.Ö.; Literature Review - Z.İ.Y.B., M.B., İ.G.; Writer - Z.İ.Y.S.; Critical Review - G.Ç., C.K., A.Ö.; Other - M.B., İ.G., Ö.G.

**Acknowledgements:** The authors thank to all blood bank employees for their contributions to this study.

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study has received no financial support.

## References

- Calhoun L, Petz LD. Erythrocyte antigens and antibodies. In: Beutler E, Lichtman MA, Coller BS, Kipps TJ, Seligsohn D, editors. Williams Hematology. New York: Mc Graw-Hill; 2001. p. 1843-58.
- Storry JR, Castilho L, Daniels G, Flegel WA, Garatty G, Haas M, et al. International Society of Blood Group Terminology: Cancun Report (2012). Vox Sang 2013; 107: 90-6. [CrossRef]
- Schwarz HP, Dörner F. Karl Landsteiner and his major contributions to haematology. Br J Haematol 2003; 121: 556-65. [CrossRef]
- Yamamoto F, Clausen H, White T, Marken J, Hakomori S. Molecular genetic basis of the histo-blood group ABO system. Nature 1990; 345: 229-33. [CrossRef]
- Bilgen H. Kan grup Antijenleri. İ.Ü. Cerrahpaşa Tıp Fakültesi Sürekli Tıp Eğitimi Etkinlikleri Herkes için Transfüzyon tıbbi Sempozyum Dizisi 2005; 40: 55-65.
- Sloan SR, Benjamin RJ, Friedmanlain DF, Webb JJ, Silberstein L. Transfusion Medicine. In: Nathan DG, Ginsburg D, Orkin SH, Look AT, editors. Nathan and Oski's Hematology of Infancy and Childhood. Philadelphia: W.B. Saunders; 2003. p. 1709-56.
- Guyton AC, Hall JE. Blood Types; Transfusion; Tissue and Organ Transplantation. In: Guyton AC, Hall JE, editors. Textbook of Medical Physiology. Philadelphia: W.B. Saunders; 2006. p. 452-53. [CrossRef]
- Garatty G, Glynn SA, McEntire R. ABO and Rh(D) phenotype frequencies of different racial/ethnic groups in the United States. Transfusion 2004; 44: 703-6.
- Gezer S, Akgün N, Akın A, Işık A. Eskişehir bölgesinde ABO kan gruplarının sıklığı. Çocuk Sağlığı ve Hastalıkları Dergisi 1987; 30: 227-31.
- Akbay T, Demiröz P, Güney Ç, Şengül A, Kocabalkan F. Türkiye'de kan gruplarının coğrafi bölgelere göre dağılımı ve bunun kan depolanmasındaki önemi. GATA Bülteni 1989; 31: 391-402.
- Gül M, Sucu R. İ. Uyar T. Şişli Etfal Eğitim ve Araştırma Hastanesi Kan Merkezi Kan Donörlerinin ABO ve Rh Kan Gruplarına göre dağılımları. KSÜ Tıp Fak Derg 2005; 2: 42-4.
- Zerin M, Karakılıç AZ, Nazlıgül Y. Şanlıurfa bölgesinde ABO ve Rh kan gruplarının dağılımı. Harran Tıp Fak Derg 2004; 1: 15-7.
- Çalışkan Ü, Yavuz H, Koç H, Odabaşı D. Konya bölgesinde ABO ve Rh kan gruplarının sıklığı. Selçuk Üniversitesi Tıp Fakültesi Dergisi, 1989; 5: 138-9.
- Dilek İ, Demir C, Bay A, Akdeniz H, Öner AF. ABO and Rh blood groups frequency in men and women living in eastern Turkey. UHOD 2006; 1: 23-6.
- Özkasap S, Dereci S, Şahin K, Dilek AR, Kalyoncuoğlu E, Zengin T, et al. Analysis of ABO and Rh blood groups distribution in East Karadeniz region of Turkey. Dicle Med J 2013; 40: 100-4. [CrossRef]
- Balcı Y, Ovet G, Covut İE, Goncu F, Yılmaz M. ABO and Rh blood groups frequency in Denizli province. UHOD 2010; 2: 103-5. [CrossRef]
- Torun YA, Kaynar LG, Karakükçü Ç, Yay M, Kurnaz F, Mutlu H, et al. ABO and Rh Blood Group Distribution in Kayseri Province, Turkey. Turk J Hematol 2012; 29: 97-8.
- Ergin A, Yardımcı S. Distribution of ABO and Rh blood groups in Turkey. Ankara Üni Tıp Fak Mec 1993; 46: 527-33.
- Kader Ç, Yolcu S, Doğan B, Pınarbaşı M, İlanbey B, Erbay A. ABO and Rh Blood Groups Distribution in Yozgat City, Turkey. J Clin Exp Invest 2014; 5: 169-72. [CrossRef]
- Coşkun Y "ABO"Rh" distribution of blood groups in Gaziantep region. Gaziantep Üniversitesi Tıp Fakültesi Dergisi 1990; 1: 13-8.
- Genç M, Aslan T. ABO ve Rh Kan grupları ve HBsAg, Anti-HIV, VDRL pozitifliği üzerine bir araştırma. Turgut Özal Tıp Merkezi Derg 1997; 4: 139-42.
- Temiz H, Altıntaş A, Gül K. Distribution of ABO and Rh Blood Groups in Diyarbakır. UHOD 2008; 4: 234-7.
- Yıldıran D, Ünal F, Ateşer G, Birtan B. Term Gebelerde Anemi Prevalansı: Kesitsel Bir Çalışma. Istanbul Med J 2009; 10: 26-8.