



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

ORACLE: a module for the description of ORganic Aerosol Composition and Evolution in the atmosphere

A. P. Tsimpidi¹, V. A. Karydis¹, A. Pozzer¹, S. N. Pandis^{2,3}, and J. Lelieveld^{1,4}

¹Max Planck Institute for Chemistry, Mainz, Germany

²Department of Chemical Engineering, University of Patras, Patras, Greece

³Department of Chemical Engineering, Carnegie Mellon University, Pittsburgh, PA, USA

⁴Energy, Environment and Water Research Center, Cyprus Institute, Nicosia, Cyprus

Received: 13 July 2014 – Accepted: 27 July 2014 – Published: 12 August 2014

Correspondence to: A. P. Tsimpidi (a.tsimpidi@mpic.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

GMDD

7, 5465–5515, 2014

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

A computationally efficient module for the description of organic aerosol (OA) partitioning and chemical aging has been developed and implemented into the EMAC atmospheric chemistry-climate model. The model simulates the formation of secondary organic aerosol (SOA) from semi-volatile (SVOCs), intermediate-volatility (IVOCs) and volatile organic compounds (VOCs). The model distinguishes SVOCs from biomass burning and all other combustion sources using two surrogate species for each source category with an effective saturation concentration at 298 K of $C^* = 0.1$ and $10 \mu\text{g m}^{-3}$. Two additional surrogate species with $C^* = 10^3$ and $10^5 \mu\text{g m}^{-3}$ are used for the IVOCs emitted by the above two source categories. Gas-phase photochemical reactions that change the volatility of the organics are taken into account. The oxidation products (SOA-sv, SOA-iv, and SOA-v) of each group of precursors (SVOCs, IVOCs, and VOCs) are simulated separately in the module to keep track of their origin. ORACLE efficiently describes the OA composition and evolution in the atmosphere and can be used to (i) estimate the relative contributions of SOA and primary organic aerosol (POA) to total OA, (ii) determine how SOA concentrations are affected by biogenic and anthropogenic emissions, and (iii) evaluate the effects of photochemical aging and long-range transport on the OA budget. Here we estimate that the predicted domain-average global surface OA concentration is $1.5 \mu\text{g m}^{-3}$ and consists of 7 % POA from fuel combustion, 11 % POA from biomass burning, 2 % SOA-sv from fuel combustion, 3 % SOA-sv from biomass burning, 15 % SOA-iv from fuel combustion, 28 % SOA-iv from biomass burning, 19 % biogenic SOA-v, and 15 % anthropogenic SOA-v. The tropospheric burden of OA components is predicted to be 0.23 Tg POA, 0.16 Tg SOA-sv, 1.41 Tg SOA-iv, and 1.2 Tg SOA-v.

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1 Introduction

Atmospheric aerosols adversely affect human health and play an important role in climate change. Depending on the physicochemical properties, aerosols affect the energy budget of the Earth's atmosphere by scattering and absorbing solar radiation (direct effect) and influencing the reflective properties of clouds, their lifetime, and precipitation formation (indirect effects). Organic aerosol (OA) is an important constituent of atmospheric particulates and varies with geographical region, accounting for 20 to 90 % of the submicron particulate mass (Zhang et al., 2007). However, the understanding of OA sources, atmospheric processing and removal is limited. Given that anthropogenic carbonaceous emissions from developing and emerging economies are expected to dramatically increase in the future (IPCC, 2013), a better understanding of the chemical evolution of OA is essential to reduce the aerosol related uncertainties in global climate simulations and improve air quality and climate assessments.

OA consists of primary material (POA, directly emitted from sources such as fossil fuel combustion and biomass burning) and secondary particulate matter (SOA, formed within the atmosphere from the oxidation of gas-phase precursors). The relative contribution of POA and SOA to the overall OA budget remains controversial. Recent studies show that OA is dominated by SOA not only in tropical regions but also in urban environments (Zhang et al., 2005). The formation of SOA is often underestimated in atmospheric chemistry-climate models (CCMs) and chemistry-transport models (CTMs) (Heald et al., 2005). Robinson et al. (2007) attributed this "unexplained" SOA to the evaporation of POA, oxidation of the resulting vapors in the gas phase and subsequent recondensation, SOA formation from intermediate volatility organic compounds (IVOCs) and chemical aging of the first generation products of the volatile organic compounds (VOCs), which were not taken into account by models. Laboratory and field studies have confirmed that the photo-oxidation of fossil fuel combustion and biomass burning emissions can lead to the formation of substantial SOA mass that cannot be explained by the traditional treatment of SOA in CTMs and CCMs (Kroll and Seinfeld,

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

CCMs have traditionally treated POA and SOA as nonvolatile and nonreactive particles that are emitted directly into the atmosphere (Kanakidou et al., 2005). Most of these models convert “hydrophobic” OA to “hydrophilic” OA using an assumed aging rate constant to account for the chemical conversion of fresh OA to more water soluble compounds and its mixing with other soluble aerosol components without any mechanistic detail. In order to describe the OA volatility changes that accompany this chemical conversion, Donahue et al. (2006) developed the volatility basis-set (VBS) framework.
This framework describes the OA absorptive partitioning, where OA is assumed to be semi-volatile and photochemically reactive and is distributed in logarithmically spaced volatility bins. With this approach, the intermediate and semi-volatile primary emissions and the SOA formation and its chemical aging can be simulated in a common framework that is well suited for regional and global modeling (Murphy and Pandis, 2009; Tsimpidi et al., 2010; Jathar et al., 2011).

Several recent regional scale modeling studies have accounted for the semi-volatile nature and chemical aging of organic compounds demonstrating improvements in reproducing the organic aerosol budget and its chemical resolution (Murphy and Pandis, 2009; Tsimpidi et al., 2010; Hodzic et al., 2010; Tsimpidi et al., 2011; Fountoukis et al., 2011; Shrivastava et al., 2011; Bergstrom et al., 2012; Athanasopoulou et al., 2013; Zhang et al., 2013; Fountoukis et al., 2014). However, such efforts on a global scale are limited. Farina et al. (2010) implemented the VBS in the Goddard Institute for Space Studies General Circulation Model II’ (GISS II’) to describe the formation of SOA from VOCs, while assuming that POA emissions were nonvolatile and non-reactive and not accounting for IVOC emissions. Pye and Seinfeld (2010) estimated the global OA production from emissions of semivolatile organic compounds (SVOCs) and IVOCs using the global CTM GEOS-Chem. However, this study simulated only the first generation of reactions neglecting the subsequent chemical aging. In contrast to other studies, their implementation of semivolatile POA led to decreases in predicted

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

total surface-level OA concentrations due to the partial evaporation of POA upon emission. Jathar et al. (2011) modified the model of Farina et al. (2010) to account explicitly for the semivolatile and reactive POA and predicted a global dominance of SOA, which brought the POA/SOA fractions into better agreement with measurements. Jo et al. (2013) employed the VBS framework in the GEOS-Chem global CTM to study the effect of photochemical aging on global SOA. They found that the model results were in better agreement with all observations relative to the simulations without aging and to those of the traditional two product approach. All of the above studies showed improved representation of POA and/or SOA in the corresponding global CTMs and that the changes brought the model predictions closer to field measurements. However, the number of studies that have employed the recent OA findings is rather limited and in most cases lack one or more potentially important processes (i.e., the aging reactions, the semivolatile character of POA, etc.).

This study aims to improve the description of organic aerosols in large-scale models, making use of the VBS approach and recent developments based on laboratory and field measurements. A new computationally efficient module for the description of organic aerosol composition and evolution in the atmosphere (ORACLE) has been developed and implemented in the ECHAM/MESSy Atmospheric Chemistry (EMAC) model (Jöckel et al., 2006) to simulate POA and SOA formation and growth. ORACLE uses a novel lumping method that allows the use of a small number of species, minimizing the computational cost. At the same time it provides valuable information about the OA sources and physicochemical evolution during its atmospheric lifetime. The ORACLE user has full control of the complexity of the OA scheme and is able to adjust the number of species and reactions, depending on the application and the desired chemical resolution (number of compounds and volatility bins), as well as the physicochemical properties of OA components (aging reaction constants, emission factors, etc.). The application of this advanced OA module can help identify gaps in our understanding of the OA formation and composition, improve the predictive capability of OA and help in

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

identifying the major uncertainties that should be addressed in future experimental and modeling studies.

2 Global model description

2.1 EMAC model

- 5 The ECHAM5/MESSy Atmospheric Chemistry model (EMAC) (Jöckel et al., 2006, 2010) is used as the host model for ORACLE. This is a numerical chemistry and climate simulation system that includes sub-models describing lower and middle atmospheric processes and their interaction with oceans, land and human influences. EMAC combines the Modular Earth Submodel System (MESSy; Jöckel et al., 2005)
- 10 and the 5th generation of the European Centre Hamburg general circulation model (ECHAM5; Röckner et al., 2006). ECHAM5 simulates the atmospheric flow with the prognostic variables vorticity, divergence, temperature, total moisture and the logarithm of the surface pressure, and is integrated in the base model layer of MESSy. The interface structure of MESSy allows the use of different modules for atmospheric chemistry,
- 15 transport, and diagnostic tools. EMAC has been extensively described and evaluated against ground-based and satellite observations, and can be run on a range of spatial resolutions (Jöckel et al., 2006; Pozzer et al., 2012; de Meij et al., 2012). In this study, the applied spectral resolution of the EMAC model is T42L31, corresponding to a horizontal grid resolution of approximately $2.8^\circ \times 2.8^\circ$ and 31 vertical layers extending to
- 20 25 km. EMAC is applied for 6 years covering the period 2004–2009 and the first year is used as spin-up.

The EMAC model simulates gas-phase species online through the Module Efficiently Calculating the Chemistry of the Atmosphere submodel (MECCA; Sander et al., 2011). MECCA calculates online the concentration of the major oxidants (OH , H_2O_2 , NO_2 , and O_3) by using a chemical scheme based on the MIM mechanism (Pöschl et al., 2000) including 218 gas phase reactions, 69 photolysis reactions, and 12 heterogeneous

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



reactions of 129 gases in total. Inorganic aerosol microphysics and gas/particle partitioning are calculated by the Global Modal-aerosol eXtension aerosol module (GMxEx; Pringle et al., 2010). This submodel is computationally efficient and is suitable for medium to long-term simulations with global and regional models.

The aerosol microphysics are described using 7 interacting lognormal modes (4 hydrophilic and 3 hydrophobic modes). The 4 hydrophilic modes cover the full aerosol size spectrum (nucleation, Aitken, accumulation, and coarse modes). Each size range has fixed size boundaries and a variable mean radius. The 3 hydrophobic modes have the same size range apart from the nucleation mode. The aerosol composition within each mode is uniform with size (internally mixed), though the composition can vary between modes (externally mixed). The removal of gas and aerosol species through dry deposition is calculated within the DRYDEP submodel (Kerkweg et al., 2006) based on the big leaf approach. The sedimentation of aerosols is calculated within the SEDI submodel (Kerkweg et al., 2006) using a first order approach. Cloud scavenging of gas and aerosol species is simulated using the SCAV submodel (Tost et al., 2006). The optical properties of aerosols (optical thickness, single scattering albedo, asymmetry factor) are calculated within the AEROPT submodel (Lauer et al., 2007) based on pre-calculated look-up tables from Mie theory. The CLOUD submodel (Jöckel et al., 2006) is used to calculate the cloud cover as well as cloud micro-physics including precipitation. The detailed two-moment liquid and ice-cloud microphysical scheme of Lohmann et al. (2007), which enables a physically based treatment of aerosol-cloud interactions, is used to compute cloud microphysical processes.

3 ORACLE module description

3.1 Module overview

The ORACLE module (i) uses logarithmically-spaced saturation concentration bins to describe the OA components based on their volatility, (ii) simulates the contribution

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



of SVOCs and IVOCs from fuel combustion and biomass burning emissions and anthropogenic and biogenic VOCs to the formation of SOA (iii) monitors the gas-phase photochemical reactions of SOA precursors, (iv) assumes bulk equilibrium between the gas and particulate phases and (v) distributes the OA in size modes. A schematic overview of the ORACLE module and how it is implemented in EMAC is provided in Fig. 1.

3.2 OA terminology

The ORACLE module describes the chemical life of a high number of different species involved in the formation of total atmospheric OA in order to cover and track the distribution of these species in multiple simultaneously occurring phases, their participation in a large number of possible formation pathways (i.e. multigenerational oxidation), their volatility distribution and the origin of their sources. The complexity of this description demands a systematic classification of the organic compounds that will be in line with ongoing field and laboratory studies and model development. Following the Murphy et al. (2014) naming convention for classifying OA components, the ORACLE classification system has the following syntax:

[Source root name] – [initial volatility]

with the “source root name” indicated by an acronym currently used in the field and the “initial volatility” of the species upon emission indicated by an alphabetical lowercase suffix. Table 1 contains a full description of the root terms and suffixes used for identifying the emitted volatility of the organic components described by ORACLE module in this application. The source root name of organic components consist of lowercase alphabetical characters (Modifiers) that describe their source type (i.e. “f” for fuel combustion), followed by three capital letters (Base Term): the first letter declares if the organic component is primary (“P”), coming from primary emissions, or secondary (“S”), formed from the oxidation of gas phase species; the second is always the letter “O” denoting organic; and the third identifies the phase of the species (“A” for

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

aerosol and “G” for gas). The exact names of all the organic compounds simulated by ORACLE are listed in Table 2.

3.3 Volatility basis set theory

The ORACLE module adopts the volatility basis set approach proposed by Donahue et al. (2006). It subdivides the thousands of organic compounds into groups (surrogate species) with logarithmically spaced effective saturation concentrations. This framework abandons the traditional distinction between POA and SOA and allows the EMAC model to efficiently treat both semivolatile primary emissions and SOA production, and then simulate the chemical evolution of these species under a unified framework. The exact volatility resolution defined and used by ORACLE can be controlled through the interface layer of the module, offering the flexibility to investigate different schemes. For the current application, eight surrogate species are used for POA divided into two groups with saturation concentrations C^* at 298 K equal to 10^{-1} , 10^1 , 10^3 , $10^5 \mu\text{g m}^{-3}$: (i) POA from biofuel combustion, fossil fuel combustion, and other urban sources (fPOA) and (ii) POA from biomass burning (bbPOA). The least volatile fraction, at $10^{-1} \mu\text{g m}^{-3}$, describes the low volatility organics in the atmosphere that are mostly in the particulate phase even in remote locations. On the other hand, even under highly polluted conditions the majority of the material in the $10^5 \mu\text{g m}^{-3}$ volatility bin will exist exclusively in the vapor phase. The user has full control of the complexity of the ORACLE module and therefore can easily add bins at either end of the range to describe and investigate unique conditions. A similar approach is followed for SOA that is formed from the VOC emissions. Following the approach of Lane et al. (2008) it is assumed that the oxidation of the anthropogenic and biogenic VOC species (aVOC and bVOC, respectively) results in four products for each precursor distributed in four volatility bins with effective saturation concentrations at 298 K equal to 1, 10 , 10^2 and $10^3 \mu\text{g m}^{-3}$ at 298 K. This range is typically constrained by available smog chamber data (Stanier et al., 2008). A wider range can also be used and tested through the

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



ORACLE interface. The different aerosol types and chemical processes modeled in the proposed framework are illustrated in Fig. 2.

3.4 Emission inventory of OA precursors

VOC emissions: the emissions of isoprene and monoterpenes are calculated online by EMAC with the ONLEM submodel and depend on ecosystem type, temperature, and solar radiation. The AIRSEA submodel estimates the oceanic isoprene emissions from chlorophyll concentrations (Pozzer et al., 2006). In this application, isoprene and monoterpene global emissions are 477 and 62.4 Tg yr⁻¹, respectively. The emissions of the aVOCs that are considered SOA precursors are derived from the IPCC RCP4.5 emission inventory (Clarke et al., 2007). These compounds were not considered by the original gas phase chemistry submodel MECCA. Therefore, six lumped aVOC species have been added to MECCA to assess the aSOA formation in ORACLE. These lumped species are grouped similarly to the Statewide Air Pollution Research Center SAPRC99 chemical mechanism (Carter, 2000) into: two alkane species, two olefins, and two aromatics. The aVOCs that are considered SOA precursors and their annual global emissions are listed in Table 3.

IVOC and SVOC emissions: the EMAC model considers the contribution of fuel combustion and biomass burning to POA emissions. The AEROCOM database is used for the aerosol emissions from fossil and biofuel combustion based on the year 2000 (Dentener et al., 2006). The biomass burning contribution is based on the Global Fire Emissions Database (GFED version 3.1) which is monthly resolved and covers the period 1997–2009 (van der Werf et al., 2010). These emission datasets treat POA as non-reactive and non-volatile. However, the POA concentration upon emission is highly sensitive to ambient conditions, including dilution and temperature (Hildemann et al., 1989; Lipsky and Robinson, 2006). ORACLE accounts for the volatility of POA emissions by distributing the traditional nonvolatile emissions into emissions of SVOCs and IVOCs. SVOCs have saturation concentrations between 0.01 and 100 µg m⁻³ and exist

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


in both the gas and particulate phases under typical ambient conditions. IVOCs have saturation concentrations between 10^3 and $10^6 \mu\text{g m}^{-3}$ and exist in the gas phase.

For the current application, the model distinguishes SVOCs from biomass burning and other combustion sources (biofuel and fossil fuel combustion, and other urban sources) using two surrogate species for each emission category with effective saturation concentration at 298 K of $C^* = 0.1$ and $10 \mu\text{g m}^{-3}$. For IVOCs two additional surrogate species with $C^* = 10^3$ and $10^5 \mu\text{g m}^{-3}$ are used for each of the above two source categories. The emission factors used for the distribution of traditional POA emissions into SVOCs and IVOCs are based on the work of Tsimpidi et al. (2010). These emission factors also account for the additional IVOC emissions that were not included in the original emission inventory used by EMAC. Traditional emission inventories account only for a small fraction of the IVOCs since they are based on samples using quartz and/or Teflon filters collected at aerosol concentrations up to $10^4 \mu\text{g m}^{-3}$ (Shrivastava et al., 2008; Robinson et al., 2010). The amount of IVOC emissions missing from traditional inventories is estimated to be between 0.25 and 2.8 times the traditional POA emissions (Schauer et al., 1999, 2001, 2002). In this study, we assume that the missing IVOC emissions are 1.5 times the traditional emission inventory (Shrivastava et al., 2008; Tsimpidi et al., 2010) and are assigned to the fourth volatility bin with $C^* = 10^5 \mu\text{g m}^{-3}$. Table 4 lists the emission factors for each SVOC and IVOC and their total annual global emissions from fuel combustion and biomass burning.

3.5 Photochemical reactions

Photo-oxidation of VOCs: the photochemical oxidation of VOCs that are considered as SOA precursors has been implemented into MECCA. The updated mechanism considers the oxidation of alkenes, aromatics and isoprene by OH, and the oxidation of olefins and monoterpene by O_3 , O, OH, and NO_3 . The oxidation products from anthropogenic (alkenes, aromatics, and olefins) and biogenic (monoterpene and isoprene) VOCs are lumped into two groups: (1) secondary organic aerosols from the oxidation of anthropogenic VOCs (aSOA-v), and (2) secondary organic aerosols from the oxidation of

biogenic VOCs (bSOA-v). These groups are further distributed into volatility bins with logarithmically spaced effective saturation concentrations (Fig. 2). The photo-oxidation of aVOCs and bVOCs is described by the following reactions:



where a_i is the aerosol yield, i is the corresponding volatility bin, and n is the total number of volatility bins. In this application, four volatility bins are considered and the corresponding aerosol yields are those used by Tsimpidi et al. (2010) based on laboratory results from smog-chamber experiments under high- NO_x conditions for aVOCs and low- NO_x conditions for bVOCs (Table 5). This is based on the assumption that urban areas are characterized by high- NO_x conditions (Tsimpidi et al., 2008; Karl et al., 2009) and forested regions by low- NO_x conditions (Pugh et al., 2010; Browne et al., 2013).

Chemical Aging Reactions: ORACLE treats all OA compounds as chemically reactive. However, only homogeneous gas-phase aging is considered since it occurs rapidly compared to heterogeneous reactions with OH (Donahue et al., 2013). The volatilities of aSOA-v reacting with OH, are reduced by a factor of 10 (Fig. 2) with a rate constant of $1 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Tsimpidi et al., 2010) and a 7.5 % increase in mass to account for the added oxygen. aSOA-v can participate in up to three generations of oxidation reaching a final organic matter to organic carbon ratio (OM/OC) of up to 2.2 (assuming an initial OM/OC of 1.8), which is within the limits (OM/OC: 1.8–2.4) of the observed OM/OC of the oxygenated organic aerosols (Aiken et al., 2008). It is assumed that the chemical aging of bSOA-v does not result in any increase of the SOA

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

concentrations due to a balance between functionalization and fragmentation reactions (Ng et al., 2006; Fountoukis et al., 2011). The chemical aging of aSOA-v is described by the following reaction:



The volatilities of SVOCs and IVOCs are reduced by a factor of 100 (Fig. 2) as a result of the OH reaction with a rate constant of $2 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Pye and Seinfeld, 2010) and a 15 % increase in mass to account for the added oxygen. This represents a more aggressive addition of oxygen and reduction in volatility compared to aSOA-v. In the present application, SVOCs and IVOCs can participate in up to three generations of oxidation reaching a final OM/OC of up to 1.8 (assuming an initial OM/OC of 1.2), which is consistent with the observed OM/OC (OM/OC: 1.8–2.4) of the oxygenated organic aerosols (Aiken et al., 2008). The oxidation products of SVOCs and IVOCs are called SOA from the oxidation of fuel combustion and biomass burning SVOCs (fSOA-sv and bbSOA-sv, respectively) and SOA from the oxidation of fuel and biomass burning IVOCs (fSOA-iv and bbSOA-iv, respectively). The photo-oxidation of SVOCs and IVOCs is described by the following reactions:

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

where i is the corresponding volatility bin. The products of reactions (R7)–(R14) are allowed to be further oxidized by OH forming species with lower saturation concentration (i.e. the oxidation of fSOA- sv_i will produce the fSOA- sv_{i-1}).

3.6 Gas/aerosol partitioning

- 5 ORACLE calculates the partitioning of organic compounds between the gas and particle phases by assuming bulk equilibrium between the two phases and that all organic compounds form a pseudo-ideal solution. The gas/aerosol partitioning is performed in two steps as follows:

10 *Bulk equilibrium:* ORACLE calculates the bulk equilibrium gas and aerosol concentrations following the approach of the SOAM II model of Strader et al. (1999). Considering partitioning of n organic compounds and assuming pseudo-ideal solution, a set of n nonlinear equations is obtained:

$$c_{a,i} = c_{t,i} - x_i c_i^* \quad \text{for } i = 1, n$$

$$x_i = \frac{c_{a,i}/M_i}{\sum_{j=1}^n c_{a,j}/M_j},$$

15

where $c_{t,i}$ and $c_{a,i}$ are the total and aerosol-phase concentrations of product i in $\mu\text{g m}^{-3}$, respectively, c_i^* is the effective saturation concentration of product i , x_i is the mole fraction of product i in the absorbing organic phase, and M_i is the molecular weight of product i . The temperature dependence of saturation concentrations is described by the Clausius–Clapeyron equation:

$$c_i^* = c_{i,0}^* \frac{T_0}{T} \exp \left[\frac{\Delta H}{R} \left(\frac{1}{T_0} - \frac{1}{T} \right) \right],$$

20 where c_i^* and $c_{i,0}^*$ are the saturation concentrations at temperature T and T_0 , respectively, R is the gas constant, and ΔH is the enthalpy of vaporization. In this application

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a ΔH of 30 kJ mol^{-1} is used for all aSOA-v and bSOA-v species and a ΔH of 106, 94, 82, and 70 kJ mol^{-1} is used for the 10^{-1} , 10^1 , 10^3 , $10^5 \mu\text{g m}^{-3}$ volatility bins, respectively, for all organic compounds from fossil fuel and biomass burning sources (Tsimpidi et al., 2010). ORACLE solves this equation set, which yields the bulk aerosol composition at equilibrium.

Aerosol size distribution: the aerosol size distribution is determined by distributing the change in aerosol mass after the bulk equilibrium into each size mode using a weighting factor (Pandis et al., 1993). Assuming pseudo-ideal solution, the fraction, $f_{i,k}$, of total flux of species i between gas and aerosol phases that condenses onto or evaporates from an aerosol mode k is given by:

$$f_{i,k} = \frac{N_k d_k (c_i - x_{i,k} c_i^*) / (\beta_k + 1)}{\sum_{l=1}^m N_l d_l (c_i - x_{i,l} c_i^*) / (\beta_l + 1)},$$

where N_k and d_k are the number and mean diameter of particles in the mode k , respectively, m is the total number of aerosol modes, $\beta_k = 2\lambda/\alpha d_k$, α is the aerosol accommodation coefficient, and λ is the mean free path of air molecules (Pandis et al., 1993). The above equation is solved iteratively at each time step and determines the OA composition of each mode.


4 Model results

4.1 Total OA concentrations

The model calculated global average surface OA concentration is $1.5 \mu\text{g m}^{-3}$ (Fig. 3). High OA concentrations are predicted over regions affected by biomass burning and biogenic VOC emissions: the tropical forests and savannas of South America (Brazil), Africa (Congo), and Southeast Asia. Considerable OA concentrations are also predicted over the industrialized regions of the Northern Hemisphere (i.e., China, Europe,

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)





and Eastern US), where strong fossil and biofuel combustion related sources are located. The model predicts a continental background OA concentration of $1\text{--}2 \mu\text{g m}^{-3}$, which is mainly formed by the condensation of oxidized low volatility organic gases. These gases are formed from the photochemical aging of VOCs, IVOCs and SVOCs, which have been emitted in the gas phase and have been efficiently transported remote from their sources. The effects of trade wind transport of bbOA off the subtropical west coasts of Africa and America are discernible in Fig. 3. In these regions, the OA concentration is increased by long-range transport of continental air masses to adjacent ocean areas.

10 4.2 Model performance evaluation

Figure 4 shows the comparison of model predictions of OA concentration against measurements from the European Monitoring and Evaluation Programme (EMEP; <http://hilu.no/projects/ccc/onlinedata/pm/>) and the Interagency Monitoring of protected Visual Environments (IMPROVE; http://vista.cira.colostate.edu/improve/Data/IMPROVE/improve_data.htm), as well as against short-term measurement data collected over East Asia as summarized by Jo et al. (2013). Each point of the scatter plots represents a monthly averaged value at a particular monitoring station. The mean bias (MB), mean absolute gross error (MAGE), normalized mean bias (NMB), normalized mean error (NME), and the root mean square error (RMSE) were also calculated (Table 6) to 15 assess the model performance:

$$\begin{aligned} \text{MAGE} &= \frac{1}{N} \sum_{i=1}^N |P_i - O_i| & \text{MB} &= \frac{1}{N} \sum_{i=1}^N (P_i - O_i) \\ \text{NME} &= \frac{\sum_{i=1}^N |P_i - O_i|}{\sum_{i=1}^N O_i} & \text{NMB} &= \frac{\sum_{i=1}^N (P_i - O_i)}{\sum_{i=1}^N O_i} \end{aligned}$$

$$\text{RMSE} = \left[\frac{1}{N} \sum_{i=1}^N (P_i - O_i)^2 \right]^{\frac{1}{2}},$$

where P_i is the predicted value of the OA mass, O_i is the observed value of OA at the same monthly averaged time, and N is the total number of data points used for the comparison. NME (in %) and MAGE (in $\mu\text{g m}^{-3}$) provide an assessment of the overall discrepancy between model predictions and observations, while NMB (in %) and MB (in $\mu\text{g m}^{-3}$) are indicative of systematic errors. RMSE (in $\mu\text{g m}^{-3}$) incorporates both the variance of the prediction and its bias. Both NME and MAGE inherently include the corresponding bias, which is the reason why their magnitude is equal or larger than NMB and MB respectively. For an unbiased prediction, NME and MAGE express the variance. When NME and NMB or MAGE and MB, respectively, are close to each other in magnitude, the discrepancy is explained as a systematic bias rather than scatter. When the magnitude of NME/MAGE is larger than NMB/MB, part of the discrepancy between predictions and observations is explained as scatter.

The model, despite its coarse resolution, captures relative well the monthly average concentrations of OA over the USA, Europe, and Asia (Table 6). This is rather encouraging given the expected uncertainties in the emission inventory and in a number of parameters used by the model such as the emission fractions for POA from fuel combustion and biomass burning, the reaction rates and aerosol yields that accompany the formation of SOA from SVOCs, IVOCs, and VOCs, etc. In addition, the fact that the formation of SOA from aqueous-phase reactions and heterogeneous reactions including processes, like oligomerization, have not been accounted for, adds to uncertainty. However, despite the above weaknesses, the model predictions for total OA concentration are generally in reasonable agreement with the measurements. More precisely, over the USA, the model overpredicts the OA with a NMB of 57 %. The high NME (89 %) indicates that part of the discrepancy between predictions and observations is explained as scatter, which is visible in Fig. 4a. Over Europe, the model underpredicts OA with a NMB of –22 %. However, this underprediction is mostly related to the measurement

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



station Ispra, located in a topographically pronounced part of Italy. The measurements of OA in Ispra are systematically high (up to $22 \mu\text{g m}^{-3}$) and are not captured by the model due to its low spatial resolution (Fig. 4b). Due to the small number of available measurement sites from the EMEP network, this discrepancy over Ispra significantly 5 affects the statistics of the model evaluation over Europe. Similarly, over East Asia, the model underpredicts OA concentrations with an NMB of -36% since it cannot capture the high values measured over Beijing and Shijiazhuang (up to $32 \mu\text{g m}^{-3}$) due to its limited spatial resolution.

In contrast to highly populated areas, long-term observations of OA over forests are 10 unavailable, therefore we have collected OA data measured during the short-term field campaigns of DABEX, DODO, and AMMA over the subtropical West Africa (Capes et al., 2008, 2009), AMAZE-08 and SAMBBA over the Amazonian rainforest (Chen et al., 2009; Brito et al., 2014), and measurement data collected over the Canadian boreal forest (Schwartz et al., 2010; Takahama et al., 2011). Capes et al. (2008) performed aircraft measurements during the DABEX and DODO field experiments (January and February of 2006) over the subtropical West Africa at altitudes up to 4000 m. This period was characterized by intense agricultural fires in the sub-Saharan part of 15 West Africa resulting in high OA concentrations ($4\text{--}16 \mu\text{g m}^{-3}$). EMAC agrees reasonably well with the observations since the predicted OA concentrations during January and February of 2006 range between 2 and $12.5 \mu\text{g m}^{-3}$, mainly due to high biomass 20 burning emissions over the area. During the wet season of the same year (July and August 2006), Capes et al. (2009) carried out aircraft measurements over the subtropical West Africa in the frame of the AMMA project. This period is characterized by low biomass burning emissions, therefore, OA consists according to EMAC mainly of biogenic SOA. The observed median concentration is $1.1 \mu\text{g m}^{-3}$ while EMAC predicts an 25 average value of $2.7 \mu\text{g m}^{-3}$. In the Northwestern Amazon basin, Chen et al. (2009) reported OA concentrations of $0.6 \mu\text{g m}^{-3}$ and $0.9 \mu\text{g m}^{-3}$ on average, during the wet season of 2008 (February and March) as part of the AMAZE-08 experiment. The corresponding predicted average OA concentration by EMAC is $2.7 \mu\text{g m}^{-3}$. During the dry

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

season (September 2012), Brito et al. (2014) reported an average OA concentration of $13.7 \mu\text{g m}^{-3}$ in the Southwestern Amazon basin during the SAMBBA field experiment. Since this value is mostly affected by biomass burning emissions (which vary significantly interannually) and EMAC model applied emissions during the years 2005–2009, a direct comparison between observations and measurements is not possible. However, EMAC captures the increased concentrations during the dry seasons since OA is predicted to vary between $9.8 \mu\text{g m}^{-3}$ and $30 \mu\text{g m}^{-3}$ over the same area during the months of September. Finally, Schwartz et al. (2010) and Takahama et al. (2011) performed measurements of OA over the boreal forest of British Columbia, Canada during May–June 2008 and March–September 2009, respectively. The reported values were $1.3 \mu\text{g m}^{-3}$ (May–June 2008), $0.6 \mu\text{g m}^{-3}$ (March–April 2009), and $4.1 \mu\text{g m}^{-3}$ (May–September 2009). The predicted OA concentrations by EMAC fit well with this range of observations with values of $1.4 \mu\text{g m}^{-3}$, $0.9 \mu\text{g m}^{-3}$, and $4.5 \mu\text{g m}^{-3}$, respectively.

4.3 Primary Organic Aerosol

POA is the fraction of the organic emissions that is emitted and contained in the aerosol phase and has not undergone chemical reactions. POA in the present application is divided into fPOA and bbPOA depending on its source as described in Sect. 3.3. fPOA and bbPOA have high concentrations close to their sources; further downwind the concentrations rapidly decrease due to dilution and evaporation (Fig. 5).

fPOA: the global average surface concentration of fPOA is $0.11 \mu\text{g m}^{-3}$ (Fig. 5a). Fossil fuels and biofuels are used in the industrial, residential, and transport sectors, therefore the fPOA concentration peaks over densely populated and highly industrialized areas. In fact, the highest fPOA concentration is predicted over Eastern China (up to $8 \mu\text{g m}^{-3}$ over Beijing), India and Bangladesh ($1\text{--}4 \mu\text{g m}^{-3}$). In Europe the highest concentration is predicted over Eastern European countries ($1\text{--}2 \mu\text{g m}^{-3}$ over Poland and Romania), Central Europe ($0.5\text{--}1 \mu\text{g m}^{-3}$ over Paris and the Rhine–Ruhr area), and Moscow ($0.5\text{--}1 \mu\text{g m}^{-3}$). Over Africa, fPOA concentrations peak over the western part of the continent ($1\text{--}3 \mu\text{g m}^{-3}$ over Nigeria), while they are also high over the cities of

Title Page	Abstract	Introduction
Conclusions	Tables	References
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
Printer-friendly Version		
Interactive Discussion		



Addis Ababa, Kampala, and Johannesburg ($0.5\text{--}1.5 \mu\text{g m}^{-3}$). Over North America, the highest fPOA levels are predicted over Los Angeles ($0.5\text{--}1 \mu\text{g m}^{-3}$), northeastern USA ($0.5\text{--}1.5 \mu\text{g m}^{-3}$), and Mexico City ($0.5\text{--}1 \mu\text{g m}^{-3}$), while over South America, fPOA is high over Rio de Janeiro ($0.5\text{--}1 \mu\text{g m}^{-3}$).

5 *bbPOA*: the global average surface concentration of bbPOA is $0.12 \mu\text{g m}^{-3}$ (Fig. 5b). The bbPOA levels are affected by emissions from forest, woodland, peatland, and savannah fires as well as agricultural waste burning. Therefore, high concentrations are predicted over the tropical rainforest and savannas in the Amazon and Congo basins ($5\text{--}10 \mu\text{g m}^{-3}$), Southeast Asia ($3\text{--}6 \mu\text{g m}^{-3}$), and the boreal forests of Alaska, Canada, and Russia.

4.4 Secondary Organic Aerosol from SVOCs

SVOCs are emitted in both the aerosol and gas phases and can be transferred from phase to phase due to cooling and dilution as the air masses travel from their sources. SVOCs in the gas phase can be oxidized and possibly re-condense to the aerosol phase forming secondary organic aerosols (SOA-sv). In ORACLE, SOA-sv is subdivided into fSOA-sv and bbSOA-sv depending on its source as described in Sect. 3.5. fSOA-sv and bbSOA-sv concentrations are high according to our simulation downwind of polluted urban areas and megacities and the major rainforests (Fig. 6). However, as most of the SVOCs are in the aerosol phase after emissions (as POA) and remain there without undergoing chemical reactions during their atmospheric lifetime, fSOA-sv and bbSOA-sv concentrations are significantly lower compared to the corresponding fPOA and bbPOA concentrations (Figs. 5 and 6).

20 *fSOA-sv*: the global average surface concentration of fSOA-sv is $0.03 \mu\text{g m}^{-3}$ (Fig. 6a). fSOA-sv disperses over the continents but remains at low levels in most cases (lower than $0.2 \mu\text{g m}^{-3}$). The highest concentrations of fSOA-sv are predicted over India and Bangladesh ($0.3\text{--}0.7 \mu\text{g m}^{-3}$), and in the Beijing area and Nigeria ($0.1\text{--}0.5 \mu\text{g m}^{-3}$).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

bbSOA-sv: the domain average surface concentration of *bbSOA-sv* is $0.05 \mu\text{g m}^{-3}$ (Fig. 6b). *bbSOA-sv* has higher concentrations in the Southern Hemisphere where the main rainforests of the world are located (Amazon, Congo, SE Asia) and the meteorological conditions favor the partition of SVOC emissions into the gas phase where they are subject to oxidation. The highest *bbSOA-sv* concentrations (up to $1 \mu\text{g m}^{-3}$) are predicted over the west coast of Central Africa, affected by the biomass burning emissions from the Congo Basin. These emissions are the source of the relatively high *bbSOA-sv* concentrations ($0.3\text{--}0.7 \mu\text{g m}^{-3}$) that are predicted over the Southern Atlantic Ocean. The predicted concentration of *bbSOA-sv* over the Amazon basin is up to $0.7 \mu\text{g m}^{-3}$. The atmosphere over the Pacific and Indian Oceans is also affected by biomass burning emissions from the Amazon rainforest and the rainforests of south-eastern Asia, respectively (*bbSOA-sv* concentration is $0.1\text{--}0.3 \mu\text{g m}^{-3}$). Over the boreal forests, *bbSOA-sv* reaches low levels (less than $0.1 \mu\text{g m}^{-3}$) as most of the biomass burning SVOCs are emitted directly into the aerosol phase forming *bbPOA*.

4.5 Secondary Organic Aerosol from IVOCs

IVOCs are emitted in the gas phase where they react with OH and produce *SOA-iv*, which is less volatile and more chemically processed secondary organic aerosol. *SOA-iv* is subdivided into *fSOA-iv* and *bbSOA-iv* depending on the source as described in Sect. 3.5. In the current application, IVOC emissions are assumed to be 4 times higher than of SVOC (Table 4). Therefore, IVOCs are a significant source of SOA and the *SOA-iv* concentration by far exceeds that of *SOA-sv* and *POA* (Figs. 5–7). Furthermore, the effective long-range transport of IVOCs results in the formation of significant *SOA-iv* amounts far from the sources (Fig. 7).

fSOA-iv: the global average surface concentration of *fSOA-iv* is $0.22 \mu\text{g m}^{-3}$ (Fig. 7a). This is two times higher than the corresponding average surface concentration of *fPOA*, even if the peak concentrations of *fSOA-iv* and *fPOA* are similar ($7 \mu\text{g m}^{-3}$ over Bangladesh and $8 \mu\text{g m}^{-3}$ over Beijing, respectively). This difference indicates

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

that fSOA-iv is more regionally distributed compared to fPOA, due to the effects of long-range transport of IVOC emissions on fSOA-iv formation. The model predicts a continental background of around $0.5 \mu\text{g m}^{-3}$ for fSOA-iv. The highest fSOA-iv concentrations are predicted over India and Bangladesh ($3\text{--}7 \mu\text{g m}^{-3}$), eastern China ($1\text{--}4 \mu\text{g m}^{-3}$) and western Africa ($1\text{--}4 \mu\text{g m}^{-3}$). The Balkan Peninsula and eastern Mediterranean are strongly influenced by long-range transport of IVOC emissions from Eastern Europe (the fSOA-iv concentration is $0.5\text{--}1 \mu\text{g m}^{-3}$). fSOA-iv concentrations are also significant over the Arabian Peninsula (around $1 \mu\text{g m}^{-3}$), eastern and South Africa ($0.5\text{--}1 \mu\text{g m}^{-3}$). Over North America fSOA-iv concentrations are relatively high over a large area covering the eastern USA, the Californian Peninsula and North Mexico ($0.5\text{--}1 \mu\text{g m}^{-3}$), while over South America high fSOA-iv concentrations are predicted mostly over Rio de Janeiro ($0.5\text{--}1 \mu\text{g m}^{-3}$). Long-range transport is also important for the predicted fSOA-iv concentrations over the oceans. This is mostly evident over the Arabian Sea where the fSOA-iv concentration is $1\text{--}2 \mu\text{g m}^{-3}$, due mainly to long-range transport of fSOA-iv from India. The Atlantic Ocean atmosphere is influenced by IVOC emissions from western Africa and the fSOA-iv concentration over this region is predicted at $0.5\text{--}1 \mu\text{g m}^{-3}$. Over the Pacific Ocean fSOA-iv is sensitive to long-range transport from the Los Angeles and Mexico City areas and from Beijing and other big cities in eastern China, resulting in significant concentrations off the west coast of California (up to $0.5\text{--}1 \mu\text{g m}^{-3}$) and the Pacific Rim (up to $0.5\text{--}1.5 \mu\text{g m}^{-3}$).

bbSOA-iv: the predicted global average surface concentration of bbSOA-iv is $0.42 \mu\text{g m}^{-3}$ (Fig. 7b). Similar to fSOA-iv, bbSOA-iv is two times higher than bbPOA and disperses over a wide area covering most of South America, Central and South Africa, Eastern Russia, Southeastern Asia and Indonesia due to emissions from the major rainforests and savannas in the Southern Hemisphere, the tropics and the boreal forests in the Northern Hemisphere. The highest bbSOA-iv concentrations are predicted over the Amazon and Congo Basins ($5\text{--}10 \mu\text{g m}^{-3}$). bbSOA-iv concentrations are also high over southeastern Asia (up to $1\text{--}5 \mu\text{g m}^{-3}$) and the boreal forests of Alaska, Canada, and Russia ($0.5\text{--}1.5 \mu\text{g m}^{-3}$). The subtropical west coasts of

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Africa, South America and Indonesia are strongly influenced by long-range transport of bbSOA-iv. The strongest transport effect is predicted over the Atlantic Ocean, where biomass burning IVOC emissions can travel thousands of kilometers from the sources (e.g., the Congo Basin), resulting in significant bbSOA-iv concentrations ($2\text{--}6 \mu\text{g m}^{-3}$).

5 4.6 Secondary Organic Aerosol from VOCs

SOA-v is formed from the oxidation of biogenic and anthropogenic VOCs. Global VOC emissions are dominated by biogenic compounds from vegetated areas (Kanakidou et al., 2005). However, anthropogenic VOCs, emitted in urban areas, can also contribute significantly to SOA formation, especially considering the aging reactions of aSOA-v.

10 bSOA-v: the global average surface concentration of bSOA-v is $0.28 \mu\text{g m}^{-3}$ (Fig. 8a). EMAC predicts the highest bSOA-v concentrations over the Amazon basin ($5\text{--}10 \mu\text{g m}^{-3}$) mostly due to the oxidation of isoprene. Over the Congo Basin, where isoprene emissions are similar to monoterpene emissions, the bSOA-v average surface concentration is $3\text{--}6 \mu\text{g m}^{-3}$. bSOA-v concentrations are also relatively high over Southeast Asia ($2\text{--}4 \mu\text{g m}^{-3}$), Southeast USA and Australia ($2\text{--}5 \mu\text{g m}^{-3}$), and Europe ($0.5\text{--}1.5 \mu\text{g m}^{-3}$).

15 aSOA-v: the global average surface concentration of aSOA-v is $0.23 \mu\text{g m}^{-3}$ (Fig. 8b). Photochemical aging adds significantly to aSOA-v since only 10 % of the predicted aSOA-v is formed from the first photooxidation step of aSOA-v. The remainder 90 % is formed through the continued oxidation of aSOA-v. The aggressive aging of aSOA-v results in a wider distribution of aSOA-v than bSOA-v and in a significant contribution of aSOA-v to the total SOA-v (45 %). The aSOA-v is predicted to have a continental background of around $0.5 \mu\text{g m}^{-3}$ and relatively high concentrations in the vicinity of urban areas. The highest concentrations are predicted over India and Bangladesh and the Persian Gulf ($2\text{--}3 \mu\text{g m}^{-3}$). Over India, aromatics dominate the aVOC emissions (57 %), followed by alkanes (29 %), and olefines (14 %). Over the Persian Gulf region, alkenes are the dominant aVOC (65 %), followed by aromatics

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(33 %), and olefines (2 %). However, these numbers do not proportionally reflect the VOC contributions to the aSOA-v formation since different types of aVOC have different aerosol yields. Over Europe, the highest aSOA-v concentrations occur over the Mediterranean Basin ($0.5\text{--}1 \mu\text{g m}^{-3}$) due to the long-range transport of aSOA-v from Central European emissions. Over the USA, aSOA-v concentrations are relatively high over California and the eastern USA (around $1 \mu\text{g m}^{-3}$).
5

4.7 Chemical composition of OA

Table 7 summarizes the predicted global average chemical composition of OA at the surface. According to these results, most of OA is SOA (82 %) formed from the oxidation of organic compounds in the gas phase. SOA consists of 52 % SOA-iv, 42 % SOA-v, and 6 % SOA-sv. The important contribution of SOA-iv to the overall SOA mass is attributed to the fact that high amounts of fuel combustion and biomass burning IVOCS react with atmospheric oxidants to form low volatility products that condense into the particle phase as SOA-iv. This result further emphasizes the importance of the oxidation of IVOCS as an additional source of OA and implies that global and regional models that do not account for IVOCS could underestimate OA formation as much as by 40 %. In addition, the model results highlight the importance of anthropogenic emissions to the global OA formation. In particular, it predicts that one third of OA originates from anthropogenic sources. Even if this fraction may be overestimated due to the aggressive aging of IVOCS and aVOCs applied in this study, it corroborates the findings of recent studies that highlight the potentially large impact of anthropogenic OA, especially SOA, on the global aerosol load (Carslaw et al., 2013; Lee et al., 2013).
10
15
20

At higher altitudes the production of SOA is enhanced since organic gases are efficiently transported vertically and can be oxidized forming lower volatile SOA (Fig. 9). In addition, the produced SOA in higher altitudes has a relatively long lifetime since it is less subjected to wet and dry deposition. This results in a higher fraction of SOA in total OA higher up than at the surface (92 % compared to 82 % at the surface). Furthermore, in contrast to the surface OA composition, the fraction of aSOA-v in the free
25

Title Page
Abstract Introduction
Conclusions References
Tables Figures

◀ ▶
◀ ▶
Back Close
Full Screen / Esc
Printer-friendly Version
Interactive Discussion



troposphere is predicted to be higher than bSOA-v since the latter is not allowed to participate in additional photochemical reactions (Table 7). The predicted tropospheric burden of POA is 0.23 Tg, the sum of SOA-sv and SOA-iv is 1.57 Tg, and SOA-v is 1.2 Tg. Pye and Seinfeld (2010) predicted 0.03 Tg of POA, 0.90 Tg of SOA-sv and SOA-iv, and 5 0.71 Tg of SOA-v, while Jathar et al. (2011) predicted 0.09 Tg of POA, 1.25 Tg of SOA-sv and SOA-iv, and 1.02 Tg of SOA-v. Farina et al. (2010) and Jo et al. (2013) did not account for SOA-sv and SOA-iv and predicted 1.1 Tg and 0.8 Tg of POA, respectively, and 0.98 Tg and 1.16 Tg of SOA-v, respectively.

5 Conclusions

10 A new module describing the organic composition of aerosols and its evolution in the atmosphere has been developed and implemented into the EMAC global model. ORACLE treats fossil fuel, biofuel and biomass burning related aerosol emissions as semi-volatile and chemically reactive, and also accounts for the emissions and oxidation of IVOCs as an additionally important source of OA. Furthermore, it considers the formation of SOA from the oxidation of anthropogenic and biogenic VOC precursors. The 15 model employs the volatility basis set theory to simulate the emissions, chemical reactions and phase partitioning of all OA components.

The comparison of the model results with measured ambient OA mass concentrations illustrates that by treating fuel combustion and biomass burning aerosol emissions 20 as semi-volatile and reactive, and accounting for emissions and oxidation of IVOCs and traditional VOCs, the model realistically predicts the sources, chemistry and properties of OA, and reproduces the measured concentrations of OA over urban and forested areas. In contrast to previous CCMs that treat POA as non-volatile and non-reactive, the ORACLE module predicts that most of the OA is SOA (82 %). Furthermore, approximately 25 one third of OA is predicted to originate from anthropogenic sources. Such high anthropogenic OA fractions challenge some previous results suggesting that anthropogenic sources do not strongly contribute to global OA concentrations.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

On average, model calculated OA at the surface consists of 18 % POA, 5 % SOA-sv, 43 % SOA-iv, and 34 % SOA-v. The tropospheric burdens of OA components are predicted to be 0.23 Tg POA, 0.16 Tg SOA-sv, 1.41 Tg SOA-iv and 1.2 Tg SOA-v. POA concentrations are particularly high close to biomass burning regions (forest and savannah fires) and fuel combustion related sources (urban and industrial regions), though decrease rapidly during atmospheric transport. SOA-sv concentrations are four to five times lower than POA concentrations since most of the SVOCs are emitted directly in the aerosol phase. IVOCs, on the other hand, are predicted to be important SOA precursors, resulting in high SOA-iv concentrations that are widely dispersed over the continents and the tropical oceans, being subject to the long-range transport of IVOC emissions. SOA-v accounts for 34 % of the total SOA concentration and originates from biogenic sources (55 %) and anthropogenic sources (45 %). In the current application, only anthropogenic SOA-v was allowed to participate in aging reactions, which considerably increases the predicted aSOA-v concentration.

The ORACLE module subdivides OA into several compounds allowing the quantification of primary vs. secondary as well as biogenic vs. anthropogenic contributions to OA concentrations. Such fundamental information can shed light on long-term changes in OA abundance, and hence project the effects of OA on future air quality and climate. ORACLE is well suited for studying the climatic impact of OA as it captures the dynamic aging of OA and its atmospheric evolution by becoming increasingly oxidized, less volatile, and more hygroscopic. Future applications therefore also include the effects on cloud condensation nuclei and indirect aerosol effects on climate. Furthermore, the detailed composition of OA, computed by ORACLE, can be used to directly compare model predictions with the latest aerosol mass spectrometer (AMS) factor analysis data, which is planned as a next application.

Overall, ORACLE is a flexible module that efficiently describes the organic aerosol composition and evolution in the atmosphere. Sensitivity studies of the influence of several parameters (such as reaction constants, aerosol yields, SVOC and IVOC emission fractions) on the model predictions will be conducted in a future study and will

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

be used, along with AMS factor analysis data from around the world, to optimize the model configuration in order to further improve the estimates of OA concentrations and investigate their role in air quality and climate.

Acknowledgements. The research leading to these results has received funding from the

- 5 European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement no 226144. V. A. Karydis acknowledges support from a FP7 Marie Curie Career Integration Grant (project reference 618349).

The service charges for this open access publication
10 have been covered by the Max Planck Society.

References

- Aiken, A. C., Decarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42, 4478–4485, 2008.
- 15 Athanasopoulou, E., Vogel, H., Vogel, B., Tsimpidi, A. P., Pandis, S. N., Knote, C., and Fountoukis, C.: Modeling the meteorological and chemical effects of secondary organic aerosols during an EUCAARI campaign, *Atmos. Chem. Phys.*, 13, 625–645, doi:10.5194/acp-13-625-2013, 2013.
- Bergström, R., Denier van der Gon, H. A. C., Prévôt, A. S. H., Yttri, K. E., and Simpson, D.:
20 Modelling of organic aerosols over Europe (2002–2007) using a volatility basis set (VBS) framework: application of different assumptions regarding the formation of secondary organic aerosol, *Atmos. Chem. Phys.*, 12, 8499–8527, doi:10.5194/acp-12-8499-2012, 2012.
- 25 Brito, J., Rizzo, L. V., Morgan, W. T., Coe, H., Johnson, B., Haywood, J., Longo, K., Freitas, S., Andreae, M. O., and Artaxo, P.: Ground based aerosol characterization during the South American Biomass Burning Analysis (SAMBBA) field experiment, *Atmos. Chem. Phys. Discuss.*, 14, 12279–12322, doi:10.5194/acpd-14-12279-2014, 2014.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

- Browne, E. C., Min, K.-E., Wooldridge, P. J., Apel, E., Blake, D. R., Brune, W. H., Cantrell, C. A., Cubison, M. J., Diskin, G. S., Jimenez, J. L., Weinheimer, A. J., Wennberg, P. O., Wisthaler, A., and Cohen, R. C.: Observations of total RONO₂ over the boreal forest: NO_x sinks and HNO₃ sources, *Atmos. Chem. Phys.*, 13, 4543–4562, doi:10.5194/acp-13-4543-2013, 2013.
- Capes, G., Johnson, B., McFiggans, G., Williams, P. I., Haywood, J., and Coe, H.: Aging of biomass burning aerosols over West Africa: aircraft measurements of chemical composition, microphysical properties, and emission ratios, *J. Geophys. Res.-Atmos.*, 113, D00C15, doi:10.1029/2008jd009845, 2008.
- Capes, G., Murphy, J. G., Reeves, C. E., McQuaid, J. B., Hamilton, J. F., Hopkins, J. R., Crosier, J., Williams, P. I., and Coe, H.: Secondary organic aerosol from biogenic VOCs over West Africa during AMMA, *Atmos. Chem. Phys.*, 9, 3841–3850, doi:10.5194/acp-9-3841-2009, 2009.
- Carslaw, K. S., Lee, L. A., Reddington, C. L., Mann, G. W., and Pringle, K. J.: The magnitude and sources of uncertainty in global aerosol, *Faraday Discuss.*, 165, 495–512, 2013.
- Carter, W. P. L.: Documentation of the SAPRC-99 chemical mechanism for VOC reactivity assessment., Report to the California air resources board, available at: <http://www.cert.ucr.edu/~carter/absts.htm#saprc99> (last access: 7 August 2014), 2000.
- Chen, Q., Farmer, D. K., Schneider, J., Zorn, S. R., Heald, C. L., Karl, T. G., Guenther, A., Allan, J. D., Robinson, N., Coe, H., Kimmel, J. R., Pauliquevis, T., Borrmann, S., Poeschl, U., Andreae, M. O., Artaxo, P., Jimenez, J. L., and Martin, S. T.: Mass spectral characterization of submicron biogenic organic particles in the Amazon Basin, *Geophys. Res. Lett.*, 36, L20806, doi:10.1029/2009gl039880, 2009.
- Clarke, L., Edmonds, J., Jacoby, H., Pitcher, H., Reilly, J., and Richels, R.: Scenarios of greenhouse gas emissions and atmospheric concentrations (Part A) and review of integrated scenario development and application (Part B). A report by the US climate change science program and the subcommittee on global change research, 2007.
- de Meij, A., Pozzer, A., Pringle, K. J., Tost, H., and Lelieveld, J.: EMAC model evaluation and analysis of atmospheric aerosol properties and distribution with a focus on the Mediterranean region, *Atmos. Res.*, 114, 38–69, 2012.
- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmos. Chem. Phys.*, 6, 4321–4344, doi:10.5194/acp-6-4321-2006, 2006.

5 Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40, 2635–2643, 2006.

Donahue, N. M., Chuang, W., Epstein, S. A., Kroll, J. H., Worsnop, D. R., Robinson, A. L., Adams, P. J., and Pandis, S. N.: Why do organic aerosols exist? Understanding aerosol lifetimes using the two-dimensional volatility basis set, *Environ. Chem.*, 10, 151–157, 2013.

10 Farina, S. C., Adams, P. J., and Pandis, S. N.: Modeling global secondary organic aerosol formation and processing with the volatility basis set: implications for anthropogenic secondary organic aerosol, *J. Geophys. Res.-Atmos.*, 115, D09202, doi:10.1029/2009jd013046, 2010.

15 Fountoukis, C., Racherla, P. N., Denier van der Gon, H. A. C., Polymeneas, P., Charalampidis, P. E., Pilinis, C., Wiedensohler, A., Dall’Osto, M., O'Dowd, C., and Pandis, S. N.: Evaluation of a three-dimensional chemical transport model (PMCAMx) in the European domain during the EUCAARI May 2008 campaign, *Atmos. Chem. Phys.*, 11, 10331–10347, doi:10.5194/acp-11-10331-2011, 2011.

20 Fountoukis, C., Megaritis, A. G., Skyllakou, K., Charalampidis, P. E., Pilinis, C., Denier van der Gon, H. A. C., Crippa, M., Canonaco, F., Mohr, C., Prévôt, A. S. H., Allan, J. D., Poulain, L., Petäjä, T., Tiitta, P., Carbone, S., Kiendler-Scharr, A., Nemitz, E., O'Dowd, C., Swietlicki, E., and Pandis, S. N.: Organic aerosol concentration and composition over Europe: insights from comparison of regional model predictions with aerosol mass spectrometer factor analysis, *Atmos. Chem. Phys. Discuss.*, 14, 7597–7635, doi:10.5194/acpd-14-7597-2014, 2014.

25 Grieshop, A. P., Logue, J. M., Donahue, N. M., and Robinson, A. L.: Laboratory investigation of photochemical oxidation of organic aerosol from wood fires 1: measurement and simulation of organic aerosol evolution, *Atmos. Chem. Phys.*, 9, 1263–1277, doi:10.5194/acp-9-1263-2009, 2009.

30 Heald, C. L., Jacob, D. J., Park, R. J., Russell, L. M., Huebert, B. J., Seinfeld, J. H., Liao, H., and Weber, R. J.: A large organic aerosol source in the free troposphere missing from current models, *Geophys. Res. Lett.*, 32, L18809, doi:10.1029/2005gl023831, 2005.

Hennigan, C. J., Miracolo, M. A., Engelhart, G. J., May, A. A., Presto, A. A., Lee, T., Sullivan, A. P., McMeeking, G. R., Coe, H., Wold, C. E., Hao, W.-M., Gilman, J. B., Kuster, W. C., de Gouw, J., Schichtel, B. A., Collett Jr., J. L., Kreidenweis, S. M., and Robinson, A. L.:

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chemical and physical transformations of organic aerosol from the photo-oxidation of open biomass burning emissions in an environmental chamber, *Atmos. Chem. Phys.*, 11, 7669–7686, doi:10.5194/acp-11-7669-2011, 2011.

Hildemann, L. M., Cass, G. R., and Markowski, G. R.: A dilution stack sampler for collection of organic aerosol emissions – design, characterization and field-tests, *Aerosol Sci. Tech.*, 10, 193–204, 1989.

Hodzic, A., Jimenez, J. L., Madronich, S., Canagaratna, M. R., DeCarlo, P. F., Kleinman, L., and Fast, J.: Modeling organic aerosols in a megacity: potential contribution of semi-volatile and intermediate volatility primary organic compounds to secondary organic aerosol formation, *Atmos. Chem. Phys.*, 10, 5491–5514, doi:10.5194/acp-10-5491-2010, 2010.

IPCC: (Intergovernmental Panel on Climate Change): The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK and New York, NY, USA, 2013.

Jathar, S. H., Farina, S. C., Robinson, A. L., and Adams, P. J.: The influence of semi-volatile and reactive primary emissions on the abundance and properties of global organic aerosol, *Atmos. Chem. Phys.*, 11, 7727–7746, doi:10.5194/acp-11-7727-2011, 2011.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.

Jo, D. S., Park, R. J., Kim, M. J., and Spracklen, D. V.: Effects of chemical aging on global secondary organic aerosol using the volatility basis set approach, *Atmos. Environ.*, 81, 230–244, 2013.

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) - a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433–444, doi:10.5194/acp-5-433-2005, 2005.

Jöckel, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: consistent simulation of ozone from the surface to the mesosphere, *Atmos. Chem. Phys.*, 6, 5067–5104, doi:10.5194/acp-6-5067-2006, 2006.

Jöckel, P., Kerkweg, A., Pozzer, A., Sander, R., Tost, H., Riede, H., Baumgaertner, A., Gromov, S., and Kern, B.: Development cycle 2 of the Modular Earth Submodel System (MESSy2), *Geosci. Model Dev.*, 3, 717–752, doi:10.5194/gmd-3-717-2010, 2010.

Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.

Karl, M., Tsigaridis, K., Vignati, E., and Dentener, F.: Formation of secondary organic aerosol from isoprene oxidation over Europe, *Atmos. Chem. Phys.*, 9, 7003–7030, doi:10.5194/acp-9-7003-2009, 2009.

Kerkweg, A., Buchholz, J., Ganzeveld, L., Pozzer, A., Tost, H., and Jöckel, P.: Technical Note: An implementation of the dry removal processes DRY DEPosition and SEDimentation in the Modular Earth Submodel System (MESSy), *Atmos. Chem. Phys.*, 6, 4617–4632, doi:10.5194/acp-6-4617-2006, 2006.

Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593–3624, 2008.

Lane, T. E., Donahue, N. M., and Pandis, S. N.: Simulating secondary organic aerosol formation using the volatility basis-set approach in a chemical transport model, *Atmos. Environ.*, 42, 7439–7451, 2008.

Lauer, A., Eyring, V., Hendricks, J., Jöckel, P., and Lohmann, U.: Global model simulations of the impact of ocean-going ships on aerosols, clouds, and the radiation budget, *Atmos. Chem. Phys.*, 7, 5061–5079, doi:10.5194/acp-7-5061-2007, 2007.

Lee, L. A., Pringle, K. J., Reddington, C. L., Mann, G. W., Stier, P., Spracklen, D. V., Pierce, J. R., and Carslaw, K. S.: The magnitude and causes of uncertainty in global model simulations of

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Pozzer, A., Jöckel, P., Sander, R., Williams, J., Ganzeveld, L., and Lelieveld, J.: Technical Note: The MESSy-submodel AIRSEA calculating the air-sea exchange of chemical species, *Atmos. Chem. Phys.*, 6, 5435–5444, doi:10.5194/acp-6-5435-2006, 2006.

Pozzer, A., de Meij, A., Pringle, K. J., Tost, H., Doering, U. M., van Aardenne, J., and Lelieveld, J.: Distributions and regional budgets of aerosols and their precursors simulated with the EMAC chemistry-climate model, *Atmos. Chem. Phys.* 12, 961–987, 2012.

Pringle, K. J., Tost, H., Message, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P., Vignati, E., and Lelieveld, J.: Description and evaluation of GMXe: a new aerosol submodel for global simulations (v1), *Geosci. Model Dev.*, 3, 391–412, doi:10.5194/gmd-3-391-2010, 2010.

Pugh, T. A. M., MacKenzie, A. R., Hewitt, C. N., Langford, B., Edwards, P. M., Furneaux, K. L., Heard, D. E., Hopkins, J. R., Jones, C. E., Karunaharan, A., Lee, J., Mills, G., Misztal, P., Moller, S., Monks, P. S., and Whalley, L. K.: Simulating atmospheric composition over a South-East Asian tropical rainforest: performance of a chemistry box model, *Atmos. Chem. Phys.*, 10, 279–298, doi:10.5194/acp-10-279-2010, 2010.

Pye, H. O. T. and Seinfeld, J. H.: A global perspective on aerosol from low-volatility organic compounds, *Atmos. Chem. Phys.*, 10, 4377–4401, doi:10.5194/acp-10-4377-2010, 2010.

Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking organic aerosols: semivolatile emissions and photochemical aging, *Science*, 315, 1259–1262, 2007.

Robinson, A. L., Grieshop, A. P., Donahue, N. M., and Hunt, S. W.: Updating the conceptual model for fine particle mass emissions from combustion systems, *J. Air Waste Manage.*, 60, 1204–1222, 2010.

Röckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L., Manzini, E., Schlese, U., and Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, *J. Climate*, 19, 3771–3791, 2006.

Sander, R., Baumgaertner, A., Gromov, S., Harder, H., Jöckel, P., Kerkweg, A., Kubistin, D., Regelin, E., Riede, H., Sandu, A., Taraborrelli, D., Tost, H., and Xie, Z.-Q.: The atmospheric chemistry box model CAABA/MECCA-3.0, *Geosci. Model Dev.*, 4, 373–380, doi:10.5194/gmd-4-373-2011, 2011.

Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 2. C-1 through C-30 organic compounds from medium duty diesel trucks, *Environ. Sci. Technol.*, 33, 1578–1587, 1999.

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 3. C-1-C-29 organic compounds from fireplace combustion of wood, *Environ. Sci. Technol.*, 35, 1716–1728, 2001.
- Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 5. C-1-C-32 organic compounds from gasoline-powered motor vehicles, *Environ. Sci. Technol.*, 36, 1169–1180, 2002.
- Schwartz, R. E., Russell, L. M., Sjostedt, S. J., Vlasenko, A., Slowik, J. G., Abbatt, J. P. D., Macdonald, A. M., Li, S. M., Liggio, J., Toom-Sauntry, D., and Leaitch, W. R.: Biogenic oxidized organic functional groups in aerosol particles from a mountain forest site and their similarities to laboratory chamber products, *Atmos. Chem. Phys.*, 10, 5075–5088, doi:10.5194/acp-10-5075-2010, 2010.
- Shrivastava, M., Fast, J., Easter, R., Gustafson Jr., W. I., Zaveri, R. A., Jimenez, J. L., Saide, P., and Hodzic, A.: Modeling organic aerosols in a megacity: comparison of simple and complex representations of the volatility basis set approach, *Atmos. Chem. Phys.*, 11, 6639–6662, doi:10.5194/acp-11-6639-2011, 2011.
- Shrivastava, M. K., Lane, T. E., Donahue, N. M., Pandis, S. N., and Robinson, A. L.: Effects of gas particle partitioning and aging of primary emissions on urban and regional organic aerosol concentrations, *J. Geophys. Res.-Atmos.*, 113, D18301, doi:10.1029/2007jd009735, 2008.
- Stanier, C. O., Donahue, N. M., and Pandis, S. N.: Parameterization of secondary organic aerosol mass fractions from smog chamber data, *Atmos. Environ.*, 42, 2276–2299, 2008.
- Strader, R., Lurmann, F., and Pandis, S. N.: Evaluation of secondary organic aerosol formation in winter, *Atmos. Environ.*, 33, 4849–4863, 1999.
- Takahama, S., Schwartz, R. E., Russell, L. M., Macdonald, A. M., Sharma, S., and Leaitch, W. R.: Organic functional groups in aerosol particles from burning and non-burning forest emissions at a high-elevation mountain site, *Atmos. Chem. Phys.*, 11, 6367–6386, doi:10.5194/acp-11-6367-2011, 2011.
- Tost, H., Jöckel, P., Kerkweg, A., Sander, R., and Lelieveld, J.: Technical note: A new comprehensive SCAVenging submodel for global atmospheric chemistry modelling, *Atmos. Chem. Phys.*, 6, 565–574, doi:10.5194/acp-6-565-2006, 2006.
- Tsimpidi, A. P., Karydis, V. A., and Pandis, S. N.: Response of fine particulate matter to emission changes of oxides of nitrogen and anthropogenic volatile organic compounds in the Eastern United States, *J. Air Waste Manage.*, 58, 1463–1473, 2008.

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
	Printer-friendly Version	
	Interactive Discussion	



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

Tsimpidi, A. P., Karydis, V. A., Zavala, M., Lei, W., Molina, L., Ulbrich, I. M., Jimenez, J. L., and Pandis, S. N.: Evaluation of the volatility basis-set approach for the simulation of organic aerosol formation in the Mexico City metropolitan area, *Atmos. Chem. Phys.*, 10, 525–546, doi:10.5194/acp-10-525-2010, 2010.

Tsimpidi, A. P., Karydis, V. A., Zavala, M., Lei, W., Bei, N., Molina, L., and Pandis, S. N.: Sources and production of organic aerosol in Mexico City: insights from the combination of a chemical transport model (PMCAMx-2008) and measurements during MILAGRO, *Atmos. Chem. Phys.*, 11, 5153–5168, doi:10.5194/acp-11-5153-2011, 2011.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Zhang, Q., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., and Jimenez, J. L.: Time- and size-resolved chemical composition of submicron particles in Pittsburgh: implications for aerosol sources and processes, *J. Geophys. Res.-Atmos.*, 110, D07s09, doi:10.1029/2004jd004649, 2005.

Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, L13801, doi:10.1029/2007gl029979, 2007.

Zhang, Q. J., Beekmann, M., Drewnick, F., Freutel, F., Schneider, J., Crippa, M., Prevot, A. S. H., Baltensperger, U., Poulain, L., Wiedensohler, A., Sciare, J., Gros, V., Borbon, A., Colomb, A., Michoud, V., Doussin, J.-F., Denier van der Gon, H. A. C., Haefelin, M., Dupont, J.-C., Siour, G., Petetin, H., Bessagnet, B., Pandis, S. N., Hodzic, A., Sanchez, O., Honoré, C., and Perrussel, O.: Formation of organic aerosol in the Paris region during the MEGAPOLI summer campaign: evaluation of the volatility-basis-set approach within the CHIMERE model, *Atmos. Chem. Phys.*, 13, 5767–5790, doi:10.5194/acp-13-5767-2013, 2013.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Table 1. Organic component name classifications in the ORACLE module.

Term	Description
Source root name	
Modifiers	
a	Mass from anthropogenic sources (i.e. aSOA)
b	Mass from biogenic sources (i.e. bSOA)
f	Mass from fossil and biofuel combustion (i.e. fPOA)
bb	Mass from biomass burning processes (i.e. bbPOA)
Base Terms	
POA	Primary organic aerosol. This is emitted in the particle phase and has not undergone chemical reaction
POG	Primary organic gas that has not undergone chemical reaction
SOA	Secondary organic aerosol formed from the oxidation of gas-phase organic species
SOG	Secondary organic gas. The gas-phase mass produced by at least one chemical reaction in the atmosphere
Initial Volatility	
Suffix	
-sv	Product of the oxidation of SVOCs
-i	Product of the oxidation of IVOCs
-v	Product of the oxidation of VOCs

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

Table 2. Description of ORACLE organic compounds in gas and aerosol phases.

Gas-phase compounds	Particle phase compounds	Description
fPOG	fPOA	Primary organic compounds from fuel combustion
bbPOG	bbPOA	Primary organic compounds from biomass burning
fSOG-sv	fSOA-sv	Secondary organic compounds from the oxidation of fuel combustion SVOCs
bbSOG-sv	bbSOA-sv	Secondary organic compounds from the oxidation of biomass burning SVOCs
fSOG-iv	fSOA-iv	Secondary organic compounds from the oxidation of fuel combustion IVOCs
bbSOG-iv	bbSOA-iv	Secondary organic compounds from the oxidation of biomass burning IVOCs
aSOG-v	aSOA-v	Secondary organic compounds from the oxidation of anthropogenic VOCs
bSOG-v	bSOA-v	Secondary organic compounds from the oxidation of biogenic VOCs

Table 3. Global annual emissions of the lumped anthropogenic VOC species introduced in the ORACLE module.

VOC lumped species	RCP4.5 species	Emission rates (Tg yr ⁻¹)
ARO1	Benzene, Toluene	18.3
ARO2	Trimethyl Benzenes, Xylene, other Aromatics	13.9
ALK4	Pentanes	15.1
ALK5	Hexanes, other higher Alkenes	21.7
OLE1	Propene	7.4
OLE2	Other Alkenes	8.2

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

Table 4. Emission factors of the primary organic compounds from fuel combustion and biomass burning sources for each volatility bin used in ORACLE. The total global annual emission rates are also shown. The SVOC/IVOC emission rates are estimated by multiplying the emission factors of each bin with the non-volatile emissions rates.

	Volatility distribution				Non volatile POA emission rates Tg yr ⁻¹	SVOC/IVOC emission rates Tg yr ⁻¹
C^* ($\mu\text{g m}^{-3}$) at 298 K	10^{-1}	10^1	10^3	10^5		
Fuel combustion OA	fPOA1	fPOA2	fPOG3	fPOG4	12.3	30.8
Biomass burning OA	bPOA1	bbPOA2	bbPOG3	bbPOG4	24.8	62
Emission factors	0.18	0.32	0.5	1.5		

VOC lumped Species	$C^* \text{ in } \mu\text{g m}^{-3} \text{ at } 298 \text{ K}$			
	1	10	10^2	10^3
ARO1	0.003	0.165	0.300	0.435
ARO2	0.002	0.195	0.300	0.435
ALK4	0.000	0.038	0.000	0.000
ALK5	0.000	0.150	0.000	0.000
OLE1	0.001	0.005	0.038	0.150
OLE2	0.003	0.026	0.083	0.270
ISOP	0.009	0.030	0.015	0.000
TERP	0.107	0.092	0.359	0.600

^a The SOA yields are based on an assumed particle density of 1.5 g cm^{-3} .

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

Table 6. Statistical evaluation of monthly averaged predicted OA against IMPROVE, EMEP and East Asia observations during 2005–2009.

Network	Number of stations	Mean Observed ($\mu\text{g m}^{-3}$)	Mean Predicted ($\mu\text{g m}^{-3}$)	MAGE ($\mu\text{g m}^{-3}$)	MB ($\mu\text{g m}^{-3}$)	NME (%)	NMB (%)	RMSE ($\mu\text{g m}^{-3}$)
IMPROVE	193 ^a	1.93	3.02	1.71	1.09	89	57	2.6
EMEP	11 ^b	2.87	2.24	2.02	-0.63	70	-22	4.15
East Asia	18 ^c	11.02	7.09	4.76	-3.93	43	-36	5.53

^a 10 202 measurements^b 361 measurements^c 74 measurements.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)





ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|](#)[|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.

**Figure 1.** Diagram showing ORACLE as part of the EMAC chemistry-climate model.

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.




Figure 2. Schematic of the VBS resolution and the formation procedure of SOA from SVOC, IVOC and VOC emissions implemented in the ORACLE submodel. Red indicates that the organic compound is in the vapor phase and blue that it is in the particulate phase. The triangles indicate the formation of secondary organic aerosol (SOA) from semivolatiles by fuel combustion and biomass burning sources, while the squares show SOA from intermediate-volatiles by fuel combustion and biomass burning sources, and the rhombi the formation of SOA from anthropogenic and biogenic VOC sources. The partitioning processes, the aging reactions of the organic compounds, and the different names of the species used to track all compounds are also shown.

* For this application it is assumed that SOA formed by biogenic VOCs does not participate in aging reactions (Lane et al., 2008; Tsimpidi et al., 2010).

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.




Figure 3. Predicted average surface concentration of total OA ($\mu\text{g m}^{-3}$) during the years 2005–2009.

Title Page

Abstract

Introduction

Conclusio

References

Tables

Figures



Back

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion






Figure 4. Scatter plot comparing model predictions of total OA concentration (in $\mu\text{g m}^{-3}$) against observations from the (a) IMPROVE network in the United States, (b) EMEP network in Europe and East Asian sites from 2005 to 2009. Each point represents a monthly average value. Also shown the 1 : 1, 2 : 1 and 1 : 2 lines.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

ORACLE: an organic aerosol module

A. P. Tsimpidi et al.




Figure 5. Predicted average surface concentrations (in $\mu\text{g m}^{-3}$) of (a) POA from fuel combustion (fPOA) and (b) POA from biomass burning emissions (bbPOA) during the years 2005–2009.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Discussion Paper | ORACLE: an organic aerosol module

A. P. Tsimpidi et al.




Figure 6. Predicted average surface concentrations (in $\mu\text{g m}^{-3}$) of (a) SOA-sv from the oxidation of SVOCs from fuel combustion (fSOA-sv) and (b) SOA-sv from the oxidation SVOCs from biomass burning emissions (bbSOA-sv) during the years 2005–2009.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.




Figure 7. Predicted average surface concentrations (in $\mu\text{g m}^{-3}$) of (a) SOA-iv from the oxidation of IVOCs from fuel combustion (fSOA-iv) and (b) SOA-iv from the oxidation of IVOCs from biomass burning emissions (bbSOA-iv) during the years 2005–2009.




Figure 8. Predicted average surface concentrations (in $\mu\text{g m}^{-3}$) of (a) SOA-v from the oxidation of biogenic VOC emissions (bSOA-v) and (b) SOA-v from the oxidation of anthropogenic VOC emissions (aSOA-v) during the years 2005–2009.

[Title Page](#) [Abstract](#) [Introduction](#) [Conclusions](#) [References](#) [Tables](#) [Figures](#) [◀](#) [▶](#) [◀](#) [▶](#) [Back](#) [Close](#) [Full Screen / Esc](#) [Printer-friendly Version](#) [Interactive Discussion](#)



ORACLE: an organic aerosol module

A. P. Tsimpidi et al.




Figure 9. Predicted average zonal concentrations (in $\mu\text{g m}^{-3}$) of (a) POA (sum of fPOA and bbPOA) and (b) SOA (sum of fSOA-sv, bbSOA-sv, fSOA-iv, bbSOA-iv, aSOA-v, and bSOA-v) during the years 2005–2009.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

