Supporting Information to:

Noniosides E – H, New Trisaccharide Fatty Acid Esters from the Fruit of *Morinda citrifolia* (Noni)

Petur W. Dalsgaard¹

Olivier Potterat¹

Fiona Dieterle¹

Thomas Paululat²

Till Kühn³

Matthias Hamburger¹

Affiliation

¹ Institute of Pharmaceutical Biology, University of Basel, Basel, Switzerland,

² Organische Chemie II, University of Siegen, Siegen, Germany

³ Bruker BioSpin AG, NMR Division, Fällanden, Switzerland

Correspondence

Prof. Dr. Matthias Hamburger

Institute of Pharmaceutical Biology

Department of Pharmaceutical Sciences

University of Basel

Klingelbergstrasse 50

4056 Basel

Switzerland

Phone: +41-61-267-1425

Fax: +41-61-267-1474

E-mail: matthias.hamburger@unibas.ch

Fig. 1S Nonioside E (4).

Fig. 2S Nonioside F (5).

© Georg Thieme Verlag KG · DOI 10.1055/s-2006-951706 (10.1055/s-2006-951734) · Planta Med 2006; 72: e342–e346 · Dalsgaard PW et al.

Fig 4S Nonioside H (7). Compound is either 2-*O*-(6-*O*-hexanoyl- β -D-glucopyranosyl)-6-*O*-(β -D-glucopyranosyl)-1-*O*-octanoyl- β -D-glucopyranose or 2-*O*-(6-*O*-octanoyl- β -D-glucopyranosyl)-6-*O*-(β -D-glucopyranosyl)-1-*O*-hexanoyl- β -D-glucopyranose.