

**Supplementary Table 1.** Primers used for generation of truncated PSPC1 and NONO

		PSPC1	NONO
PN1	Forward*	Primer 1 CTGGTGCCACGCGTTCTATGGGGTTCCTACTATCGACATC Primer 2 AAGGATCCAGACTACAAGGACGACGATGACAAAGTCTGGTGCCACGCGGTTCT	CCCATATGCGTCTTTTTGTGGGAAATC
	Reverse	CCGAATTCTTAAAACCTGCTCCATGGGTTCCAC	CCGCTCGAGTTATCTCATTAGCATGACCTGGTG
PN2	Forward	CAGGATCCAGAAAACCTGTATTTTCAGGGCATGGGGTTCCTACTATCGACATC	CGCATATGGAAGGCTTGACTATTGAC
	Reverse	CCGAATTCTTAAAACCTGCTCCATGGGTTCCAC	AACTCGAGTTATTGGCGCCTCATCAAATC
PN3	Forward	CAGGATCCAGAAAACCTGTATTTTCAGGGCATGGGGTTCCTACTATCGACATC	CGCATATGGAAGGCTTGACTATTGAC
	Reverse	CGGAATTCTTACTTCCATCGAGATGCATAC	CCGCTCGAGTTACTTCCAGCGCATGGCATATTC
PN4	Forward	CAGGATCCAGAAAACCTGTATTTTCAGGGCATGGGGTTCCTACTATCGACATC	CGCATATGGAAGGCTTGACTATTGAC
	Reverse	CGGAATTCTTAGGCTTCTCTGATGTTTC	CCGCTCGAGTTAAGCCTCCTTGATGTTGCG
PN5	Forward	CAGGATCCAGAAAACCTGTATTTTCAGGGCATGGGGTTCCTACTATCGACATC	CGCATATGGAAGGCTTGACTATTGAC
	Reverse	CGGAATTCTTACATTAGCATTAAATTGGTG	CCGCTCGAGTTACATTAGCATGACCTGGTG
PN6	Forward	CAGGATCCAGAAAACCTGTATTTTCAGGGCATGGGGTTCCTACTATCGACATC	CGCATATGGAAGGCTTGACTATTGAC
	Reverse	CGGAATTCTTATTGGTTTCTGAGTTCTTC	CCGCTCGAGTTATTGGTTGTGCAGCTCTTC

\* Two PCRs were performed to generate the PSPC1 construct of PN1; in the first reaction, the forward primer 1 and the reverse primer were used. The product of this reaction was used as the template of the subsequent PCR using the forward primer 2 and the same reverse primer to generate the final PCR product for cloning into pETDuet-1.