SUPPLEMENTARY MATERIAL

Predicting XAFS scattering path cumulants and XAFS spectra for metals (Cu, Ni, Fe, Ti, Au) using molecular dynamics simulations

M.A. Karolewski^{1,*}, R.G. Cavell¹, R.A. Gordon², C.J. Glover³, M. Cheah³, and M.C. Ridgway⁴.

Contents

Figure S1. Third cumulants (C_3) for several XAFS scattering paths predicted by MD simulations for Ni at 300 K.

Figure S2. Third cumulants (C_3) for several XAFS scattering paths predicted by MD simulations for Fe at 300 K.

Figure S3. Third cumulants (C_3) for several XAFS scattering paths predicted by MD simulations for Ti at 300 K.

Figure S4. Experimental XAFS spectrum for Ni, and XAFS spectra fitted using scaled MD cumulants.

Figure S5. Experimental XAFS spectrum for Fe, and XAFS spectra fitted using scaled MD cumulants.

Figure S6. Experimental XAFS spectrum for Ti, and XAFS spectra fitted using scaled MD cumulants.

Figure S7. Experimental XAFS spectrum for Au, and XAFS spectra fitted using scaled MD cumulants.

¹ Department of Chemistry, University of Alberta, Edmonton, Alberta T6G 2G2, Canada.

² Pacific Northwest Consortium Synchrotron Radiation Facility, APS Sector 20, 9700 S. Cass Ave. Bldg. 435E, Argonne, IL, USA 60439.

³ Australian Synchrotron, Clayton, VIC 3168, Australia.

⁴ Department of Electronic Materials Engineering, Research School of Physics and Engineering, Australian National University, ACT 0200, Australia.

^{*} Corresponding author. E-mail address: <u>karolewski@alum.mit.edu</u>.




Figure S1. Third cumulants (C_3) for several XAFS scattering paths predicted by MD simulations for Ni at 300 K. Data for multiple-scattering paths are indicated by arrows.




Figure S2. Third cumulants (C_3) for several XAFS scattering paths predicted by MD simulations for Fe at 300 K. Data for multiple-scattering paths are indicated by arrows.




Figure S3. Third cumulants (C_3) for several XAFS scattering paths predicted by MD simulations for Ti at 300 K. Data for multiple-scattering paths are indicated by arrows.




Figure S4. Symbols: experimental XAFS spectrum for Ni; solid lines: XAFS spectra fitted using scaled MD cumulants (solid lines; see main text for fitting details).




Figure S5. Symbols: experimental XAFS spectrum for Fe; solid lines: XAFS spectra fitted using scaled MD cumulants (solid lines; see main text for fitting details).




Figure S6. Symbols: experimental XAFS spectrum for Ti; solid lines: XAFS spectra fitted using scaled MD cumulants (solid lines; see main text for fitting details).




Figure S7. Symbols: experimental XAFS spectrum for Au; solid lines: XAFS spectra fitted using scaled MD cumulants (solid lines; see main text for fitting details).