



STRUCTURAL
CHEMISTRY

Volume 74 (2018)

Supporting information for article:

Purification, characterization, crystal structure and NO production inhibitory activity of three new sesquiterpenoids from *homalomena occulta*

Qi Zhang, Li Ma, Zhaoxia Qu, Guige Hou, Yanan Wang, Chunhua Wang and Feng Zhao




Figure S1 The ¹H NMR Spectrum of Compound **1** in MeOH-*d*₄ (600 MHz) (a); The ¹³C NMR Spectrum of Compound **1** in MeOH-*d*₄ (150 MHz) (b); The HSQC Spectrum of Compound **1** in MeOH-*d*₄ (600 MHz for ¹H) (c); The HMBC Spectrum of Compound **1** in MeOH-*d*₄ (600 MHz for ¹H) (d); The ROESY Spectrum of Compound **1** in MeOH-*d*₄ (600 MHz) (e).




Figure S2 The ^1H NMR Spectrum of Compound (2) in acetone- d_6 (500 MHz) (a); The ^{13}C NMR Spectrum of Compound (2) in acetone- d_6 (125 MHz) (b); The ^1H - ^1H COSY Spectrum of Compound (2) in acetone- d_6 (500 MHz) (c); The HSQC Spectrum of Compound (2) in acetone- d_6 (500 MHz for ^1H) (d); The HMBC Spectrum of Compound (2) in acetone- d_6 (500 MHz for ^1H) (e); The NOESY Spectrum of Compound (2) in acetone- d_6 (500 MHz) (f).




Figure S3 The ^1H NMR Spectrum of Compound (3) in acetone- d_6 (500 MHz) (a); The ^{13}C NMR Spectrum of Compound (3) in acetone- d_6 (125 MHz) (b); The ^1H - ^1H COSY Spectrum of Compound (3) in acetone- d_6 (500 MHz) (c); The HSQC Spectrum of Compound (3) in acetone- d_6 (500 MHz for ^1H) (d); The HMBC Spectrum of Compound (3) in acetone- d_6 (500 MHz for ^1H) (e); The NOESY Spectrum of Compound (3) in acetone- d_6 (500 MHz) (f).