



STRUCTURAL BIOLOGY  
COMMUNICATIONS

**Volume 71 (2015)**

**Supporting information for article:**

**Stabilization of porcine pancreatic elastase (PPE) crystals by  
glutaraldehyde cross-linking**

**Stefan Hofbauer, José A. Brito, Jalmira Mulchande, Przemyslaw Nogly, Miguel  
Pessanha, Rui Moreira and Margarida Archer**

**Table S1** Selected crystallization conditions of PPE and PPE-inhibitor complexes.

Condition	pH	Reference
0.2 - 0.3 M sodium sulfate 0.1 M sodium acetate	5.1	(Weiss <i>et al.</i> , 2002, Würtele <i>et al.</i> , 2000, Oliveira <i>et al.</i> , 2007)
0.1 M sodium acetate 0.05 M sodium citrate 0.005 M calcium chloride	5.1	(Gilliland <i>et al.</i> , 1994, Weiss <i>et al.</i> , 2002)
0.2 M ammonium sulfate 0.1 M sodium MES 30% (w/v) PEG 5000	6.5	(Hilpert <i>et al.</i> , 2004)
0.3 M sodium chloride 0.05 M tris(hydroxymethyl)aminomethane	7.0	(Kinoshita <i>et al.</i> , 2006)
68 - 74% (w/v) 2-methyl-2,4-pentanediol 0.01 M sodium phosphate	5.9	(Tsunemi <i>et al.</i> , 1996)
0.2 M bicine 0.06 M sodium citrate	8.1	(Dementiev <i>et al.</i> , 2006)