

Supporting Information
for
Crystal Structure of the Homoleptic Zinc(II) Complex Based on *bis*(3,5-Diisopropylpyrazol-1-yl)acetate

Josiah G. Elsberg, Nicholas G. Spiropulos, Adam C. Colson, Eric C. Brown*

†Department of Chemistry, Boise State University, Boise, ID 83725

Tel: 208-426-1186

Fax: 208-426-1311

E-mail: ericbrown3@boisestate.edu

*To whom all correspondence should be addressed

Contents

Figure S1. ¹ H NMR spectrum of 2	S2
Figure S2. ¹³ C { ¹ H} NMR spectrum of 2	S2
Figure S3. HMBC NMR spectrum of 2	S3
Figure S4. HSQC NMR spectrum of 2	S3
Figure S5 COSY NMR spectrum of 2	S4
Figure S6. ESI Mass spectrometry of 2	S4




Figure S1. ^1H NMR spectrum of **2** in CDCl_3 .




Figure S2. ^{13}C NMR spectrum of **2** in CDCl_3 .




Figure S3. HMBC spectrum of **2** in CDCl₃.




Figure S4. HSQC spectrum of **2** in CDCl₃. Expansion shown in bottom right.




Figure S5. COSY NMR spectrum of **2** in CDCl_3 .




Figure S6. Experimental (left) and theoretical (right) ESI-MS spectra of **2**.