1 **APPENDIX 1: SUPPLEMENTARY DATA**

2 **Data collection methodology**

3 Images of published seismic-reflection and outcrop cross-sections were geo-
4 rectified, allowing for trajectory and channel form measurements using ArcGIS.
5 Previously unpublished measurements were taken directly from 3D seismic volumes
6 using Petrel. Data were compiled, stored and analyzed using Matlab. We collected
7 trajectory and channel form data according to simple guidelines based on observable
8 seismic/outcrop facies and geometries. Trajectory data (L_c , L_a) for a particular channel
9 belt were collected at the basal terminations of inclined reflectors/surfaces (Fig 1B). In
10 areas of high aggradation, points are located at the center of concave-up high-amplitude,
11 laterally discontinuous reflectors interpreted as channel thalweg deposits (Fig. 1B).
12 Because seismic resolution is variable, these points record the most reliable
13 spatiotemporal trajectory, not the specific number of discrete channel migration events
14 (cf. Maier et al., 2012). Data quality varied due to limited seismic resolution, orientation
15 of cross section relative to channel, and image quality. No de-compaction of the
16 trajectory or channel form measurements was attempted. We also note that it is difficult
17 to map points in phases of channel evolution characterized by significant lateral
18 migration and/or vertical degradation due to low preservation potential. Therefore,
19 mapped channel belt widths are minimum values as they preferentially record
20 aggradational phases of channel evolution. This also may also create a bias towards

aggradational trajectories in our dataset, but it exists for both fluvial and submarine examples.

Channel dimensions

Fig. DR1 shows the cumulative distribution of aspect ratios (B/H) of submarine and fluvial channels from modern and ancient settings. This compiled dataset shows quite similar channel aspect ratios for submarine and fluvial systems, validating our findings that channel kinematics rather than the channel form are responsible for differences in channel-belt shape and stratigraphic architecture. However, slight differences between these compiled data are evident. For example, the variability between distributions of submarine channel aspect ratios (Fig. DR1) likely results from the fact that defining bankfull width and depth in submarine channels is problematic (see discussion in Pirmez and Imran, 2003). Fluvial channel aspect ratios collected from this study are very similar to data compiled by Wilkerson and Parker (2011) (Fig. DR1). However, data from Jerolmack and Mohrig (2007) (Fig. DR1) have much larger values of B/H , resulting in consistently smaller mobility numbers as compared to this study (Fig. 3B, C). Whereas Jerolmack and Mohrig (2007) presented many fluvial systems with $M < 1$, our study calculated very few fluvial values of $M_{si} < 1$ and no values of $M_{sb} < 1$ (Fig. 3B, C). Interestingly, the condition for branching found by Jerolmack and Mohrig (2007) occurred at $M \sim 1$, and the majority of the very large values of B/H are from these distributary systems (Fig. DR1; Fig. 2 of Jerolmack and Mohrig, 2007).

FIGURE AND TABLE CAPTIONS

Fig. DR1. Cumulative distribution of channel aspect ratios (B/H) from modern and ancient data.

Table DR1. Data sources and citations used in this study.

REFERENCES CITED (Supplementary text)

- Adams, M.M., and Bhattacharya, J.P., 2005, No Change in Fluvial Style Across a Sequence Boundary, Cretaceous Blackhawk and Castlegate Formations of Central Utah, U.S.A.: *Journal of Sedimentary Research*, v. 75, p. 1038–1051, doi: 10.2110/jsr.2005.080.
- Alqahtani, F.A., Johnson, H.D., Jackson, C.A.L., and Som, M.R.B., 2015, Nature, origin and evolution of a Late Pleistocene incised valley-fill, Sunda Shelf, Southeast Asia: *Sedimentology*, v. 62, p. 1198–1232, doi: 10.1111/sed.12185.
- Armstrong, C. P., 2012, 3D seismic geomorphology and stratigraphy of the late Miocene to Pliocene Mississippi River Delta: fluvial systems and dynamics. MS Thesis, University of Texas at Austin, 119 p.
- Auchter, N., 2012, Evolution and Architecture of an Incised Valley in the Upper Cretaceous Eagle Formation in South-Central Montana. MS thesis, University of Montana, 132 p.

66 Auchter, N.C., Romans, B.W., Hubbard, S.M., Stright, L., Reimchen, A.P., and Daniels,
67 B.G., 2014, Slope evolution revealed by analysis of sandstone body architecture,
68 Tres Pasos Formation at Cerro Mirador, Chile. AAPG Annual Meeting, Houston.

69 Babonneau, N., Savoye, B., Cremer, M., and Klein, B., 2002, Morphology and
70 architecture of the present canyon and channel system of the Zaire deep-sea fan:
71 Marine and Petroleum Geology, v. 19, p. 445–467.

72 Babonneau, N., Savoye, B., Cremer, M., and Bez, M., 2004, Multiple terraces within the
73 deep incised Zaire Valley (ZaiAngo Project): are they confined levees?:
74 Geological Society, London, Special Publications, v. 222, p. 91–114, doi:
75 10.1144/GSL.SP.2004.222.01.06.

76 Babonneau, N., Savoye, B., Cremer, M., and Bez, M., 2010, Sedimentary Architecture in
77 Meanders of a Submarine Channel: Detailed Study of the Present Congo
78 Turbidite Channel (Zaiango Project): Journal of Sedimentary Research, v. 80, p.
79 852–866, doi: 10.2110/jsr.2010.078.

80 Darmadi, Y., Willis, B.J., and Dorobek, S.L., 2007, Three-Dimensional Seismic
81 Architecture of Fluvial Sequences on the Low-Gradient Sunda Shelf, Offshore
82 Indonesia: Journal of Sedimentary Research, v. 77, p. 225–238, doi:
83 10.2110/jsr.2007.024.

84 Deptuck, M.E., Steffens, G.S., Barton, M., and Pirmez, C., 2003, Architecture and
85 evolution of upper fan channel-belts on the Niger Delta slope and in the Arabian
86 Sea: Marine and Petroleum Geology, v. 20, p. 649–676, doi:
87 10.1016/j.marpetgeo.2003.01.004.

88 Deptuck, M.E., Sylvester, Z., Pirmez, C., and O'Byrne, C., 2007, Migration–aggradation
89 history and 3-D seismic geomorphology of submarine channels in the Pleistocene
90 Benin-major Canyon, western Niger Delta slope: *Marine and Petroleum Geology*,
91 v. 24, p. 406–433, doi: 10.1016/j.marpetgeo.2007.01.005.

92 Di Celma, C.N., Brunt, R.L., Hodgson, D.M., Flint, S.S., and Kavanagh, J.P., 2011,
93 Spatial and Temporal Evolution of a Permian Submarine Slope Channel-Levee
94 System, Karoo Basin, South Africa: *Journal of Sedimentary Research*, v. 81, p.
95 579–599, doi: 10.2110/jsr.2011.49.

96 Droz, L., Rigaut, F., Cochonat, P., and Tofani, R., 1996, Morphology and recent
97 evolution of the Zaire turbidite system (Gulf of Guinea): *Geological Society of*
98 *America Bulletin*, v. 108, p. 253–269, doi: 10.1130/0016-
99 7606(1996)108<0253:MAREOT>2.3.CO;2.

100 Droz, L., Marsset, T., Ondras, H., and Lopez, M., 2003, Architecture of an active mud-
101 rich turbidite system: The Zaire Fan (Congo–Angola margin southeast Atlantic):
102 Results from ZaAngo 1 and 2 cruises: *AAPG Bulletin*, v. 87, p. 1145–1168.

103 Flood, R.D., and Damuth, J.E., 1987, Quantitative characteristics of sinuous distributary
104 channels on the Amazon Deep-Sea Fan: *Geological Society of America Bulletin*,
105 v. 98, p. 728–738, doi: 10.1130/0016-7606(1987)98<728:QCOSDC>2.0.CO;2.

106 Fustic, M., Hubbard, S.M., Spencer, R., Smith, D.G., Leckie, D.A., Bennett, B., and
107 Larter, S., 2012, Recognition of down-valley translation in tidally influenced
108 meandering fluvial deposits, Athabasca Oil Sands (Cretaceous), Alberta, Canada:
109 *Marine and Petroleum Geology*, v. 29, p. 219–232, doi:
110 10.1016/j.marpetgeo.2011.08.004.

111 Ghinassi, M., Nemec, W., Aldinucci, M., Nehyba, S., Özaksoy, V., and Fidolini, F.,
112 2014, Plan-form evolution of ancient meandering rivers reconstructed from
113 longitudinal outcrop sections (N. Mountney, Ed.): *Sedimentology*, v. 61, p. 952–
114 977, doi: 10.1111/sed.12081.

115 Hartley, A.J., Owen, A., Swan, A., Weissmann, G.S., Holzweber, B.I., Howell, J.,
116 Nichols, G., and Scuderi, L., 2015, Recognition and importance of amalgamated
117 sandy meander belts in the continental rock record: *Geology*, v. 43, p. 679–682,
118 doi: 10.1130/G36743.1.

119 Jegou, Isabelle, 2008, Étude de la transition chenal-levées-lobe dans les systèmes
120 turbiditiques récents. Ph.D. thesis, Université de Brest, 351 p.

121 Jobe, Z.R., Lowe, D.R., and Uchytíl, S.J., 2011, Two fundamentally different types of
122 submarine canyons along the continental margin of Equatorial Guinea: *Marine*
123 *and Petroleum Geology*, v. 28, p. 843–860, doi:
124 10.1016/j.marpetgeo.2010.07.012.

125 Jobe, Z.R., Sylvester, Z., Parker, A.O., Howes, N., Slowey, N., and Pirmez, C., 2015,
126 Rapid Adjustment of Submarine Channel Architecture To Changes In Sediment
127 Supply: *Journal of Sedimentary Research*, v. 85, p. 729–753, doi:
128 10.2110/jsr.2015.30.

129 Kolla, V., Bandyopadhyay, A., Gupta, P., Mukherjee, B., and Ramana, D.V., 2012,
130 Morphology and internal structure of a recent upper Bengal fan-valley complex,
131 in Prather, B.E., Deptuck, M.E., Mohrig, D., van Hoorn, B., and Wynn, R., eds.,
132 *Application of the Principles of Seismic Geomorphology to Continental-Slope*

133 and Base-of-Slope Systems: Case Studies from Seafloor and Near-Seafloor
134 Analogues: SEPM, Special Publication 99, p. 347–369.

135 Kolla, V., Posamentier, H.W., and Wood, L.J., 2007, Deep-water and fluvial sinuous
136 channels—Characteristics, similarities and dissimilarities, and modes of
137 formation: *Marine and Petroleum Geology*, v. 24, p. 388–405, doi:
138 10.1016/j.marpetgeo.2007.01.007.

139 Kenyon, N.H., Amir, A., and Cramp, A., 1995, Geometry of the younger sediment bodies
140 on the Indus Fan, in Pickering, K.T., Hiscott, R.N., Kenyon, N.H., Ricci Lucchi,
141 F., and Smith, R.D.A., eds., *Atlas of Deepwater Environments; Architectural
142 Style in Turbidite Systems*: London, Chapman & Hall, p. 212–215.

143 Labourdette, R., and Jones, R.R., 2007, Characterization of fluvial architectural elements
144 using a three-dimensional outcrop data set: Escanilla braided system, South-
145 Central Pyrenees, Spain: *Geosphere*, v. 3, p. 422, doi: 10.1130/GES00087.1.

146 Macauley, R.V., and Hubbard, S.M., 2013, Slope channel sedimentary processes and
147 stratigraphic stacking, Cretaceous Tres Pasos Formation slope system, Chilean
148 Patagonia: *Marine and Petroleum Geology*, v. 41, p. 146–162, doi:
149 10.1016/j.marpetgeo.2012.02.004.

150 Maier, K.L., Fildani, A., McHargue, T.R., Paull, C.K., Graham, S.A., and Caress, D.W.,
151 2012, Punctuated Deep-Water Channel Migration: High-Resolution Subsurface
152 Data from the Lucia Chica Channel System, Offshore California, U.S.A: *Journal
153 of Sedimentary Research*, v. 82, p. 1–8, doi: 10.2110/jsr.2012.10.

154 Manson, S, 2009, Avulsion process: stratigraphic and lithologic records - Application to
155 the Amazon and Zaire turbidite systems. Universite de Bretagne occidentale -
156 Brest, 242 p.

157 Maynard, J.R., Feldman, H.R., and Alway, R., 2010, From Bars to Valleys: The
158 Sedimentology and Seismic Geomorphology of Fluvial to Estuarine Incised-
159 Valley Fills of the Grand Rapids Formation (Lower Cretaceous), Iron River Field,
160 Alberta, Canada: *Journal of Sedimentary Research*, v. 80, p. 611–638, doi:
161 10.2110/jsr.2010.060.

162 Miall, A.D., and Jones, B.G., 2003, Fluvial Architecture of the Hawkesbury Sandstone
163 (Triassic), Near Sydney, Australia: *Journal of Sedimentary Research*, v. 73, p.
164 531–545, doi: 10.1306/111502730531.

165 Musial, Geoffray, Richard Labourdette, Jessica Franco, and Jean-Yves Reynaud, 2013,
166 Modeling of a tide-influenced point-bar heterogeneity distribution and Impacts on
167 steam-assisted gravity drainage production: Example from Steepbank River,
168 McMurray Formation, Canada, in F. J. Hein, D. Leckie, S. Larter, and J. R. Suter,
169 eds., *Heavy-oil and oil-sand petroleum systems in Alberta and beyond: AAPG*
170 *Studies in Geology* 64, p. 545 – 564.

171 O’Byrne, C. J., Barton, M. D., Steffens, G. S., Pirmez, C., & Buergisser, H., 2007,
172 Architecture of a laterally migrating channel complex: Channel 4, Isaac
173 Formation, Windermere Supergroup, Castle Creek North, British Columbia,
174 Canada: in *Atlas of Deep-Water Outcrops*, American Association of Petroleum
175 Geologists, *Studies in Geology*, 56, 115-118.

176 Pickering, K.T., Hodgson, D.M., Platzman, E., Clark, J.D., and Stephens, C., 2001, A
 177 new type of bedform produced by backfilling processes in a submarine channel,
 178 late Miocene, Tabernas-Sorbas basin, SE Spain: *Journal of Sedimentary Research*,
 179 v. 71, p. 692–704.

180 Piper, D., Pirmez, C., Manley, P.L., Long, D., Flood, R.D., Normark, W.R., and Showers,
 181 W., 1997, Mass-transport deposits of the Amazon Fan, in *Proceedings of the*
 182 *Ocean Drilling Program. Scientific Results. Vol. 155*, p. 109–146.

183 Popescu, I., Lericolais, G., Panin, N., Wong, H.K., and Droz, L., 2001, Late Quaternary
 184 channel avulsions on the Danube deep-sea fan, Black Sea: *Marine Geology*, v.
 185 179, p. 25–37, doi: 10.1016/S0025-3227(01)00197-9.

186 Pranter, M.J., Ellison, A.I., Cole, R.D., and Patterson, P.E., 2007, Analysis and modeling
 187 of intermediate-scale reservoir heterogeneity based on a fluvial point-bar outcrop
 188 analog, Williams Fork Formation, Piceance Basin, Colorado: *AAPG Bulletin*, v.
 189 91, p. 1025–1051, doi: 10.1306/02010706102.

190 Reijenstein, H.M., Posamentier, H.W., and BHATTACHARYA, J.P., 2011, Seismic
 191 geomorphology and high-resolution seismic stratigraphy of inner-shelf fluvial,
 192 estuarine, deltaic, and marine sequences, Gulf of Thailand: *AAPG Bulletin*, v. 95,
 193 p. 1959–1990, doi: 10.1306/03151110134.

194 Reimchen, A.P., Hubbard, S.M., Auchter, N.C., Daniels, B., Jackson, A., Stright, L., and
 195 Romans, B.W., 2014, The record of sediment transfer on deep-water slopes:
 196 Characterization of an aggradational channel complex 25 km from the shelf edge,
 197 Tres Pasos Formation, Chile. AAPG Annual Meeting, Houston.

198 Tjallingii, R., Stattegger, K., Wetzel, A., and Van Phach, P., 2010, Infilling and flooding
199 of the Mekong River incised valley during deglacial sea-level rise: Quaternary
200 Science Reviews, v. 29, p. 1432–1444, doi: 10.1016/j.quascirev.2010.02.022.

201 Ullah, M.S., Bhattacharya, J.P., and Dupre, W.R., 2015, Confluence Scours Versus
202 Incised Valleys: Examples From the Cretaceous Ferron Notom Delta,
203 Southeastern Utah, U.S.A: Journal of Sedimentary Research, v. 85, p. 445–458,
204 doi: 10.2110/jsr.2015.34.

205 Wood, L.J., and Mize-Spansky, K.L., 2009, Quantitative seismic geomorphology of a
206 Quaternary leveed-channel system, offshore eastern Trinidad and Tobago,
207 northeastern South America: AAPG Bulletin, v. 93, p. 101–125, doi:
208 10.1306/08140807094.

209

210

Figure DR1

Table DR1. Data sources and citations used in this study.

System	Environment of Deposition	Data Type	Data Source	# of trajectories	Latitude	Longitude
Bengal Fan-Valley	Submarine	Seismic	Kolla et al. (2012) doi: 10.2110/pec.12.99.0347	69	18.42286	88.71923
Niger Delta Slope: Benin Major Canyon	Submarine	Seismic	Deptuck et al. (2007) doi:10.1016/j.marpetgeo.2007.01.005 Kolla et al (2007) doi:10.1016/j.marpetgeo.2007.01.007	56	5.24521	4.21526
Niger Delta Slope: Y Canyon	Submarine	Seismic	Jobe et al., 2015 doi:10.2110/jsr.2015.30 and this study	20	4.72031	4.46132
Niger Delta Slope: Bonga Canyon	Submarine	Seismic	Auchter, unpublished (this study)	15	4.47796	4.49091
Zaire/Congo submarine channel	Submarine	Seismic	Babonneau et al. (2002) doi:10.1016/S0264-8172(02)00009-0 Babonneau et al. (2004) doi: 10.1144/GSL.SP.2004.222.01.06 Babonneau et al. (2010) doi: 10.2110/jsr.2010.078	11	-5.65162	10.61062
Indus: Upper Fan Channels	Submarine	Seismic	Deptuck et al. (2003) doi:10.1016/j.marpetgeo.2003.01.004; Sylvester, unpublished data (this study)	24	23.24681	65.76993
Gulf of Mexico	Submarine	Seismic	Kolla et al. (2007) doi:10.1016/j.marpetgeo.2007.01.007	8	27.94598	-89.30778
Lucia Chica Channel System	Submarine	Seismic	Maier et al. (2012) doi: 10.2110/jsr.2012.10	12	35.74116	-121.81659
Amazon Fan Channels	Submarine	Seismic	Flood & Damuth (1987) doi: 10.1130/0016-7606(1987)98<728:QCOSDC>2.0.CO;2 Piper et al. (1997) doi:10.2973/odp.proc.sr.155.212.1997	9	5.63388	-47.71058
Borneo	Submarine	Seismic	Kolla et al. (2007) doi:10.1016/j.marpetgeo.2007.01.007	1	-0.64159	117.99938
Krishna Basin	Submarine	Seismic	Kolla et al. (2007) doi:10.1016/j.marpetgeo.2007.01.007	1	15.72219	81.62802
Equatorial Guinea slope	Submarine	Seismic	Jobe et al. (2011) doi:10.1016/j.marpetgeo.2010.07.012	9	1.68872	9.32117
Moki Formation, Taranaki, New Zealand	Submarine	Seismic	Shumaker and Jobe, unpublished (this study)	16	-39.44102	173.39802
Columbus Basin	Submarine	Seismic	Wood & Mize-Spansky (2009) doi:10.1306/08140807094	6	11.11256	-60.15098
Karoo Basin, South Africa	Submarine	Outcrop	Di Celma et al. (2011) doi: 10.2110/jsr.2011.49	4	-33.20047	20.78796
Tres Pasos Formation, Magallanes Basin, Chile	Submarine	Outcrop	Macauley & Hubbard (2013) doi:10.1016/j.marpetgeo.2012.02.004 Auchter et al. (2014) AAPG abstract Reimchen et al. (2014) AAPG abstract	9	-51.38507	-72.43274
Capistrano Formation, San Clemente, CA	Submarine	Outcrop	Jeanette and Bellian, unpublished (this study)	2	33.40343	-117.60507
Isaac Formation, Windermere, Canada	Submarine	Outcrop	O'Byrne et al. (2007) doi:10.1306/12401073St563290	1	53.05407	-120.44988
Ross Formation, Clare Basin, Ireland	Submarine	Outcrop	Jobe and Sylvester, unpublished data (this study)	1	52.59475	-9.85446
Solitary channel, Tabernas Basin, Spain	Submarine	Outcrop	Pickering et al. (2001) doi:10.1306/2DC40960-0E47-11D7-8643000102C1865D	1	37.01486	-2.46423
Tourelle Fm, Canada	Submarine	Outcrop	This study	1	49.15250	-66.43648
Natuna Basin, Gulf of Thailand	Fluvial	Seismic	Kolla et al. (2007) doi:10.1016/j.marpetgeo.2007.01.007	7	4.52001	106.60268
Grand Rapids Formation, Alberta	Fluvial	Seismic	Maynard et al. (2010) doi: 10.2110/jsr.2010.060	6	55.99832	-113.55601
Mekong River, Vietnam	Fluvial	Seismic	Tjallingii et al. (2010) doi:10.1016/j.quascirev.2010.02.022	1	9.51837	106.93548
Sunda Shelf, Gulf of Thailand	Fluvial	Seismic	Alqahtani et al. (2015) doi: 10.1111/sed.12185 Reijenstein et al. (2007) doi:10.1306/0315111013 Darmadi et al. (2007) doi: 10.2110/jsr.2007.024	7	9.05603	101.67400
McMurray Formation, Alberta, Canada	Fluvial	Outcrop	Fustic et al. (2012) doi:10.1016/j.marpetgeo.2011.08.004 Musial et al. (2012) doi:10.1016/j.sedgeo.2011.04.020	3	57.01236	-111.46643
Boyabat Basin, Turkey	Fluvial	Outcrop	Ghinassi et al. (2014) doi: 10.1111/sed.12081	4	41.57610	34.74013
Blackhawk Formation, Utah	Fluvial	Outcrop	Adams & Bhattacharya (2005) doi: 10.2110/jsr.2005.080	1	38.88048	-111.58324
Escanilla Formation, Ainsa Basin, Spain	Fluvial	Outcrop	Labourdette & Jones (2007) doi: 10.1130/GES00087.1	1	42.27190	0.12257
Hawkesbury Sandstone, Australia	Fluvial	Outcrop	Miall & Jones (2003) doi: 10.1306/111502730531	1	-34.00836	151.22861
Morrison Formation, Utah	Fluvial	Outcrop	Hartley et al. (2015) doi:10.1130/G36743.1	1	38.40499	-111.00964
Ferron Sandstone, Utah	Fluvial	Outcrop	Ullah et al. (2015) doi: http://dx.doi.org/10.2110/jsr.2015.34	3	38.40874	-110.85496
Williams Fork Formation, Colorado	Fluvial	Outcrop	Pranter et al. (2007) doi:10.1306/02010706102	3	39.13684	-108.36776
Eagle Formation, Montana	Fluvial	Outcrop	Auchter (2012) unpublished thesis	3	45.78476	-108.75089
Mississippi River (Pliocene)	Fluvial	Seismic	Armstrong (2012) unpublished thesis	1	29.70636	-89.70126