Comment and Reply
Dilational fault slip and pit chain formation on Mars, David A. Ferrill et al., *GSA Today*, v. 14, no. 10, p. 4–12.

Comment

Daniel R. Shawe, U.S. Geological Survey, Emeritus, 1805 S. Balsam no. 323, Lakewood, Colorado 80232, USA

I am greatly intrigued by the cover article in the October issue of *GSA Today* by Ferrill et al. (2004) that describes grabens and pit craters in the Alba Patera region of Mars. The thrust of the article is that north-trending grabens and pit craters on north-northeast–striking structures are the result of east-west dilation (extension). Block diagrams show the pit craters forming along graben-bounding faults; loose surficial material flowed into open space along the dilated graben faults. However, the images of the area (particularly the cover illustration) show clearly that the pit craters formed along the transverse, north-northeast–striking structures, not along the graben-bounding faults.

The question arises, therefore: What is the mechanism of pit crater formation? I suggest that the transverse structures are strike-slip faults. Although the article states that the faults in the Alba Patera show “no consistent evidence of strike-slip displacement” (p. 6) (and I agree that the graben fault traces show no clear indication of such), the orientations of the graben faults nevertheless have the relation of gash fractures along left-lateral strike-slip faults defined by the aligned pit craters, and lateral movement on such a structure could result in dilation of segments that are not aligned perfectly with the direction of slip.

Grabens are common structures associated with major strike-slip faults, as along the Alpine fault in New Zealand, the San Andreas in California, and the Great Basin of Nevada. Such an association on Mars would not be a surprise. I have interpreted development of grabens in western Nevada to be related to lateral movement on strike-slip faults (Shawe, 1965). Whatever the case, I think that the origin of the pit craters on Mars is more complex than the October 2004 *GSA Today* cover article infers.

REFERENCES CITED

- Ferrill, D.A., Wyrick, D.Y., Morris, A.P., Sims, D.W., and Franklin, N.M., 2004, Dilational fault slip and pit chain formation on Mars: *GSA Today*, v. 14, no. 10, p. 4–12.
- Shawe, D.R., 1965, Strike-slip control of Basin-Range structure indicated by historical faults in western Nevada: *Geological Society of America Bulletin*, v. 76, p. 1361–1378.

Reply

David A. Ferrill, CNWRA, Southwest Research Institute, 6220 Culebra Road, San Antonio, Texas 78238, USA, dferrill@swri.edu; **Alan P. Morris**, Department of Earth and Environmental Science, University of Texas, San Antonio, Texas 78249, USA; **Danielle Y. Wyrick**, **Darrell W. Sims**, and **Nathan M. Franklin**, CNWRA, Southwest Research Institute, 6220 Culebra Road, San Antonio, Texas 78238, USA

We appreciate Daniel Shawe's interest in our article (Ferrill et al., 2004). We agree that normal faults commonly form in association with strike-slip deformation (e.g., Ramsay and Huber, 1987). In addition, certain geometric features commonly associated with strike-slip deformation, such as en echelon fault arrangement, can form in normal fault stress regimes without strike-slip displacement (Ferrill et al., 1999).

As Shawe noted, the fault pattern in the cover illustration and Figure 3 of Ferrill et al. (2004) shows that pit chains are common along northeast-striking faults and crosscut north-striking faults. Faults that host pit chains have significant vertical components of displacement (e.g., 100–600 m throw, shown in Fig. 1A and in the cover illustration of Ferrill et al., 2004). Where drainage channels are cut by faults, they lack evidence of strike-slip deformation. These observations support our normal fault interpretation for the Alba Patera fault system. Rather than calling on strike-slip deformation, we interpret this fault pattern to reflect evolution of a regional stress field responding to a combination of volcano inflation and deflation and regional extension. This does not negate the possibility that these normal faults could be part of a larger scale strike-slip system. However, in a comprehensive review Schultz (1989) found only small scale strike-slip faults in the Coprates region southeast of the Valles Marineris. The paucity of evidence for strike-slip faulting on Mars, compared with Earth, may be attributable to the apparent lack of large-scale horizontal plate motion and the corresponding geometric requirement for transform plate-boundary displacement. Although we do not see evidence for strike-slip faulting at Alba Patera on Mars, there is active pit crater formation on extensional stepovers within strike-slip systems in southwest Iceland. Pit formation associated with dilational faults in the uppermost crust is expected, regardless of the dilational fault origin.

REFERENCES CITED

- Ferrill, D.A., Wyrick, D.Y., Morris, A.P., Sims, D.W., and Franklin, N.M., 2004, Dilational fault slip and pit chain formation on Mars: *GSA Today*, v. 14, no. 10, p. 4–12.
- Ferrill, D.A., Stamatakis, J.A., and Sims, D.W., 1999, Normal fault corrugation: Implications for growth and seismicity of active normal faults: *Journal of Structural Geology*, v. 21, p. 1027–1038.
- Ramsay, J.G., and Huber, M.I., 1987, The techniques of modern structural geology, volume 2: Folds and Fractures: London, Academic Press, 462 p.
- Schultz, R.A., 1989, Strike-slip faulting of ridged plains near Valles Marineris, Mars: *Nature*, v. 341, p. 424–426.