SUPPLEMENTAL MATERIALS TO “ASYMPTOTIC DISTRIBUTION-FREE CHANGE-POINT DETECTION FOR MULTIVARIATE AND NON-EUCLIDEAN DATA”

BY LYNNA CHU AND HAO CHEN

University of California, Davis

CONTENTS

A	Test statistics for changed interval alternative	1
A.1	Generalized edge-count scan statistic for changed interval alternative	1
A.2	Weighted edge-count scan statistic for a changed interval alternative	3
B	Exact analytic expressions for third moments	4
C	Proofs for Lemmas, Propositions and Theorems	6
C.1	Proof of Lemma 3.1	7
C.2	Proof of Theorem 4.1	7
C.3	Proof of Theorem 4.3	14
C.4	Proof of Proposition 4.4	19
D	Checking analytic p -value approximations	22
D.1	Single change-point alternative	22
D.2	Changed interval alternative	23
E	Power tables	30
F	Additional illustrations of the NYC taxi dataset	32
G	More discussions on the conditions	34
G.1	Critical Values for k -MST	34
G.2	Critical Values for k -NNG	34
H	Relationship of the three test statistics and an extension to the max-type edge count test	38
I	Block permutation for local dependency	42
	References	43

SUPPLEMENT A: TEST STATISTICS FOR CHANGED INTERVAL ALTERNATIVE

A.1. Generalized edge-count scan statistic for changed interval alternative. Here, we present the generalized edge-count two-sample test

statistic for testing the null H_0 (1.1) versus the changed interval alternative H_2 (1.3). Similar to the single change-point case, each possible interval $(t_1, t_2]$ divides the observations into two groups: one group containing all observations observed during $(t_1, t_2]$, and the other group containing all observations observed outside of this interval.

Let $g_i(t_1, t_2) = I_{t_1 < i \leq t_2}$. For an edge $e = (i, j)$, we define

$$J_e(t_1, t_2) = \begin{cases} 0 & \text{if } g_i(t_1, t_2) \neq g_j(t_1, t_2) \\ 1 & \text{if } g_i(t_1, t_2) = g_j(t_1, t_2) = 1, \\ 2 & \text{if } g_i(t_1, t_2) = g_j(t_1, t_2) = 0. \end{cases}$$

Then, for any candidate changed interval $(t_1, t_2]$ of $(\tau_1, \tau_2]$,

$$(A.1) \quad R_k(t_1, t_2) = \sum_{e \in G} I_{J_e(t_1, t_2) = k}, \quad k = 0, 1, 2,$$

and the test statistic is defined as

$$S(t_1, t_2) = \begin{pmatrix} R_1(t_1, t_2) - \mathbf{E}(R_1(t_1, t_2)) \\ R_2(t_1, t_2) - \mathbf{E}(R_2(t_1, t_2)) \end{pmatrix}^T \Sigma^{-1}(t_1, t_2) \begin{pmatrix} R_1(t_1, t_2) - \mathbf{E}(R_1(t_1, t_2)) \\ R_2(t_1, t_2) - \mathbf{E}(R_2(t_1, t_2)) \end{pmatrix}.$$

Under the permutation null, the explicit expression for $\mathbf{E}(R_1(t_1, t_2))$, $\mathbf{E}(R_2(t_1, t_2))$ and the covariance matrix can be obtained similarly as in the single change-point setting.

$$\mathbf{E}(R_1(t_1, t_2)) = |G| \frac{(t_2 - t_1)(t_2 - t_2 - 1)}{n(n-1)}.$$

$$\mathbf{E}(R_2(t_2, t_2)) = |G| \frac{(n - (t_2 - t_1))(n - (t_2 - t_1) - 1)}{n(n-1)}.$$

$$\begin{aligned} \Sigma_{11}(t_1, t_2) &= \mathbf{E}(R_1(t_1, t_2))(1 - \mathbf{E}(R_1(t_1, t_2))) \\ &\quad + \left(\sum_{i=1}^n |G_i|^2 - 2|G| \right) \frac{(t_2 - t_1)(t_2 - t_1 - 1)(t_2 - t_1 - 2)}{n(n-1)(n-2)} \\ &\quad + \left(|G|^2 - \sum_{i=1}^n |G_i|^2 + |G| \right) \frac{(t_2 - t_1)(t_2 - t_1 - 1)(t_2 - t_1 - 2)(t_2 - t_1 - 3)}{n(n-1)(n-2)(n-3)}. \end{aligned}$$

$$\begin{aligned} \Sigma_{22}(t_1, t_2) &= \mathbf{E}(R_2(t_1, t_2))(1 - \mathbf{E}(R_2(t_1, t_2))) \\ &\quad + \left(\sum_{i=1}^n |G_i|^2 - 2|G| \right) \frac{(n - t_2 - t_1)(n - t_2 + t_1 - 1)(n - t_2 + t_1 - 2)}{n(n-1)(n-2)} \\ &\quad + \left(|G|^2 - \sum_{i=1}^n |G_i|^2 + |G| \right) \frac{(n - t_2 + t_1)(n - t_2 + t_1 - 1)(n - t_2 + t_1 - 2)(n - t_2 + t_1 - 3)}{n(n-1)(n-2)(n-3)}. \end{aligned}$$

$$\begin{aligned}\Sigma_{12}(t_1, t_2) &= \Sigma_{21}(t_1, t_2) = \\ &\left(|G|^2 - \sum_{i=1}^n |G_i|^2 + |G| \right) \frac{(t_2-t_1)(t_2-t_1-1)(n-t_2+t_1)(n-t_2+t_1-1)}{n(n-1)(n-2)(n-3)} \\ &- \mathbf{E}(R_1(t_1, t_2))\mathbf{E}(R_2(t_1, t_2)).\end{aligned}$$

The scan statistic involves a maximization over t_1 and t_2 ,

$$(A.2) \quad \max_{\substack{1 \leq t_1 < t_2 \leq n \\ l_0 \leq t_2 - t_1 \leq l_1}} S(t_1, t_2),$$

where l_0 and l_1 are constraints on the window size. For example, we can set $l_1 = n - l_0$ so that only alternatives where the numbers of observations in either group is larger than l_0 are considered.

A.2. Weighted edge-count scan statistic for a changed interval alternative. Here, we present the weighted edge-count two-sample test statistic for testing the null H_0 (1.1) versus the changed interval alternative H_2 (1.3). For any candidate interval $(t_1, t_2]$, the test statistic is

$$\begin{aligned}R_w(t_1, t_2) &= q(t_1, t_2)R_1(t_1, t_2) + p(t_1, t_2)R_2(t_1, t_2), \\ \text{with } p(t_1, t_2) &= \frac{t_2-t_1-1}{n-2}, \quad q(t_1, t_2) = 1 - p(t_1, t_2)\end{aligned}$$

Similarly, $R_w(t_1, t_2)$ is standardized:

$$(A.3) \quad Z_w(t_1, t_2) = \frac{R_w(t_1, t_2) - \mathbf{E}[R_w(t_1, t_2)]}{\sqrt{\mathbf{Var}[R_w(t_1, t_2)]}}.$$

Analytical expressions for the expectation and variance of $R_w(t_1, t_2)$ under the permutation null are:

$$\begin{aligned}\mathbf{E}(R_w(t_1, t_2)) &= |G| \frac{(t_2-t_1-1)(n-t_2+t_1-1)}{(n-1)(n-2)}, \\ \mathbf{Var}(R_w(t_1, t_2)) &= \frac{(t_2-t_1)(t_2-t_1-1)(n-t_2+t_1)(n-t_2+t_1-1)}{n(n-1)(n-2)(n-3)} \times \\ &\quad \left(|G| - \frac{1}{n-2} \left(\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2}{n} \right) - \frac{2}{n(n-1)} |G|^2 \right).\end{aligned}$$

To test H_0 versus H_2 , the scan statistic involves a maximization over t_1 and t_2 ,

$$(A.4) \quad \max_{\substack{1 \leq t_1 < t_2 \leq n \\ l_0 \leq t_2 - t_1 \leq l_1}} Z_w(t_1, t_2),$$

where l_0 and l_1 are constraints on the window size.

SUPPLEMENT B: EXACT ANALYTIC EXPRESSIONS FOR THIRD
MOMENTS

The following lemma provides the exact analytic expressions for $\mathbf{E}(R_w^3(t))$ and $\mathbf{E}(R_{\text{diff}}^3(t))$.

LEMMA B.1. *We have*

$$\begin{aligned} \mathbf{E}(R_w^3(t)) = & q^3(t)\mathbf{E}(R_1^3(t)) + 3q^2(t)p(t)\mathbf{E}(R_1^2(t)R_2(t)) + \\ & 3q(t)p^2(t)\mathbf{E}(R_1(t)R_2^2(t)) + p^3(t)\mathbf{E}(R_2^3(t)), \end{aligned}$$

$$\mathbf{E}(R_{\text{diff}}^3(t)) = \mathbf{E}(R_1^3(t)) - 3\mathbf{E}(R_1^2(t)R_2(t)) + 3\mathbf{E}(R_1(t)R_2^2(t)) - \mathbf{E}(R_2^3(t)),$$

where

$$\begin{aligned} \mathbf{E}(R_1^3(t)) = & p_1(t)|G| + p_2(t) \left(3 \sum_i |G_i|(|G_i| - 1) + 2 \sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| \right) \\ & + p_3(t) \left(3|G|(|G| - 1) + \sum_i |G_i|(|G_i| - 1)(|G_i| - 5) \right) \\ & + 6p_3(t) \left(\sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) - \sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| \right) \\ & + 3p_4(t) \left(\sum_i |G_i|(|G_i| - 1)(|G| - |G_i|) + 2 \sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| \right) \\ & - 12p_4(t) \sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) \\ & + p_5(t) \left(|G|(|G| - 1)(|G| - 2) + 6 \sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) \right) \\ & - 2p_5(t) \sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| \\ & + p_5(t) \left(\sum_i |G_i|(|G_i| - 1)(3|G| - 2|G_i| - 2) \right), \end{aligned}$$

$$\mathbf{E}(R_1^2(t)R_2(t))$$

$$\begin{aligned}
&= p_6(t) \left(|G|(|G| - 1) - \sum_i |G_i|(|G_i| - 1) \right) \\
&+ p_7(t) \left(\sum_i |G_i|(|G_i| - 1)(|G| - |G_i|) - 4 \sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) + \right. \\
&\quad \left. 2 \sum_{(i,j) \in G} |\{k : (i, k), (j, k) \in G\}| \right) \\
&+ p_8(t) \left(|G|(|G| - 1)(|G| - 2) - 2 \sum_{(i,j) \in G} |\{k : (i, k), (j, k) \in G\}| + \right. \\
&\quad \left. 6 \sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) - \sum_i |G_i|(|G_i| - 1)(3|G| - 2|G_i| - 2) \right),
\end{aligned}$$

$$\begin{aligned}
&\mathbf{E}(R_1(t)R_2^2(t)) \\
&= p_6(n-t) \left(|G|(|G| - 1) - \sum_i |G_i|(|G_i| - 1) \right) \\
&+ p_7(n-t) \left(\sum_i |G_i|(|G_i| - 1)(|G| - |G_i|) - 4 \sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) + \right. \\
&\quad \left. 2 \sum_{(i,j) \in G} |\{k : (i, k), (j, k) \in G\}| \right) \\
&+ p_8(n-t) \left(|G|(|G| - 1)(|G| - 2) - 2 \sum_{(i,j) \in G} |\{k : (i, k), (j, k) \in G\}| + \right. \\
&\quad \left. 6 \sum_{(i,j) \in G} (|G_i| - 1)(|G_j| - 1) - \sum_i |G_i|(|G_i| - 1)(3|G| - 2|G_i| - 2) \right),
\end{aligned}$$

$$\begin{aligned}
&\mathbf{E}(R_2^3(t)) \\
&= p_1(n-t)|G| \\
&+ p_2(n-t) \left(3 \sum_i |G_i|(|G_i| - 1) + 2 \sum_{(i,j) \in G} |\{k : (i, k), (j, k) \in G\}| \right)
\end{aligned}$$

$$\begin{aligned}
& + p_3(n-t) \left(3|G|(|G|-1) + \sum_i |G_i|(|G_i|-1)(|G_i|-5) \right) \\
& + 6p_3(n-t) \left(\sum_{(i,j) \in G} (|G_i|-1)(|G_j|-1) - \sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| \right) \\
& + 3p_4(n-t) \left(\sum_i |G_i|(|G_i|-1)(|G|-|G_i|) \right) \\
& + 6p_4(n-t) \left(\sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| - 2 \sum_{(i,j) \in G} (|G_i|-1)(|G_j|-1) \right) \\
& + p_5(n-t) \left(|G|(|G|-1)(|G|-2) + 6 \sum_{(i,j) \in G} (|G_i|-1)(|G_j|-1) \right) \\
& - 2p_5(n-t) \sum_{(i,j) \in G} |\{k : (i,k), (j,k) \in G\}| \\
& - p_5(n-t) \sum_i |G_i|(|G_i|-1)(3|G|-2|G_i|-2),
\end{aligned}$$

with

$$\begin{aligned}
p_j(t) &= \prod_{i=0}^j \frac{(t-i)}{(n-i)}, \quad j = 1, \dots, 5, \\
p_6(t) &= \frac{t(t-1)(n-t)(n-t-1)}{n(n-1)(n-2)(n-3)}, \\
p_7(t) &= \frac{t(t-1)(n-t)(n-t-1)(t-2)}{n(n-1)(n-2)(n-3)(n-4)}, \\
p_8(t) &= \frac{t(t-1)(n-t)(n-t-1)(t-2)(t-3)}{n(n-1)(n-2)(n-3)(n-4)(n-5)}.
\end{aligned}$$

The proof of this lemma can be followed straightforwardly from the proof for Lemma 3.8 in [Chen and Zhang \(2015\)](#) and is omitted here.

SUPPLEMENT C: PROOFS FOR LEMMAS, PROPOSITIONS AND THEOREMS

C.1. Proof of Lemma 3.1. Let $\mathbf{R}(t) = \begin{pmatrix} R_1(t) \\ R_2(t) \end{pmatrix}$, $\mathbf{A}(t) = \begin{pmatrix} 1 & -1 \\ q(t) & p(t) \end{pmatrix}$.

It is easy to see that $\mathbf{A}(t)$ is invertible. Then

$$\begin{aligned} S(t) &= (\mathbf{R}(t) - \mathbf{E}(\mathbf{R}(t)))^T (\boldsymbol{\Sigma}(t))^{-1} (\mathbf{R}(t) - \mathbf{E}(\mathbf{R}(t))) \\ &= (\mathbf{R}(t) - \mathbf{E}(\mathbf{R}(t)))^T \mathbf{A}^T(t) (\mathbf{A}^T(t))^{-1} \boldsymbol{\Sigma}^{-1}(t) \\ &\quad \times \mathbf{A}^T(t) (\mathbf{A}^T(t))^{-1} (\mathbf{R}(t) - \mathbf{E}(\mathbf{R}(t))) \\ &= (\mathbf{A}(t)(\mathbf{R}(t) - \mathbf{E}(\mathbf{R}(t))))^T (\mathbf{A}(t)\boldsymbol{\Sigma}(t)\mathbf{A}^T(t))^{-1} (\mathbf{A}(t)(\mathbf{R}(t) - \mathbf{E}(\mathbf{R}(t)))). \end{aligned}$$

The elements of $\boldsymbol{\Sigma}(t)$ can be found in Section 3.1 and re-organized as

$$\begin{aligned} \Sigma_{11}(t) &= \frac{(n-t)t(n-t-1)(t-1)}{n(n-1)(n-2)(n-3)} \left(|G| + \frac{t-2}{n-t-1} \left(\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2}{n} \right) - \frac{2}{n(n-1)} |G|^2 \right), \\ \Sigma_{12}(t) &= \frac{(n-t)t(n-t-1)(t-1)}{n(n-1)(n-2)(n-3)} \left(|G| - \left(\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2}{n} \right) - \frac{2}{n(n-1)} |G|^2 \right), \\ \Sigma_{22}(t) &= \frac{(n-t)t(n-t-1)(t-1)}{n(n-1)(n-2)(n-3)} \left(|G| + \frac{n-t-2}{t-1} \left(\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2}{n} \right) - \frac{2}{n(n-1)} |G|^2 \right). \end{aligned}$$

With the above reorganization, it is not hard to show that

$$\begin{aligned} &\mathbf{A}(t)\boldsymbol{\Sigma}(t)\mathbf{A}^T(t) \\ &= \begin{pmatrix} \Sigma_{11}(t) + \Sigma_{22}(t) - 2\Sigma_{12}(t) & 0 \\ 0 & q(t)\Sigma_{11}(t) + (p(t) - q(t))\Sigma_{12}(t) - p(t)\Sigma_{22}(t) \end{pmatrix} \end{aligned}$$

Then,

$$\begin{aligned} S(t) &= \left(\frac{R_1(t) - R_2(t) - (\mathbf{E}(R_1(t)) - \mathbf{E}(R_2(t)))}{\sqrt{\mathbf{Var}(R_1(t) - R_2(t))}} \right)^2 \\ &\quad + \left(\frac{q(t)R_1(t) + p(t)R_2(t) - (q(t)\mathbf{E}(R_1(t)) + p(t)\mathbf{E}(R_2(t)))}{\sqrt{\mathbf{Var}(q(t)R_1(t) + p(t)R_2(t))}} \right)^2 \\ &= Z_{\text{diff}}^2(t) + Z_w^2(t). \end{aligned}$$

The proof for $S(t_1, t_2) = Z_{\text{diff}}^2(t_1, t_2) + Z_w^2(t_1, t_2)$ can be shown in a similar manner.

C.2. Proof of Theorem 4.1. Here, we prove that $\{Z_w([nu]) : 0 < u < 1\}$ and $\{Z_{\text{diff}}([nu]) : 0 < u < 1\}$ converge to independent Gaussian processes in finite dimensional distributions. The proof for the convergence

of $\{Z_w([nu], [nv]) : 0 < u < v < 1\}$ and $\{Z_{\text{diff}}([nu], [nv]) : 0 < u < v < 1\}$ to two-dimensional Gaussian random fields in finite dimensional distributions can be done in a similar way.

To prove the two processes converges to two independent Gaussian processes in finite dimensional distributions, we first prove that

$$(Z_w([nu_1]), \dots, Z_w([nu_K]), Z_{\text{diff}}([nu_1]), \dots, Z_{\text{diff}}([nu_K]))$$

converges to a multivariate Gaussian distribution as $n \rightarrow \infty$ for any $0 < u_1 < u_2 < \dots < u_K < 1$ and fixed K and then show that $\text{cov}(Z_w(u), Z_{\text{diff}}(v)) = 0$ for any $0 < u, v < 1$ as $n \rightarrow \infty$. For notation simplicity, let $t_k = [nu_k]$, $k = 1, \dots, K$.

To prove $(Z_w(t_1), \dots, Z_w(t_K), Z_{\text{diff}}(t_1), \dots, Z_{\text{diff}}(t_K))$ converges to a multivariate Gaussian distribution, we revisit the permutation distribution. In permutation distribution, we permute the order of the observations. Let $\pi(i)$ be the observed time of $\mathbf{y}_i$ after permutation. Then $(\pi(1), \pi(2), \dots, \pi(n))$ is a permutation of $1, \dots, n$. To obtain the permutation distribution, we can do it in two steps: (1) For each i , $\tilde{\pi}(i)$ is sampled uniformly from 1 to n ; (2) only those $(\tilde{\pi}(1), \tilde{\pi}(2), \dots, \tilde{\pi}(n))$ such that each value in $\{1, \dots, n\}$ is sampled once are retained. It is not hard to see that each permutation has the same occurrence probability after these two steps. We call the distribution resulting from only performing the first step the bootstrap distribution, and use $\mathbf{P}_B$, $\mathbf{E}_B$, $\mathbf{Var}_B$, $\mathbf{Cov}_B$ to denote the probability, expectation, variance, and covariance under the bootstrap distribution, respectively. In this section, the corresponding quantities with the subscript P are used to denote the equivalences under the permutation distribution.

Given that the $\tilde{\pi}(i)$'s are independent under the bootstrap null distribution, we have

$$\begin{aligned} \mathbf{E}_B(R_1(t)) &= \frac{t^2}{n^2}|G|, \\ \mathbf{E}_B(R_2(t)) &= \frac{(n-t)^2}{n^2}|G|, \\ \mathbf{Var}_B(R_1(t)) &= \frac{t^2(n-t)^2}{n^4}|G| + \frac{t^3(n-t)}{n^4} \sum_{i=1}^n |G_i|^2, \\ \mathbf{Var}_B(R_2(t)) &= \frac{t^2(n-t)^2}{n^4}|G| + \frac{t(n-t)^3}{n^4} \sum_{i=1}^n |G_i|^2, \\ \mathbf{Cov}_B(R_1(t), R_2(t)) &= \frac{t^2(n-t)^2}{n^4}|G| - \frac{t^2(n-t)^2}{n^4} \sum_{i=1}^n |G_i|^2. \end{aligned}$$

For $R_w(t) = q(t)R_1(t) + p(t)R_2(t)$, we have

$$\begin{aligned}\mathbf{E}_B(R_w(t)) &= \frac{t^2(n-t-1) + (n-t)^2(t-1)}{n^2(n-2)}|G| := \mu_w^B(t)|G|, \\ \mathbf{Var}_B(R_w(t)) &= \frac{t^2(n-t)^2}{n^4}|G| + \frac{t(n-t)(n-2t)^2}{n^4(n-2)^2} \sum_{i=1}^n |G_i|^2 \\ &:= (\sigma_w^B(t))^2.\end{aligned}$$

For $R_{\text{diff}}(t) = R_1(t) - R_2(t)$, we have

$$\begin{aligned}\mathbf{E}_B(R_{\text{diff}}(t)) &= \frac{(2t-n)}{n}|G| := \mu_{\text{diff}}^B(t)|G|, \\ \mathbf{Var}_B(R_{\text{diff}}(t)) &= \frac{2t(n-t)^2(n-2t)}{n^4} \sum_{i=1}^n |G_i|^2 := (\sigma_{\text{diff}}^B(t))^2.\end{aligned}$$

Let

$$\begin{aligned}Z_w^B(t) &= \frac{R_w - \mathbf{E}_B(R_w(t))}{\mathbf{Var}_B(R_w(t))}, \\ Z_{\text{diff}}^B(t) &= \frac{R_{\text{diff}} - \mathbf{E}_B(R_{\text{diff}}(t))}{\mathbf{Var}_B(R_{\text{diff}}(t))}, \\ X^B(t) &= \frac{n^B(t) - t}{\sqrt{t(1-t/n)}}, \text{ where } n^B(t) = \sum_{i=1}^n I_{\tilde{\pi}(i) \leq t}.\end{aligned}$$

To prove $(Z_w(t_1), \dots, Z_w(t_K), Z_{\text{diff}}(t_1), \dots, Z_{\text{diff}}(t_K))$ converges to a multivariate Gaussian distribution under the conditions on the graph in Theorem 4.1, as $n \rightarrow \infty$, we only need to prove the following two lemmas:

LEMMA C.1. *When $|G| = O(n^\alpha)$, $1 \leq \alpha < 1.5$, $\sum_{e \in G} |A_e||B_e| = o(n^{1.5\alpha})$, $\sum_{e \in G} |A_e|^2 = o(n^{\alpha+0.5})$, and $\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2}{n} = O(\sum_{i=1}^n |G_i|^2)$, for $0 < u_1 < u_2 < \dots < u_k < 1$, as $n \rightarrow \infty$, under the bootstrap distribution ,*

$$(C.1) \quad (Z_w^B(t_1), \dots, Z_w^B(t_k), Z_{\text{diff}}^B(t_1), \dots, Z_{\text{diff}}^B(t_k), X^B(t_1), \dots, X^B(t_k))$$

is multivariate normal and the covariance matrix of

$$(X^B(t_1), X^B(t_2), \dots, X^B(t_k))$$

is positive definite.

LEMMA C.2. *We have,*

1. $\frac{\mathbf{Var}_B(R_w(t))}{\mathbf{Var}_P(R_w(t))} \rightarrow 1$
2. $\frac{\mathbf{Var}_B(R_{\text{diff}}(t))}{\mathbf{Var}_P(R_{\text{diff}}(t))} \rightarrow c_1$
3. $\frac{\mathbf{E}_B(R_w(t)) - \mathbf{E}_P(R_w(t))}{\sqrt{\mathbf{Var}_P(R_w(t))}} \rightarrow 0$
4. $\frac{\mathbf{E}_B(R_{\text{diff}}(t)) - \mathbf{E}_P(R_{\text{diff}}(t))}{\sqrt{\mathbf{Var}_P(R_{\text{diff}}(t))}} \rightarrow 0$

where c_1 is a positive constant.

From Lemma C.1,

$$(Z_w^B(t_1), \dots, Z_w^B(t_k), Z_{\text{diff}}^B(t_1), \dots, Z_{\text{diff}}^B(t_k) | X^B(t_1), X^B(t_2), \dots, X^B(t_k))$$

is multivariate normal under the bootstrap distribution. Since $(Z_w^B(t_1), \dots, Z_w^B(t_k), Z_{\text{diff}}^B(t_1), \dots, Z_{\text{diff}}^B(t_k) | X^B(t_1) = 0, \dots, X^B(t_k) = 0)$ under the bootstrap distribution has the same distribution as $(Z_w^B(t_1), \dots, Z_w^B(t_k), Z_{\text{diff}}^B(t_1), \dots, Z_{\text{diff}}^B(t_k))$ under the permutation distribution, and

$$Z_w(t) = \frac{\mathbf{Var}_B(R_w(t))}{\mathbf{Var}_P(R_w(t))} \left(Z_w^B(t) + \frac{\mathbf{E}_B(R_w(t)) - \mathbf{E}_P(R_w(t))}{\sqrt{\mathbf{Var}_B(R_w(t))}} \right),$$

$$Z_{\text{diff}}(t) = \frac{\mathbf{Var}_B(R_{\text{diff}}(t))}{\mathbf{Var}_P(R_{\text{diff}}(t))} \left(Z_{\text{diff}}^B(t) + \frac{\mathbf{E}_B(R_{\text{diff}}(t)) - \mathbf{E}_P(R_{\text{diff}}(t))}{\sqrt{\mathbf{Var}_B(R_{\text{diff}}(t))}} \right),$$

with Lemma C.2, we conclude that

$$(Z_w([nu_1]), Z_w([nu_2]), \dots, Z_w([nu_k]), Z_{\text{diff}}([nu_1]), Z_{\text{diff}}([nu_2]), \dots, Z_{\text{diff}}([nu_k]))$$

is multivariate Gaussian under the permutation distribution.

To prove C.1, we only need to show that $\sum_{k=1}^K (a_k Z_w^B(t_k) + b_k Z_{\text{diff}}^B(t_k) + c_k X^B(t_k))$ is normal for any fixed a_k , b_k , and c_k for the non-degenerating case that $\mathbf{Var}_B(W) := \mathbf{Var}_B(a_k Z_w^B(t_k) + b_k Z_{\text{diff}}^B(t_k) + c_k X^B(t_k)) > 0$. We prove Gaussianity of $\sum_{k=1}^K (a_k Z_w^B(t_k) + b_k Z_{\text{diff}}^B(t_k) + c_k X^B(t_k))$ by Stein's method.

Consider sums of the form $W = \sum_{i \in \mathcal{J}} \xi_i$ where $\mathcal{J}$ is an index set and ξ are random variables with $E[\xi_i] = 0$, and $E[W^2] = 1$. The following assumption restricts the dependence among $\{\xi_i : i \in \mathcal{J}\}$.

ASSUMPTION C.3. (*Chen and Shao (2005), pg. 17*) For each $j \in \mathcal{J}$ there exists $S_i \subset T_i \subset \mathcal{J}$ such that ξ_i is independent of $\xi_{S_i^c}$ and ξ_{S_i} is independent of $\xi_{T_i^c}$.

We will use the following specific form of Stein's method.

THEOREM C.4. (*Chen and Shao (2005), Theorem 3.4*) Under Assumption C.3, we have

$$\sup_{h \in Lip(1)} |\mathbf{E}h(W) - \mathbf{E}h(Z)| \leq \delta$$

where $Lip(1) = \{h : \mathbb{R} \rightarrow \mathbb{R}, \|h'\| \leq 1\}$, Z has $\mathcal{N}(0, 1)$ distribution and

$$\delta = 2 \sum_{j \in \mathcal{J}} (\mathbf{E}|\xi_i \eta_i \theta_i| + |\mathbf{E}(\xi_i \eta_i)| |\mathbf{E}|\theta_i||) + \sum_{j \in \mathcal{J}} \mathbf{E}|\xi_i \eta_i^2|$$

with $\eta_i = \sum_{j \in S_i} \eta_j$ and $\theta_i = \sum_{j \in T_i} \theta_j$, where S_i and T_i are defined in Assumption C.3.

We adopt the same notations with the index set $\mathcal{J} = \{e : e \in G\} \cup \{1, \dots, n\}$. Let

$$\begin{aligned} \xi_{e,k} = & \frac{1}{\sigma_w^B(t_k)} \left(\frac{n-t_k-1}{n-2} I_{\tilde{\pi}(e_-) \leq t_k, \tilde{\pi}(e_+) \leq t_k} + \frac{t_k-1}{n-2} I_{\tilde{\pi}(e_-) > t_k, \tilde{\pi}(e_+) > t_k} - \mu_w^B(t_k) \right) \\ & + \frac{1}{\sigma_{\text{diff}}^B(t_k)} \left(I_{\tilde{\pi}(e_-) \leq t_k, \tilde{\pi}(e_+) \leq t_k} - I_{\tilde{\pi}(e_-) > t_k, \tilde{\pi}(e_+) > t_k} - \mu_{\text{diff}}^B(t_k) \right) \end{aligned}$$

and

$$\xi_{i,k} = \frac{I_{\tilde{\pi}(i) \leq t_k} - \frac{t_k}{n}}{\sqrt{t_k(1 - t_k/n)}}.$$

Let $\xi_e = \sum_k a_k \xi_{e,k}$, $\xi_i = \sum_k b_k \xi_{i,k}$. Then $W = \sum_{j \in \mathcal{J}} \xi_j = \sum_k (a_k Z_w^B(t_k) + b_k Z_{\text{diff}}^B(t_k) + c_k X^B(t_k))$. We have $\mathbf{E}_B(W) = 0$, $\mathbf{E}_B(W^2) = 1$. Let $a = \max(\max_k a_k, \max_k b_k, \max_k c_k)$, $\sigma_B = \min(\min_k \sigma_w^B(t_k), \min_k \sigma_{\text{diff}}^B(t_k))$, and $\sigma_0 = \min_k(\sqrt{t_k(1 - t_k/n)})$. Then

$$|\xi_e| \leq \frac{2aK}{\sigma_B}, \forall e \in G; \quad |\xi_i| \leq \frac{aK}{\sigma_0}, \forall i \in \{1, \dots, n\}.$$

For $e \in G$, let

$$S_e = \{A_e, e_-, e_+\},$$

$$T_e = B_e \cup \{\text{Nodes in } A_e\},$$

where A_e, B_e are defined in 4.7 and 4.8. Then S_e and T_e satisfy Assumption C.3.

For $i = 1, \dots, n$, let

$$S_i = \{e \in G_i\}$$

$$T_i = \{e \in G_{i,2}\} \cup \{\text{Nodes in } G_i\},$$

where $G_{i,2}$ is the subgraph of G including all edges connect to any node in G_i . Then S_i and T_i satisfy Assumption C.3.

For $j \in \mathcal{J}$, let $\eta_j = \sum_{k \in K_j} \xi_k$, $\theta_j = \sum_{k \in L_j} \xi_k$. By Theorem C.4, we have $\sum_{h \in Lip(1)} |\mathbf{E}_B h(W) - \mathbf{E}_B h(Z)| \leq \delta$ for $Z \sim \mathcal{N}(0, 1)$, where

$$\begin{aligned} \delta &= \frac{1}{\sqrt{\mathbf{Var}_B(W)}} \left(2 \sum_{j \in \mathcal{J}} (\mathbf{E}_B |\xi_j \eta_j \theta_j| + |\mathbf{E}_B(\xi_j \eta_j)| |\mathbf{E}_B \theta_j|) + \sum_{j \in \mathcal{J}} \mathbf{E}_B |\xi_j \eta_j^2| \right) \\ &\leq \frac{a^3 K^3}{\sqrt{\mathbf{Var}_B(W)}} \left(10 \sum_{e \in G} \frac{1}{\sigma_B} \left(\frac{|A_e|}{\sigma_B} + \frac{2}{\sigma_0} \right) \left(\frac{|B_e|}{\sigma_B} + 2 \frac{|A_e|}{\sigma_0} \right) \right. \\ &\quad \left. + 5 \sum_{i=1}^n \frac{1}{\sigma_0} \left(\frac{|G_i|}{\sigma_B} + \frac{1}{\sigma_0} \right) \left(\frac{|G_{i,2}|}{\sigma_B} + 2 \frac{|G_i|}{\sigma_0} \right) \right). \end{aligned}$$

Since σ_B is at least of order $|G|^{0.5}$ and $\sigma_0 = O(n^{0.5})$, as long as the following results hold, we have $\delta \rightarrow 0$ as $n \rightarrow \infty$.

$$(C.2) \quad \sum_{e \in G} |A_e| |B_e| = o(|G|^{1.5}),$$

$$(C.3) \quad \sum_{e \in G} |A_e|^2 = o(|G| n^{0.5}),$$

$$(C.4) \quad \sum_{e \in G} |B_e| = o(|G| n^{0.5}),$$

$$(C.5) \quad \sum_{e \in G} |A_e| = o(|G|^{0.5} n),$$

$$(C.6) \quad \sum_{i=1}^n |G_i| |G_{i,2}| = o(|G| n^{0.5})$$

$$(C.7) \quad \sum_{i=1}^n |G_i|^2 = o(|G|^{0.5} n)$$

$$(C.8) \quad \sum_{i=1}^n |G_{i,2}| = o(|G|^{0.5} n)$$

$$(C.9) \quad \sum_{i=1}^n |G_i| = o(n^{1.5}).$$

To show the above 9 results, recall the conditions in Theorem 4.1 are

1. $|G| = O(n^\alpha)$, $1 \leq \alpha < 1.5$,

2. $\sum_{e \in G} |A_e| |B_e| = o(n^{1.5\alpha})$,
3. $\sum_{e \in G} |A_e|^2 = o(n^{\alpha+0.5})$.

Substituting $|G|$ by $O(n^\alpha)$, then (C.2), (C.3), and (C.9) follow immediately.

Since $|B_e| \leq \sum_{e^* \in A_e} |A_{e^*}|$, we have $\sum_{e \in G} |B_e| \leq \sum_{e \in G} \sum_{e^* \in A_e} |A_{e^*}| = \sum_{e \in G} |A_e|^2$. So (C.3) ensures (C.4).

By Cauchy-Schwartz, we have $\sum_{e \in G} |A_e| \leq \sqrt{\sum_{e \in G} |A_e|^2 \sum_{e \in G} 1^2} = o(|G|n^{0.25})$. Since $(\alpha + 0.25) - (0.5\alpha + 1) = 0.5\alpha - 0.75 < 0$ when $\alpha < 1.5$, then (C.5) follows.

We use $\mathcal{V}_{G_i}$ to denote the vertex set of G_i . Since $|G_{i,2}| \leq \sum_{j \in \mathcal{V}_{G_i}} |G_j|$, we have $\sum_{i=1}^n |G_{i,2}| = \sum_{i=1}^n \sum_{j \in \mathcal{V}_{G_i}} |G_j| = \sum_{(i,j) \in G} (|G_i| + |G_j|) \leq \sum_{e \in G} 2|A_e|$. So (C.5) ensures (C.8).

Next, $\sum_{e \in G} |A_e| = \sum_{(i,j) \in G} (|G_i| + |G_j| - 1) = \sum_{i=1}^n \sum_{j \in \mathcal{V}_{G_i}} |G_j| - |G| = \sum_{i=1}^n |G_i|^2 - |G|$. Since $|G| = o(n^{0.5\alpha+1})$ when $|G| = O(n^\alpha)$, $\alpha < 1.5$ and so (C.5) and (C.7) are equivalent.

For (C.6), we have $\sum_{i=1}^n |G_i| |G_{i,2}| \leq \sum_{i=1}^n |G_i| \sum_{j \in \mathcal{V}_{G_i}} |G_j| = \sum_{i=1}^n \sum_{j \in G_j} |G_i| |G_j| = 2 \sum_{(i,j) \in G} |G_i| |G_j| \leq 2 \sum_{e \in A_e} |A_e|^2$. So (C.3) ensures (C.6).

Hence, (C.2) - (C.9) can be derived from conditions in Theorem 4.1.

The proof for the second part of Lemma C.1 (showing that the covariance matrix of $(X^B(t_1), X^B(t_2), \dots, X^B(t_k))$ is positive definite) can be found in the Appendix of Chen and Zhang (2015) (Theorem 3.1) and is omitted here.

Now we prove Lemma C.2:

We have,

$$\lim_{n \rightarrow \infty} \frac{\text{Var}_P(R_w(t))}{\text{Var}_B(R_w(t))} = 1 - \lim_{n \rightarrow \infty} \left(\frac{\sum_{i=1}^n |G_i|^2}{n|G|} - \frac{4|G|}{n^2} + \frac{2|G|}{n^2} \right).$$

Since $|G| = O(n^\alpha)$, $\alpha < 1.5$ and we have $\sum_{i=1}^n |G_i|^2 = o(n^{0.5\alpha+1})$ from the proof for Lemma C.1, we have

$$\lim_{n \rightarrow \infty} \left(\frac{\sum_{i=1}^n |G_i|^2}{n|G|} - \frac{4|G|}{n^2} + \frac{2|G|}{n^2} \right) = 0$$

which leads to $\lim_{n \rightarrow \infty} \frac{\text{Var}_P(R_w(t))}{\text{Var}_B(R_w(t))} = 1$.

Since $\lim_{n \rightarrow \infty} \mathbf{E}_B(R_w(t)) - \mathbf{E}_P(R_w(t)) = \frac{t(n-t)}{n^2(n-1)} |G|$ we have

$$\lim_{n \rightarrow \infty} \frac{\mathbf{E}_B(R_w(t)) - \mathbf{E}_P(R_w(t))}{\sqrt{\text{Var}_P(R_w(t))}} = \lim_{n \rightarrow \infty} \frac{\sqrt{|G|}}{n-2} = 0,$$

when $|G| = O(n^\alpha)$, $\alpha < 1.5$.

Similarly for $R_{\text{diff}}(t)$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\mathbf{Var}_P(R_{\text{diff}}(t))}{\mathbf{Var}_B(R_{\text{diff}}(t))} &= \lim_{n \rightarrow \infty} \frac{\frac{n^2}{2(n-1)(n-t)(n-2t)} (\sum_{i=1}^n n|G_i|^2 - 4|G|^2)}{\sum_{i=1}^n |G_i|^2} \\ &= \lim_{n \rightarrow \infty} \left(\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2/n}{\sum_{i=1}^n |G_i|^2} \right). \end{aligned}$$

By Cauchy-Schwartz inequality $\sum_{i=1}^n |G_i|^2 \geq \frac{4|G|^2}{n}$ and the condition that $\sum_{i=1}^n |G_i|^2 - \frac{4|G|^2}{n} = O(\sum_{i=1}^n |G_i|^2)$, we have $\lim_{n \rightarrow \infty} \left(\frac{\sum_{i=1}^n |G_i|^2 - 4|G|^2/n}{\sum_{i=1}^n |G_i|^2} \right)$ converges to a positive constant.

Since $\mathbf{E}_B(R_{\text{diff}}(t)) - \mathbf{E}_P(R_{\text{diff}}(t)) = 0$, we have

$$\lim_{n \rightarrow \infty} \frac{\mathbf{E}_B(R_{\text{diff}}(t)) - \mathbf{E}_P(R_{\text{diff}}(t))}{\sqrt{\mathbf{Var}_P(R_{\text{diff}}(t))}} = 0$$

What's left to show is as $n \rightarrow \infty$, $\mathbf{Cov}_P(Z_w(s), Z_{\text{diff}}(t)) = 0 \forall s, t$. WLOG, let $s < t$ and $\lim_{n \rightarrow \infty} s/n = u$, $\lim_{n \rightarrow \infty} t/n = v$.

Since

$$\mathbf{Cov}_P(Z_w(s), Z_{\text{diff}}(t)) = \frac{\mathbf{E}_P(R_w(s)R_{\text{diff}}(t)) - \mathbf{E}_P(R_w(s))\mathbf{E}_P(R_{\text{diff}}(t))}{\sqrt{\mathbf{Var}_P(R_w(s))\mathbf{Var}_P(R_{\text{diff}}(t))}},$$

from the proof for Theorem 4.3 (see Section C.3), after simplification we have

$$\begin{aligned} &\mathbf{E}_P(R_w(s)R_{\text{diff}}(t)) \\ &= q(s) (\mathbf{E}_P(R_1(s)R_1(t)) - \mathbf{E}_P(R_1(s)R_2(t))) \\ &\quad + p(s) (\mathbf{E}_P(R_2(s)R_1(t)) - \mathbf{E}_P(R_2(s)R_2(t))) \\ &= |G|^2 \frac{(n-2t)(s-1)(s-n+1)}{n(n-1)(n-2)}. \end{aligned}$$

Then,

$$\begin{aligned} \lim_{n \rightarrow \infty} (\mathbf{E}_P(R_w(s)R_{\text{diff}}(t)) - \mathbf{E}_P(R_w(s))\mathbf{E}_P(R_{\text{diff}}(t))) &= \\ |G|^2(1-2v)u(u-1) - |G|u(1-u)|G|(v^2 - (1-v)^2) &= 0. \end{aligned}$$

C.3. Proof of Theorem 4.3. Here, we show in details the derivation for $\rho_{\text{diff}}^*(u, v)$, and that for $\rho_w^*(u, v)$ can be done in a similar manner.

Let $\rho_{\text{diff}}(u, v) := \mathbf{Cov}(Z_{\text{diff}}([nu]), Z_{\text{diff}}([nv]))$, then

$$\rho_{\text{diff}}^*(u, v) = \lim_{n \rightarrow \infty} \rho_{\text{diff}}(u, v).$$

Let $s = [nu], t = [nv]$. We first show for $u \leq v$. Then $s \leq t$. Since

$$\mathbf{Cov}(Z_{\text{diff}}(s), Z_{\text{diff}}(t)) = \frac{\mathbf{E}[(R_1(s) - R_2(s))(R_1(t) - R_2(t)) - \mathbf{E}[R_1(s) - R_2(s)]\mathbf{E}[R_1(t) - R_2(t)]]}{\sqrt{\mathbf{Var}(R_1(s) - R_2(s))\mathbf{Var}(R_1(t) - R_2(t))}},$$

where the expressions for $\mathbf{E}[R_1(s) - R_2(s)]$, $\mathbf{E}[R_1(t) - R_2(t)]$, $\mathbf{Var}(R_1(s) - R_2(s))$, $\mathbf{Var}(R_1(t) - R_2(t))$ follow easily from $\mathbf{E}(R_1(t))$, $\mathbf{E}(R_2(t))$, $\mathbf{Var}(R_1(t))$, $\mathbf{Var}(R_2(t))$ and $\mathbf{Cov}(R_1(t), R_2(t))$ provided in Section 3.1. So we only need to figure out

$$\begin{aligned} \mathbf{E}[(R_1(s) - R_2(s))(R_1(t) - R_2(t))] = \\ \mathbf{E}(R_1(s)R_1(t)) - \mathbf{E}(R_1(s)R_2(t)) - \mathbf{E}(R_2(s)R_1(t)) + \mathbf{E}(R_2(s)R_2(t)). \end{aligned}$$

We derive the expressions for (i) $\mathbf{E}(R_1(s)R_1(t))$, (ii) $\mathbf{E}(R_1(s)R_2(t))$, (iii) $\mathbf{E}(R_2(s)R_1(t))$, and (iv) $\mathbf{E}(R_2(s)R_2(t))$ in the following.

(i) We have

$$\mathbf{E}(R_1(s)R_1(t)) = \sum_{e_1, e_2 \in G} \mathbf{P}(J_{e_1}(s) = 1, J_{e_2}(t) = 1).$$

Let edge $e_1 = (i, j)$ and $e_2 = (k, l)$. By considering the different ways of placing i, j, k, l , we have

$$\mathbf{P}(g_i(s) = 0, g_j(s) = 0, g_k(t) = 0, g_l(t) = 0) = \begin{cases} \frac{s(s-1)}{n(n-1)} := a_1(s), & \text{if } \begin{cases} i = k, j = l \\ i = l, j = k \end{cases} \\ \frac{s(s-1)(t-2)}{n(n-1)(n-2)} := a_2(s, t), & \text{if } \begin{cases} i = k, j \neq l \\ i = l, j \neq k \\ j = k, i \neq l \\ j = l, i \neq k \end{cases} \\ \frac{s(s-1)(t-2)(t-3)}{n(n-1)(n-2)(n-3)} := a_3(s, t), & \text{if } i, j, k, l \text{ all different.} \end{cases}$$

Then,

$$\begin{aligned} \mathbf{E}(R_1(s)R_1(t)) \\ = \sum_{(i,j), (k,l) \in G} \mathbf{P}(g_i(s) = 0, g_j(s) = 0, g_k(t) = 0, g_l(t) = 0) \end{aligned}$$

$$\begin{aligned}
&= \sum_{(i,j) \in G} a_1(s) + \sum_{(i,j),(k,l) \in G} a_2(s,t) + \sum_{\substack{(i,j),(k,l) \in G \\ i,j,k,l \text{ all different}}} a_3(s,t) \\
&= (a_1(s) - 2a_2(s,t) + a_3(s,t))|G| + (a_2(s,t) - a_3(s,t)) \sum_{i=1}^n |G_i|^2 + a_3(s,t)|G|^2 \\
&= \frac{s(s-1)(n-t)(n-t-1)}{n(n-1)(n-2)(n-3)} \left(|G| + \frac{t-2}{n-t-1} \left(\sum_{i=1}^n |G_i|^2 + \frac{t-3}{n-t} |G|^2 \right) \right).
\end{aligned}$$

(ii) Since

$$\mathbf{E}(R_1(s)R_2(t)) = \sum_{e_1, e_2 \in G} \mathbf{P}(J_{e_1}(s) = 1, J_{e_2}(t) = 2),$$

and

$$\mathbf{P}(g_i(s) = 0, g_j(s) = 0, g_k(t) = 1, g_l(t) = 1) = \frac{s(s-1)(n-t)(n-t-1)}{n(n-1)(n-2)(n-3)} := b_1(s, t),$$

we have

$$\begin{aligned}
\mathbf{E}(R_1(s)R_2(t)) &= \sum_{(i,j),(k,l)} \mathbf{P}(g_i(s) = 0, g_j(s) = 0, g_k(t) = 1, g_l(t) = 1) \\
&= b_1(s, t)(|G|^2 - \sum_{i=1}^n |G_i|^2 + |G|).
\end{aligned}$$

(iii) Since

$$\mathbf{E}(R_2(s)R_1(t)) = \sum_{e_1, e_2 \in G} \mathbf{P}(J_{e_1}(s) = 2, J_{e_2}(t) = 1),$$

and

$$\mathbf{P}(g_i(s) = 1, g_j(s) = 1, g_k(t) = 0, g_l(t) = 0) = \begin{cases} \frac{(t-s)(t-s-1)}{n(n-1)} := c_1(s, t), & \text{if } \begin{cases} i = k, j = l \\ i = l, j = k \end{cases} \\ \frac{(t-s)[(t-s-1)(t-2) + (n-t)(t-1)]}{n(n-1)(n-2)} := c_2(s, t), & \text{if } \begin{cases} i = k, j \neq l \\ i = l, j \neq k \\ j = k, i \neq l \\ j = l, i \neq k \end{cases} \\ \frac{(t-s)(n-s-2)[(t-s-1)(n-s-3) + 2s(n-s-1)]}{n(n-1)(n-2)(n-3)} + \frac{s(s-1)(n-s)(n-s-1)}{n(n-1)(n-2)(n-3)} := c_3(s, t), & \text{if } i, j, k, l \text{ all different,} \end{cases}$$

we have,

$$\begin{aligned}
& \mathbf{E}(R_2(s)R_1(t)) \\
&= \sum_{(i,j),(k,l) \in G} \mathbf{P}(g_i(t) = 1, g_j(t) = 1, g_k(t) = 0, g_l(t) = 0) \\
&= \sum_{(i,j) \in G} c_1(s) + \sum_{(i,j),(k,l) \in G} c_2(s, t) + \sum_{\substack{(i,j),(k,l) \in G \\ i,j,k,l \text{ all different}}} c_3(s, t) \\
&= (c_1(s) - 2c_2(s, t) + c_3(s, t))|G| + (c_2(s, t) - c_3(s, t)) \sum_{i=1}^n |G_i|^2 + c_3(s, t)|G|^2.
\end{aligned}$$

(iv) Since

$$\mathbf{E}(R_2(s)R_2(t)) = \sum_{e_1, e_2 \in G} \mathbf{P}(J_{e_1}(s) = 2, J_{e_2}(t) = 2),$$

and

$$\mathbf{P}(g_i(s) = 1, g_j(s) = 1, g_k(t) = 1, g_l(t) = 1) = \begin{cases} \frac{(n-t)(n-t-1)}{n(n-1)} := d_1(t), & \text{if } \begin{cases} i = k, j = l \\ i = l, j = k \end{cases} \\ \frac{(n-t)(n-t-1)(n-s-2)}{n(n-1)(n-2)} := d_2(s, t), & \text{if } \begin{cases} i = k, j \neq l \\ i = l, j \neq k \\ j = k, i \neq l \\ j = l, i \neq k \end{cases} \\ \frac{(n-t)(n-t-1)(n-s-2)(n-s-3)}{n(n-1)(n-2)(n-3)} := d_3(s, t), & \text{if } i, j, k, l \text{ all different,} \end{cases}$$

we have

$$\begin{aligned}
& \mathbf{E}(R_2(s)R_2(t)) \\
&= \sum_{(i,j),(k,l) \in G} \mathbf{P}(g_i(t) = 1, g_j(t) = 1, g_k(t) = 1, g_l(t) = 1) \\
&= \sum_{(i,j) \in G} d_1(s) + \sum_{(i,j),(k,l) \in G} d_2(s, t) + \sum_{\substack{(i,j),(k,l) \in G \\ i,j,k,l \text{ all different}}} d_3(s, t) \\
&= (d_1(s) - 2d_2(s, t) + d_3(s, t))|G| + (d_2(s, t) - d_3(s, t)) \sum_{i=1}^n |G_i|^2 + d_3(s, t)|G|^2 \\
&= \frac{s(s-1)(n-t)(n-t-1)}{n(n-1)(n-2)(n-3)} \left(|G| + \frac{n-s-2}{s-1} \left(\sum_{i=1}^n |G_i|^2 + \frac{n-s-3}{s} |G|^2 \right) \right).
\end{aligned}$$

By plugging in and simplifying the expressions, we have

$$\begin{aligned} & \mathbf{E}[(R_1(s) - R_2(s))(R_1(t) - R_2(t))] - \mathbf{E}(R_1(s) - R_2(s))\mathbf{E}(R_1(t) - R_2(t)) \\ &= \frac{s(n-t)(\sum |G_i|^2 - 4|G|^2/n)}{n(n-1)}. \end{aligned}$$

Then,

$$\begin{aligned} \mathbf{Cov}(Z_{\text{diff}}(s), Z_{\text{diff}}(t)) &= \frac{\frac{s(n-t)(\sum |G_i|^2 - 4|G|^2/n)}{n(n-1)}}{\sqrt{\frac{s(n-s)}{n(n-1)}(\sum |G_i|^2 - \frac{4|G|^2}{n})\frac{t(n-t)}{n(n-1)}(\sum |G_i|^2 - \frac{4|G|^2}{n})}} \\ &= \frac{s(n-t)}{\sqrt{st(n-s)(n-t)}}. \end{aligned}$$

Hence, for $u \leq v$

$$\rho_{\text{diff}}^*(u, v) = \lim_{n \rightarrow \infty} \mathbf{Cov}(Z_{\text{diff}}(s), Z_{\text{diff}}(t)) = \frac{u(1-v)}{\sqrt{u(1-u)v(1-v)}}.$$

Similarly, for $u \geq v$, we have

$$\rho_{\text{diff}}^*(u, v) = \frac{v(1-u)}{\sqrt{v(1-v)u(1-u)}},$$

and the result in the proposition follows.

We can derive $\mathbf{Cov}(Z_w(s), Z_w(t))$ in a similar manner. Since

$$\mathbf{Cov}(Z_w(s), Z_w(t)) = \frac{\mathbf{E}[(R_w(s))(R_w(t)) - \mathbf{E}[R_w(s)]\mathbf{E}[R_w(t)]]}{\sqrt{\mathbf{Var}(R_w(s))\mathbf{Var}(R_w(t))}}$$

then after some simplification we have,

$$\begin{aligned} & \mathbf{Cov}(Z_w(s), Z_w(t)) \\ &= \frac{\frac{s(s-1)(n-t)(n-t-1)(|G|(n-1)(n-2) - \sum_{i=1}^n |G_i|^2(n-1) + 2|G|^2)}{n(n-1)^2(n-2)^2(n-3)}}{\frac{|G|(n-1)(n-2) - \sum_{i=1}^n |G_i|^2(n-1) + 2|G|^2}{n(n-1)^2(n-2)^2(n-3)}\sqrt{st(s-1)(t-1)(n-s)(n-t)(n-s-1)(n-t-1)}} \\ &= \frac{s(s-1)(n-t)(n-t-1)}{\sqrt{st(s-1)(t-1)(n-s)(n-t)(n-s-1)(n-t-1)}}. \end{aligned}$$

Hence, for $u \leq v$

$$\rho_w^*(u, v) = \lim_{n \rightarrow \infty} \mathbf{Cov}(Z_w(s), Z_w(t)) = \frac{u(1-v)}{v(1-u)}.$$

Similarly, for $u \geq v$, we have

$$\rho_w^*(u, v) = \frac{v(1-u)}{u(1-v)},$$

and the result in the proposition follows.

C.4. Proof of Proposition 4.4. To prove this proposition, we adopt Woodroffe's method (Woodroffe, 1976, 1978) by conditioning on the last cross-over and use the technique developed in Siegmund (1988). Let $S^*(u) = (Z_{\text{diff}}^*(u))^2 + (Z_w^*(u))^2$. Observe that $(S^*(u))^{1/2} = \sup_{0 \leq \omega < 2\pi} [Z_{\text{diff}}^*(u) \cos(\omega) + Z_w^*(u) \sin(\omega)]$. Then,

$$\begin{aligned} & \mathbf{P} \left(\max_{n_0 \leq t_1 \leq n_1} S^*(\frac{t_1}{n}) > b \right) \\ &= \sum_{n_0 \leq t_1 \leq n_1} \int_0^{2\pi} \int_0^\infty \mathbf{P}(S^*(\frac{t_1}{n}) \in b + dx, \arg(Z_{\text{diff}}^*(\frac{t_1}{n}), Z_w^*(\frac{t_1}{n})) \in d\omega) \\ & \quad \times \mathbf{P} \left(\max_{t_1 < t_2 \leq n_1} S^*(\frac{t_2}{n}) < b \mid S^*(\frac{t_1}{n}) = b + x, \arg(Z_{\text{diff}}^*(\frac{t_1}{n}), Z_w^*(\frac{t_1}{n})) = \omega \right) \end{aligned}$$

For the first probability in the product, we have

$$\mathbf{P}(S^*(\frac{t_1}{n}) \in b + dx, \arg(Z_{\text{diff}}^*(\frac{t_1}{n}), Z_w^*(\frac{t_1}{n})) \in d\omega) = \frac{1}{4\pi} e^{-(b+x)/2} dx d\omega.$$

For the second probability in the product, we have

$$\begin{aligned} & \mathbf{P} \left(\max_{t_1 < t_2 \leq n_1} S^*(\frac{t_2}{n}) < b \mid S^*(\frac{t_1}{n}) = b + x, Z_{\text{diff}}^*(\frac{t_1}{n}) = \sqrt{b} \cos(\omega) \right) \\ &= \mathbf{P} \left(\max_{t_1 < t_2 \leq n_1} (S^*(\frac{t_2}{n}) - S^*(\frac{t_1}{n})) < -x \mid S^*(\frac{t_1}{n}) = b + x, Z_{\text{diff}}^*(\frac{t_1}{n}) = \sqrt{b} \cos(\omega) \right). \end{aligned}$$

When $x = o(b)$, x is negligible to b . When $x = O(b)$, the first probability tends to 0. Hence,

$$\begin{aligned} & \mathbf{P} \left(\max_{n_0 \leq t_1 \leq n_1} S^*(t_1/n) > b \right) \\ &\approx \sum_{n_0 \leq t_1 \leq n_1} \int_0^{2\pi} \int_0^\infty \frac{1}{4\pi} e^{-(b+x)/2} \\ & \quad \times \mathbf{P} \left(\max_{t_1 < t_2 \leq n_1} (S^*(\frac{t_2}{n}) - S^*(\frac{t_1}{n})) < -x \mid S^*(\frac{t_1}{n}) = b, Z_{\text{diff}}^*(\frac{t_1}{n}) = \sqrt{b} \cos(\omega) \right) \\ & \quad dx d\omega. \end{aligned}$$

Let $u = t_1/n$, $v = t_2/n$. Then $u < v$. By chain rule,

$$S^{*'}(u) = 2Z_{\text{diff}}^*(u)Z_{\text{diff}}^{*'}(u) + 2Z_w^*(u)Z_w^{*'}(u),$$

For v close to u , $S^*(v) - S^*(u)$ can be approximated by

$$2Z_{\text{diff}}^*(u)(Z_{\text{diff}}^*(v) - Z_{\text{diff}}^*(u)) + 2Z_w^*(u)(Z_w^*(v) - Z_w^*(u)).$$

Notice that

$$\begin{aligned} & 2Z_{\text{diff}}^*(u)(Z_{\text{diff}}^*(v) - Z_{\text{diff}}^*(u))|(Z_{\text{diff}}^*(u) = \sqrt{b} \cos(\omega)) \\ & \sim \mathcal{N}((\rho_{\text{diff}}^*(u, v) - 1)2b \cos^2(\omega), (1 - (\rho_{\text{diff}}^*(u, v))^2)4b \cos^2(\omega)). \end{aligned}$$

Let $\delta = v - u$, by Taylor expansion,

$$\begin{aligned} \rho_{\text{diff}}^*(u, v) &= 1 + \lim_{v \searrow u} \frac{\partial \rho_{\text{diff}}^*(u, v)}{\partial v} \delta + O(\delta^2) = 1 - h_{\text{diff}}^*(u) \delta + O(\delta^2), \\ (\rho_{\text{diff}}^*(u, v))^2 &= 1 + 2 \lim_{v \searrow u} \frac{\partial \rho_{\text{diff}}^*(u, v)}{\partial u} \delta + O(\delta^2) = 1 - 2h_{\text{diff}}^*(u) \delta + O(\delta^2). \end{aligned}$$

For $\delta = O(n^{-1})$,

$$\begin{aligned} & 2Z_{\text{diff}}^*(u)(Z_{\text{diff}}^*(v) - Z_{\text{diff}}^*(u))|(Z_{\text{diff}}^*(u) = \sqrt{b} \cos(\omega)) \\ & \dot{\sim} \mathcal{N}(-2h_{\text{diff}}^*(u)b \cos^2(\omega)\delta, 8h_{\text{diff}}^*(u)b \cos^2(\omega)\delta), \end{aligned}$$

where $\dot{\sim}$ means approximated distributed as. Similarly, for $\delta = O(n^{-1})$,

$$\begin{aligned} & 2Z_w^*(u)(Z_w^*(v) - Z_w^*(u))|(Z_w^*(u) = \sqrt{b} \sin(\omega)) \\ & \dot{\sim} \mathcal{N}(-2h_w^*(u)b \sin^2(\omega)\delta, 8h_w^*(u)b \sin^2(\omega)\delta). \end{aligned}$$

Let $\mu(u) = 2b(h_{\text{diff}}^*(u) \cos^2(\omega) + h_w^*(u) \sin^2(\omega))$ and $\sigma^2(u) = 4\mu(u)$. Then, for $\delta = O(n^{-1})$,

$$S^*(v) - S^*(u) \dot{\sim} \mathcal{N}(-\mu(v)\delta, \sigma^2(v)\delta).$$

One can show that for $b = b_0 \sqrt{n}$ and $n \rightarrow \infty$,

$$\lim_{k \rightarrow \infty} \limsup_{n \rightarrow \infty} \sum_{|i-t| > k} \mathbf{P}(S^*(i/n) > b | S^*(t/n) = b + x) = 0.$$

Let $W_m^{(t_1)}$ be a random walk with $W_1^{(t_1)} \sim \mathcal{N}(\mu^{(t_1)}, (\sigma^{(t_1)})^2)$, where

$$\mu^{(t_1)} = \frac{2b}{n}(h_{\text{diff}}^*(u) \cos^2(\omega) + h_w^*(u) \sin^2(\omega)) \text{ and } (\sigma^{(t_1)})^2 = 4\mu^{(t_1)}.$$

Then

$$\begin{aligned} & \mathbf{P}\left(\max_{t_1 < t_2 < n_1} S^*\left(\frac{t_2}{n}\right) - S^*\left(\frac{t_1}{n}\right) < -x | S^*\left(\frac{t_1}{n}\right) = b, \arg(Z_{\text{diff}}^*\left(\frac{t_1}{n}\right), Z_w^*\left(\frac{t_2}{n}\right)) = \omega\right) \\ & \approx \mathbf{P}\left(\max_{t_1 < t_2 < n_1} -W_{t_2-t_1}^{(t_1)} < -x\right) \approx \mathbf{P}\left(\min_{m \geq 1} W_m^{(t_1)} > x\right). \end{aligned}$$

Together with the fact

$$\int_0^\infty \exp(-2\mu x/\sigma^2) \mathbf{P}(\min_{m \geq 1} W_m > x) dx = \mu \nu(2\mu/\sigma)$$

for a random walk $W_1 \sim \mathcal{N}(\mu, \sigma)$ with $\mu > 0$ (Siegmund (1992)), we have

$$\begin{aligned} & \mathbf{P}\left(\max_{n_0 \leq t_1 \leq n_1} S^*(t_1/n) > b\right) \\ & \approx \sum_{n_0 \leq t_1 \leq n_1} \frac{e^{-b/2}}{4\pi} \int_0^{2\pi} \mu^{(t_1)} \nu(\sqrt{\mu^{(t_1)}}) d\omega. \\ & = \frac{e^{-b/2}}{4\pi} \sum_{n_0 \leq t_1 \leq n_1} \int_0^{2\pi} 2b_0 u^*(t_1/n, \omega) \nu(\sqrt{2b_0 u^*(t_1/n, \omega)}) d\omega. \\ & \approx \frac{e^{-b/2}}{4\pi} \int_{x_0}^{x_1} \int_0^{2\pi} 2b_0 u^*(x, \omega) \nu(\sqrt{2b_0 u^*(x, \omega)}) d\omega n dx \\ & = \frac{e^{-b/2}}{4\pi} \int_{x_0}^{x_1} \int_0^{2\pi} 2b u^*(x, \omega) \nu(\sqrt{2b_0 u^*(x, \omega)}) d\omega dx, \end{aligned}$$

where $u^*(x, \omega) = h_{\text{diff}}^*(x) \cos^2(\omega) + h_w^*(x) \sin^2(\omega)$.

Next, we show for the changed interval statistic, which follows from Siegmund (1988, 1992). We include only key intermediate steps. Let

$$\rho_{\text{diff},(u_1, u_2)}^*(\delta_1, \delta_2) := \mathbf{Cov}(Z_{\text{diff}}^*(u_1 + \delta_1, u_2 + \delta_2), Z_{\text{diff}}^*(u_1, u_2))$$

$$\rho_{w,(u_1, u_2)}^*(\delta_1, \delta_2) := \mathbf{Cov}(Z_w^*(u_1 + \delta_1, u_2 + \delta_2), Z_w^*(u_1, u_2)).$$

Given that they are both differentiable with the derivative being continuous except at $\delta_1 = 0$ and $\delta_2 = 0$, a key intermediate form is:

$$\begin{aligned} & \mathbf{P}\left(\max_{n_0 \leq t_2 - t_1 \leq n_1} S^*(t_1/n, t_2/n) > b\right) \approx \\ & \sum_{n_0 \leq t_2 - t_1 \leq n_1} \frac{b^2 e^{-\frac{b}{2}}}{\pi} \int_0^{2\pi} C_1(t_1, t_2) C_2(t_1, t_2) \nu(\sqrt{2b C_1(t_1, t_2)}) \nu(\sqrt{2b C_2(t_1, t_2)}) d\omega \end{aligned}$$

where C_1, C_2 are the partial derivatives

$$\begin{aligned} C_1(nu_1, nu_2) &= \frac{\cos^2(\omega)}{n} \frac{\partial \rho_{\text{diff},(u_1, u_2)}^*(\delta_1, 0)}{\partial \delta_1} \Big|_{\delta_1=0} + \frac{\sin^2(\omega)}{n} \frac{\partial \rho_{w,(u_1, u_2)}^*(\delta_1, 0)}{\partial \delta_1} \Big|_{\delta_1=0}, \\ C_2(nu_1, nu_2) &= \frac{\cos^2(\omega)}{n} \frac{\partial \rho_{\text{diff},(u_1, u_2)}^*(0, \delta_2)}{\partial \delta_2} \Big|_{\delta_2=0} + \frac{\sin^2(\omega)}{n} \frac{\partial \rho_{w,(u_1, u_2)}^*(0, \delta_2)}{\partial \delta_2} \Big|_{\delta_2=0}. \end{aligned}$$

Under the permutation null, the processes derived from the perturbation of the left and right end points,

$$S^*((t_1 + k)/n, t_2/n), k = \dots, -2, -1, 0, 1, 2, \dots$$

and

$$S^*(t_1/n, (t_2 - k)/n), k = \dots, -2, -1, 0, 1, 2, \dots$$

are identical in distribution the process

$$S^*((t_2 - t_1 - k)/n), k = \dots, -2, -1, 0, 1, 2, \dots$$

Thus, the partial derivatives are equal to the derivative in the one change-point scenario,

$$C_1(t_1, t_2) = C_2(t_1, t_2) = \frac{1}{n} (h_{\text{diff}}^*(u_2 - u_1) \cos^2(\omega) + h_w^*(u_2 - u_1) \sin^2(\omega)),$$

with $u_1 = t_1/n, u_2 = t_2/n$. This leads to

$$\begin{aligned} \mathbf{P} \left(\max_{n_0 \leq t_2 - t_1 \leq n_1} S^*(t_1/n, t_2/n) > b \right) \\ \approx \frac{e^{-b/2}}{4\pi} 4b^2 \int_{x_0}^{x_1} \int_0^{2\pi} \left(u^*(x, \omega) \nu(\sqrt{2b_0 u^*(x, \omega)}) \right)^2 (1-x) d\omega dx. \end{aligned}$$

SUPPLEMENT D: CHECKING ANALYTIC P -VALUE APPROXIMATIONS

In this supplement, we present the complete results on checking analytic p -value approximations for the single change-point alternative (D.1) and changed interval alternative (D.2).

D.1. Single change-point alternative. Here, we present the critical values obtained from (4.15), (4.17), (4.19), (4.24), and (4.26) to the critical values obtained from doing 10,000 permutations directly, under various simulation settings for the single change-point alternative. Tables D.1, D.2, and D.3 are the complete versions of Tables 2, 3, and 4, respectively, presented in Section 4.4.

In each simulation, sequences of length 1,000 were generated from a given distribution F_0 in $\mathbb{R}^d$. We considered three distributions (multivariate normal, multivariate t with 5 degrees of freedom, and multivariate log-normal)

under various dimensions ($d = 10$, $d = 100$, and $d = 1000$). The analytical approximations depend on constraints on the sequence in which the change-point is searched over (n_0 and n_1). To make things simple, we let $n_1 = n - n_0$.

On the other hand, the skewness corrected p -value approximations and permutation p -values depend on certain characteristics of the structure of the graph G . In this simulation, the MST is used. As the structure of MST depends on the observations, the critical value vary by simulation runs. We show results for 4 randomly simulated sequences in each setting. Two characteristics of the graph are also reported: the sum of squared node degrees ($\sum_i |G_i|^2$) and the maximum node degree ($\max_i |G_i|$). These quantities give some intuitions on the size and density of the hubs in the graph. The notation is identical to that in described in Section 4.4.

D.2. Changed interval alternative. Tables D.4, D.5, and D.6 show the results of the p -value approximations for the changed interval alternative. The notation and simulation settings are identical to those for the single change-point alternative in Section 4.4 and Section D.1, except that n_0 is replaced by l_0 for the smallest window size (l_1 is set to $n - l_0$).

From the tables, conclusions similar to those for the single change-point alternative can be drawn. The analytical approximation after skewness correction performs much better than the analytic approximation, especially when the window size decreases. As the window size decreases, the skewness-corrected p -value approximations still do quite well when the dimension is not too high ($d = 10$, $d = 100$) and $n_0 \geq 50$. When the $d = 1000$, the skewness-corrected p -values are still doing pretty well for $n_0 \geq 75$ in general.

TABLE D.1
Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} S(t)$ based on MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$		
A1	13.10	13.38	13.70	14.11		

	Critical Values				Graph	
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	$\sum G_i ^2$	$d_{\max}$
	Per	Per	Per	Per		
$N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	13.02	13.41	14.20	15.51	5518	10
	12.98	13.34	13.94	15.38	5442	8
	12.87	13.29	14.04	15.17	5394	8
	13.02	13.42	13.71	15.65	5368	8
Multivariate t , 5 df $d = 10$	12.73	13.39	13.83	15.17	5336	9
	13.20	13.34	13.81	15.39	5358	7
	13.13	13.44	13.98	15.45	5346	8
	12.91	13.44	13.84	15.20	5326	8
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	12.93	13.35	13.92	15.26	5106	8
	13.11	13.13	13.90	15.12	5020	7
	13.07	13.29	13.66	15.17	5022	6
	13.02	13.34	13.89	15.62	5098	9
$N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	13.07	13.86	15.15	17.71	9852	40
	13.08	13.79	14.82	17.17	9888	29
	13.48	14.28	15.72	18.49	13106	49
	13.48	13.78	14.53	17.17	9500	30
Multivariate t , 5 df $d = 100$	13.47	14.20	15.48	17.81	14302	42
	13.32	13.77	14.96	17.11	12424	39
	13.94	14.82	16.03	19.07	18574	64
	13.22	14.00	15.08	17.36	11266	31
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	13.42	13.82	14.89	17.67	9482	31
	13.31	14.19	15.54	18.21	10056	44
	13.18	13.68	14.65	16.43	8526	27
	13.32	13.72	14.76	16.46	8442	24
$N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	13.78	14.39	16.37	19.49	17560	70
	13.87	14.77	16.19	19.86	17866	67
	13.50	13.94	15.73	18.36	15364	46
	13.24	14.21	15.15	17.14	12772	36
Multivariate t , 5 df $d = 1000$	14.11	14.91	17.03	20.73	30180	92
	14.04	15.09	17.05	20.67	29556	77
	15.14	17.52	20.04	28.66	51128	179
	14.12	14.68	16.62	20.06	24128	69
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	14.99	16.81	19.26	25.74	63580	155
	16.12	18.38	22.00	29.07	106524	208
	14.50	15.83	18.14	21.96	46876	83
	16.81	19.13	23.42	33.46	117598	286

TABLE D.2
Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} Z_w(t)$ based on MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$		$n_0 = 75$		$n_0 = 50$		$n_0 = 25$			
A1	2.98		3.02		3.08		3.14			

	Critical Values								Graph	
	$n_0 = 100$		$n_0 = 75$		$n_0 = 50$		$n_0 = 25$		$\sum G_i ^2$	$d_{\max}$
	A2	Per	A2	Per	A2	Per	A2	Per		
$N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	3.05	3.02	3.12	3.11	3.22	3.22	3.4	3.48	5518	10
	3.05	3.05	3.12	3.14	3.22	3.25	3.4	3.45	5442	8
	3.05	3.05	3.12	3.14	3.22	3.27	3.4	3.53	5394	8
	3.05	3.03	3.12	3.10	3.22	3.23	3.4	3.50	5368	8
Multivariate t , 5 df $d = 10$	3.05	3.06	3.12	3.15	3.22	3.25	3.4	3.49	5336	9
	3.05	3.06	3.12	3.10	3.22	3.23	3.4	3.51	5358	7
	3.05	3.07	3.12	3.11	3.22	3.27	3.4	3.47	5346	8
	3.05	3.05	3.12	3.15	3.22	3.22	3.4	3.45	5326	8
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	3.05	3.02	3.12	3.12	3.22	3.22	3.4	3.45	5106	8
	3.05	3.01	3.12	3.13	3.22	3.25	3.4	3.50	5020	7
	3.05	3.05	3.12	3.11	3.22	3.26	3.4	3.52	5022	6
	3.05	3.01	3.12	3.12	3.22	3.21	3.4	3.47	5098	9
$N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	3.05	3.06	3.12	3.13	3.22	3.23	3.39	3.60	9852	40
	3.05	3.03	3.12	3.12	3.22	3.25	3.39	3.52	9888	29
	3.05	3.09	3.12	3.13	3.22	3.28	3.39	3.58	13106	49
	3.05	3.01	3.12	3.14	3.22	3.27	3.39	3.56	9500	30
Multivariate t , 5 df $d = 100$	3.05	3.04	3.12	3.15	3.22	3.31	3.39	3.62	14302	42
	3.05	3.06	3.12	3.13	3.22	3.29	3.39	3.54	12424	39
	3.05	3.05	3.12	3.21	3.21	3.33	3.39	3.65	18574	64
	3.05	3.07	3.12	3.16	3.22	3.24	3.39	3.57	11266	31
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	3.05	3.07	3.12	3.11	3.22	3.27	3.39	3.57	9482	31
	3.05	3.10	3.12	3.15	3.22	3.28	3.39	3.50	10056	44
	3.05	3.07	3.12	3.12	3.22	3.24	3.39	3.49	8526	27
	3.05	3.04	3.12	3.11	3.22	3.26	3.39	3.55	8442	24
$N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	3.05	3.08	3.12	3.16	3.22	3.31	3.39	3.59	17560	70
	3.05	3.05	3.12	3.13	3.22	3.29	3.39	3.63	17866	67
	3.05	3.08	3.12	3.12	3.22	3.25	3.39	3.62	15364	46
	3.05	3.07	3.12	3.17	3.22	3.31	3.39	3.59	12772	36
Multivariate t , 5 df $d = 1000$	3.05	3.08	3.12	3.20	3.21	3.33	3.39	3.71	30180	92
	3.04	3.08	3.12	3.21	3.21	3.33	3.39	3.73	29556	77
	3.04	3.11	3.11	3.20	3.21	3.42	3.38	3.81	51128	179
	3.05	3.06	3.12	3.17	3.21	3.33	3.39	3.70	24128	69
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	3.04	3.14	3.11	3.29	3.20	3.47	3.37	3.89	63580	155
	3.03	3.20	3.10	3.40	3.19	3.61	3.35	3.99	106524	208
	3.04	3.11	3.11	3.25	3.21	3.37	3.38	3.82	46876	83
	3.03	3.19	3.10	3.35	3.19	3.56	3.35	4.03	117598	286

TABLE D.3

Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} M(t)$ based on MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
A1	3.23	3.27	3.32	3.38

	Critical Values								Graph	
	$n_0 = 100$		$n_0 = 75$		$n_0 = 50$		$n_0 = 25$		$\sum G_i ^2$	$d_{\max}$
	A2	Per	A2	Per	A2	Per	A2	Per		
$N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	3.27	3.26	3.33	3.34	3.41	3.42	3.56	3.66	5518	10
	3.27	3.29	3.33	3.34	3.41	3.44	3.56	3.67	5442	8
	3.27	3.29	3.33	3.33	3.41	3.43	3.56	3.64	5394	8
	3.27	3.30	3.33	3.36	3.41	3.44	3.55	3.66	5368	8
Multivariate t , 5 df $d = 10$	3.27	3.27	3.33	3.32	3.41	3.45	3.56	3.68	5336	9
	3.27	3.26	3.33	3.34	3.41	3.43	3.55	3.64	5358	7
	3.27	3.31	3.33	3.31	3.41	3.44	3.55	3.63	5346	8
	3.27	3.27	3.33	3.36	3.41	3.41	3.56	3.67	5326	8
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	3.27	3.28	3.33	3.34	3.41	3.45	3.55	3.63	5106	8
	3.27	3.29	3.33	3.36	3.41	3.41	3.55	3.62	5020	7
	3.27	3.27	3.33	3.33	3.41	3.44	3.55	3.63	5022	6
	3.27	3.28	3.33	3.34	3.41	3.45	3.55	3.61	5098	9
$N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	3.30	3.32	3.37	3.40	3.47	3.55	3.66	3.92	9852	40
	3.28	3.31	3.35	3.40	3.45	3.53	3.62	3.83	9888	29
	3.31	3.33	3.39	3.44	3.50	3.57	3.69	4.00	13106	49
	3.28	3.32	3.35	3.36	3.45	3.51	3.62	3.80	9500	30
Multivariate t , 5 df $d = 100$	3.30	3.33	3.38	3.44	3.48	3.55	3.67	3.89	14302	42
	3.29	3.31	3.36	3.40	3.46	3.54	3.64	3.85	12424	39
	3.33	3.30	3.41	3.44	3.53	3.64	3.73	4.04	18574	64
	3.29	3.26	3.36	3.40	3.46	3.53	3.64	3.79	11266	31
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	3.29	3.29	3.37	3.41	3.47	3.50	3.65	3.82	9482	31
	3.32	3.33	3.40	3.42	3.51	3.60	3.71	3.89	10056	44
	3.28	3.27	3.35	3.38	3.45	3.45	3.62	3.80	8526	27
	3.28	3.30	3.35	3.37	3.45	3.54	3.62	3.76	8442	24
$N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	3.33	3.34	3.42	3.45	3.53	3.64	3.74	4.05	17560	70
	3.33	3.37	3.41	3.45	3.53	3.62	3.73	4.07	17866	67
	3.31	3.32	3.38	3.40	3.49	3.59	3.68	3.95	15364	46
	3.29	3.30	3.35	3.39	3.45	3.54	3.63	3.83	12772	36
Multivariate t , 5 df $d = 1000$	3.33	3.34	3.41	3.44	3.53	3.68	3.73	4.13	30180	92
	3.33	3.39	3.41	3.49	3.53	3.67	3.73	4.13	29556	77
	3.43	3.42	3.54	3.64	3.69	3.98	3.95	4.84	51128	179
	3.32	3.33	3.40	3.43	3.51	3.64	3.71	4.04	24128	69
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	3.37	3.43	3.47	3.61	3.60	3.87	3.83	4.47	63580	155
	3.39	3.51	3.49	3.75	3.63	4.06	3.88	4.58	106524	208
	3.33	3.34	3.41	3.49	3.53	3.69	3.74	4.22	46876	83
	3.44	3.50	3.55	3.71	3.71	4.15	3.97	5.05	117598	286

TABLE D.4
Critical values for the single change-point scan statistic $\max_{l_0 \leq t_2 - t_1 \leq l_1} S(t)$ based on MST at 0.05 significance level. $n = 1000$.

	$l_0 = 100$	$l_0 = 75$	$l_0 = 50$	$l_0 = 25$
A1	22.83	23.33	23.96	24.90

	Critical Values				Graph	
	$l_0 = 100$	$l_0 = 75$	$l_0 = 50$	$l_0 = 25$	$\sum G_i ^2$	$d_{\max}$
	Per	Per	Per	Per		
$N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	21.58	22.91	26.51	37.07	5496	8
	21.65	23.14	26.08	36.48	5432	8
	21.54	23.14	26.23	36.89	5524	8
	21.40	22.95	26.08	36.70	5432	9
Multivariate t , 5 df $d = 10$	21.25	23.17	26.27	37.07	5378	8
	21.39	22.79	26.31	36.26	5416	7
	21.54	22.89	25.69	36.77	5334	8
	21.37	22.58	25.98	36.60	5356	8
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	21.15	22.69	25.53	35.50	5066	8
	21.47	22.65	25.51	35.65	5082	7
	21.04	22.83	25.78	35.35	5068	8
	21.15	22.80	25.57	35.95	5068	6
$N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	24.41	27.59	33.79	52.29	11634	57
	23.63	26.95	32.96	50.49	11098	50
	23.51	25.88	30.81	45.21	9992	24
	22.90	24.83	28.68	43.05	9250	21
Multivariate t , 5 df $d = 100$	24.32	27.07	33.34	49.99	13908	41
	24.91	28.47	35.80	56.96	16786	64
	26.10	30.78	39.52	66.40	22336	97
	24.69	28.02	33.95	52.33	16344	40
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	24.51	27.18	33.52	51.18	10384	43
	23.41	25.99	30.78	46.20	9110	30
	24.05	26.92	32.62	49.33	9822	42
	23.23	26.11	31.05	47.32	9264	34
$N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	25.64	29.44	36.89	57.03	21204	69
	25.59	29.35	36.11	56.07	18710	67
	25.87	30.21	38.54	62.41	19402	78
	24.02	27.01	33.19	49.58	14076	47
Multivariate t , 5 df $d = 1000$	28.77	34.29	44.96	75.38	52082	145
	25.93	29.93	38.09	60.99	27188	69
	29.89	35.82	46.94	79.97	52498	162
	27.09	31.66	40.77	65.00	30584	101
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	29.07	34.62	45.10	75.23	49088	143
	36.74	46.61	65.84	121.42	173558	380
	31.06	38.47	53.10	92.49	75060	223
	31.69	39.44	54.75	95.64	80592	234

TABLE D.5

Critical values for the changed interval change-point scan statistic $\max_{l_0 \leq t_2 - t_1 \leq l_1} Z_w(t)$ based on MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
A1	4.08	4.14	4.22	4.33

	Critical Values								Graph	
	$n_0 = 100$		$n_0 = 75$		$n_0 = 50$		$n_0 = 25$		$\sum G_i ^2$	$d_{\max}$
	A2	Per	A2	Per	A2	Per	A2	Per		
$N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	4.29	4.27	4.47	4.48	4.77	4.85	5.46	5.85	5404	8
	4.29	4.27	4.47	4.46	4.77	4.85	5.46	5.83	5436	10
	4.29	4.28	4.47	4.46	4.77	4.81	5.46	5.90	5372	7
	4.29	4.26	4.47	4.47	4.77	4.81	5.46	5.85	5458	8
Multivariate t , 5 df $d = 10$	4.29	4.27	4.47	4.47	4.77	4.84	5.46	5.85	5390	8
	4.29	4.26	4.47	4.48	4.77	4.81	5.46	5.85	5360	8
	4.29	4.25	4.47	4.46	4.77	4.83	5.46	5.83	5348	9
	4.29	4.25	4.47	4.47	4.77	4.85	5.46	5.89	5326	7
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	4.29	4.27	4.47	4.44	4.77	4.83	5.46	5.82	5078	6
	4.29	4.25	4.47	4.45	4.77	4.81	5.46	5.84	5078	8
	4.29	4.23	4.47	4.44	4.77	4.84	5.46	5.85	5044	7
	4.29	4.26	4.47	4.47	4.77	4.84	5.46	5.85	5018	7
$N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	4.29	4.30	4.46	4.56	4.76	5.02	5.45	6.31	12576	38
	4.29	4.30	4.46	4.51	4.76	4.94	5.45	6.13	9796	27
	4.29	4.31	4.46	4.56	4.76	5.06	5.44	6.46	13042	50
	4.29	4.31	4.46	4.57	4.76	4.96	5.45	6.12	10616	44
Multivariate t , 5 df $d = 100$	4.29	4.37	4.46	4.71	4.75	5.24	5.43	6.76	21738	96
	4.29	4.35	4.46	4.65	4.76	5.13	5.44	6.51	15074	45
	4.29	4.34	4.46	4.61	4.76	5.13	5.44	6.54	16122	49
	4.29	4.39	4.46	4.62	4.75	5.15	5.44	6.56	17584	68
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	4.29	4.29	4.46	4.50	4.76	4.92	5.45	6.08	8688	27
	4.29	4.33	4.46	4.54	4.76	4.99	5.45	6.22	10260	35
	4.29	4.34	4.46	4.58	4.76	5.06	5.44	6.29	11976	54
	4.29	4.28	4.46	4.56	4.76	4.94	5.45	6.21	9692	41
$N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	4.29	4.34	4.46	4.58	4.76	5.07	5.44	6.41	14052	40
	4.28	4.43	4.45	4.75	4.75	5.38	5.43	6.88	24632	101
	4.29	4.32	4.46	4.60	4.76	5.05	5.44	6.36	13690	42
	4.29	4.36	4.46	4.61	4.75	5.14	5.44	6.56	18774	62
Multivariate t , 5 df $d = 1000$	4.28	4.47	4.45	4.81	4.74	5.47	5.42	7.16	33052	106
	4.28	4.40	4.45	4.71	4.75	5.30	5.43	6.85	24644	60
	4.29	4.38	4.46	4.71	4.75	5.23	5.43	6.73	21496	52
	4.28	4.50	4.45	4.87	4.74	5.54	5.41	7.17	35786	119
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	4.20	5.17	4.35	5.79	4.60	6.94	5.19	9.39	194376	378
	4.28	4.49	4.45	4.80	4.74	5.44	5.42	7.06	31896	80
	4.28	4.59	4.44	4.95	4.73	5.73	5.39	7.62	48656	129
	4.26	4.71	4.43	5.08	4.71	6.03	5.37	8.02	65810	139

TABLE D.6
Critical values for the changed interval change-point scan statistic $\max_{l_0 \leq t_2 - t_1 \leq l_1} M(t)$
based on MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
A1	4.21	4.27	4.34	4.45

	Critical Values								Graph	
	$n_0 = 100$		$n_0 = 75$		$n_0 = 50$		$n_0 = 25$		$\sum G_i ^2$	$d_{\max}$
	A2	Per	A2	Per	A2	Per	A2	Per		
$N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	4.40	4.32	4.56	4.50	4.83	4.85	5.48	5.85	5392	8
	4.40	4.35	4.56	4.54	4.83	4.88	5.47	5.85	5474	8
	4.41	4.33	4.56	4.51	4.84	4.85	5.48	5.82	5392	9
	4.41	4.33	4.56	4.52	4.83	4.87	5.48	5.88	5444	8
Multivariate t , 5 df $d = 10$	4.40	4.33	4.56	4.54	4.83	4.89	5.47	5.88	5344	8
	4.40	4.36	4.56	4.51	4.83	4.84	5.48	5.86	5296	8
	4.42	4.34	4.58	4.52	4.85	4.90	5.49	5.92	5428	11
	4.40	4.34	4.56	4.51	4.83	4.86	5.48	5.85	5428	7
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 10$	4.39	4.34	4.55	4.52	4.82	4.83	5.47	5.83	5010	6
	4.39	4.34	4.55	4.50	4.82	4.81	5.47	5.85	5020	6
	4.39	4.31	4.55	4.49	4.82	4.84	5.47	5.82	5026	7
	4.39	4.34	4.54	4.52	4.81	4.84	5.47	5.82	5064	7
$N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	4.59	4.47	4.79	4.69	5.11	5.16	5.74	6.37	10578	33
	4.52	4.44	4.71	4.67	5.01	5.05	5.63	6.20	9228	22
	4.65	4.48	4.86	4.74	5.20	5.22	5.84	6.43	10738	42
	4.70	4.53	4.93	4.77	5.28	5.29	5.94	6.59	13276	48
Multivariate t , 5 df $d = 100$	4.55	4.47	4.74	4.69	5.04	5.19	5.80	6.36	10820	28
	4.88	4.59	5.16	4.98	5.56	5.55	6.28	6.98	19662	66
	4.74	4.55	4.98	4.83	5.34	5.36	6.01	6.71	15250	59
	4.66	4.55	4.88	4.83	5.22	5.33	5.86	6.54	13820	47
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 100$	4.58	4.43	4.79	4.69	5.10	5.13	5.73	6.24	8690	32
	4.58	4.46	4.78	4.71	5.09	5.13	5.72	6.27	9416	28
	4.53	4.42	4.72	4.66	5.01	5.06	5.63	6.15	8028	26
	4.51	4.40	4.69	4.63	4.98	5.05	5.60	6.12	8400	22
$N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	4.78	4.60	5.03	4.88	5.41	5.40	6.09	6.78	16810	64
	4.78	4.57	5.03	4.87	5.41	5.21	6.09	6.80	16168	61
	4.57	4.47	4.77	4.69	5.09	5.18	5.71	6.38	11748	32
	4.85	4.61	5.12	4.94	5.52	5.54	6.23	6.95	22126	81
Multivariate t , 5 df $d = 1000$	5.44	4.76	5.82	5.24	6.35	6.20	7.24	8.29	68100	211
	4.97	4.69	5.27	5.09	5.70	5.75	6.45	7.37	33762	107
	4.79	4.66	5.05	5.01	5.43	5.62	6.11	7.16	29350	80
	4.65	4.53	4.87	4.82	5.20	5.38	5.84	6.73	20578	52
$\ln N_d(\mathbf{0}, \mathbf{I})$ $d = 1000$	5.19	4.91	5.53	5.41	6.01	6.22	6.83	8.16	65522	165
	5.34	4.88	5.71	5.39	6.22	6.26	7.08	8.38	82076	220
	5.74	5.22	6.17	5.82	6.75	6.98	7.72	9.37	206368	419
	5.36	4.83	5.73	5.60	6.25	6.16	7.12	8.15	71082	212

SUPPLEMENT E: POWER TABLES

Here, we present additional simulation settings to examine the performance of the three new test statistics. In each simulation, we generated a sequence of $n = 200$ observations for various dimensions d from the following model:

$$\mathbf{y}_t \sim \begin{cases} F_0, & t = 1, \dots, \tau; \\ F_1, & t = \tau + 1, \dots, 200. \end{cases}$$

Here, τ is the change-point. When there is a mean difference, we use Δ to denote the Euclidean distance of the means of F_0 and F_1 . When there is a variance difference, to make the change less significant, only the first $\lfloor d/5 \rfloor$ of the diagonal elements of the covariance matrix differ with a multiple of σ , and the rest are unchanged. In the following simulation settings, for simplicity, we set the graph to be the 5-MST constructed using Euclidean distance.

Under the Gaussian setting, if one assumes that, at the change-point, only the mean (μ) may change, the scan statistic over Hotelling's T^2 statistics can be used: $\max_{n_0 \leq t \leq n_1} HT(t)$, with $HT(t) = \frac{t(n-t)}{n} (\bar{\mathbf{y}}_t - \bar{\mathbf{y}}_t^*)^T \tilde{\Sigma}_t^{-1} (\bar{\mathbf{y}}_t - \bar{\mathbf{y}}_t^*)$ where $\bar{\mathbf{y}}_t = \sum_{i=1}^t \mathbf{y}_i / t$, $\bar{\mathbf{y}}_t^* = \sum_{i=t+1}^n \mathbf{y}_i / (n-t)$, and $\tilde{\Sigma}_t = (\sum_{i=1}^t (\mathbf{y}_i - \bar{\mathbf{y}}_t)(\mathbf{y}_i - \bar{\mathbf{y}}_t)^T + \sum_{i=t+1}^n (\mathbf{y}_i - \bar{\mathbf{y}}_t^*)(\mathbf{y}_i - \bar{\mathbf{y}}_t^*)^T) / (n-2)$. If the variance may also change at the change-point, the scan statistic over the generalized likelihood ratio statistic can be used: $\max_{n_0 \leq t \leq n_1} GLR(t)$ with $GLR(t) = n \log |\hat{\Sigma}_n| - t \log |\hat{\Sigma}_t| - (n-t) \log |\hat{\Sigma}_t^*|$, where $\hat{\Sigma}_t = \frac{\sum_{i=1}^t (\mathbf{y}_i - \bar{\mathbf{y}}_t)(\mathbf{y}_i - \bar{\mathbf{y}}_t)^T}{t}$ and $\hat{\Sigma}_t^* = \frac{\sum_{i=t+1}^n (\mathbf{y}_i - \bar{\mathbf{y}}_t^*)(\mathbf{y}_i - \bar{\mathbf{y}}_t^*)^T}{n-t}$.

The performance of six methods are compared: two methods based on normal theory ($\max HT(t)$, $\max GLR(t)$), the method in [Chen and Zhang \(2015\)](#) ($\max Z(t)$), and three new tests ($\max S(t)$, $\max R_w(t)$, $\max M(t)$). The estimated power is calculated as the number of trials, out of 100, that the null hypothesis is rejected at 0.05 level for each of these methods, with p -values determined by 10,000 permutation runs for fairness in comparison. To examine the accuracy of the estimated change-point, the number of trials where the estimated change-point is within 20 from the true change-point is provided in parentheses. Under each setting, the specific alternative is chosen so that the tests have moderate power to be comparable. The best one for each scenario is made bold. In the following, we use 'HT' to refer to the scan statistic over the Hotelling's T^2 statistic and use 'GLR' to refer to the scan statistic over the generalized likelihood ratio statistic.

When there is scale change only (Table [E.1](#)) and the data is multivariate Gaussian, HT has no power as there is no mean change. GLR dominates in

power when $d = 10$, but starting at $d = 20$, the graph-based methods exceed GLR in power. Among the graph-based methods, $S(t)$ and $M(t)$ have much higher power, with $M(t)$ slightly outperforming $S(t)$ as the dimension increases. On the other hand, this setting is not well-suited for $Z_w(t)$ and its power suffers as d increases. $Z(t)$ has reasonable power across all d but struggles to accurately estimate the change-point (the numbers in the parenthesis are much smaller).

Other distributions are considered as well. Tables E.2 and E.3 show results from multivariate log-normal distribution. For the log-normal distribution, a change in mean difference leads to a change in location and scale. When the change is at the center of the sequence, the method in Chen and Zhang (2015) and $Z_w(t)$ have comparable performance. However, once the change is no longer at the center of the sequence, the method in Chen and Zhang (2015) fails badly in estimating the location of the change-point for $d \geq 50$. In this particular parameter choice, $Z_w(t)$ outperforms the rest but $S(t)$ and $M(t)$ are also on par.

TABLE E.1
Multivariate Gaussian data, scale difference, τ at center

d	10	20	50	75	500	2000
σ	1.7	1.5	1.3	1.28	1.1	1.05
<i>HT</i>	6 (2)	9 (2)	6 (2)	2 (2)	-	-
GLR	72 (63)	15 (3)	8 (0)	-	-	-
<i>Z</i>	71 (53)	61 (45)	51 (17)	64 (30)	42 (15)	52 (30)
Z_w	60 (54)	46 (35)	22 (8)	25 (16)	7 (4)	6 (1)
<i>S</i>	69 (60)	70 (56)	60 (39)	80 (62)	57 (50)	73 (58)
<i>M</i>	59 (48)	64 (47)	65 (49)	81 (62)	68 (57)	83 (65)

TABLE E.2
Multivariate Log-normal data, mean difference, τ at center

d	10	50	100	150	175	500	2000
Δ	1	1.2	1.4	1.5	1.5	1.6	2
<i>HT</i>	78 (65)	62 (48)	29 (23)	16 (10)	9 (6)	-	-
GLR	13 (9)	6 (0)	-	-	-	-	-
<i>Z</i>	86 (79)	83 (80)	88 (87)	91 (91)	91 (91)	81 (81)	87 (86)
Z_w	81 (62)	81 (70)	88 (75)	95 (81)	90 (79)	75 (62)	83 (68)
<i>S</i>	72 (57)	73 (67)	78 (69)	86 (70)	82 (69)	60 (49)	73 (59)
<i>M</i>	80 (61)	79 (69)	83 (74)	91 (77)	85 (75)	64 (52)	78 (64)

TABLE E.3
Multivariate Log-normal data, mean difference, τ at three quarters

d	10	50	100	150	175	500	2000
Δ	1	1.2	1.4	1.5	1.5	1.6	2
HT	84 (72)	69 (64)	43 (39)	24 (16)	6 (3)	-	-
GLR	14 (6)	14 (10)	-	-	-	-	-
Z	62 (56)	45 (23)	44 (17)	40 (6)	38 (6)	28 (2)	39 (2)
Z_w	61 (55)	64 (56)	81 (75)	76 (68)	71 (64)	53 (49)	64 (56)
S	59 (52)	56 (50)	69 (63)	66 (60)	63 (58)	42 (38)	51 (47)
M	59 (53)	58 (51)	69 (64)	75 (68)	68 (62)	48 (43)	58 (52)

SUPPLEMENT F: ADDITIONAL ILLUSTRATIONS OF THE NYC
TAXI DATASET

(a) Oct. 3, 2015

(b) Oct. 18, 2015

(c) Oct. 29, 2015

(d) Nov. 06, 2015

FIG 1. *Density heatmap of taxi drop-offs for ten selected days.*

SUPPLEMENT G: MORE DISCUSSIONS ON THE CONDITIONS

G.1. Critical Values for k -MST. The conditions on $|A_e|$ and $|B_e|$ stated in Theorem 4.1 are sufficient conditions, but not a necessity. It may be that in practice, these conditions are violated. For example, for k -MST where $k = O(1)$, the conditions on A_e and B_e boil down to one: $\sum_{e \in G} |A_e||B_e| = o(n^{1.5})$. Suppose that the largest hub, u , is of order n^β , then $\sum_{e \in G} \geq \sum_{v \in G_u} |A_{(u,v)}||B_{(u,v)}| \geq \sum_{v \in G_u} n^\beta n^\beta = n^{3\beta}$. Hence, $\beta < 0.5$. For $n = 1000$, $n^{0.5} \approx 32$.

Table G.1 lists the critical values for the generalized edge-count test obtained from asymptotic results and from doing 10,000 random permutations under three different settings: (1) multivariate standard normal with $d = 1000$, (2) multivariate t_5 with $d = 1000$, and (3) multivariate log-normal with $d = 1000$. The largest degree ($d_{\max}$) are all larger than 32. Some are much larger, such as for the 10-MST under the Multivariate log-normal distribution, the largest node degree is 768. We also directly computed $\sum_{e \in G} |A_e||B_e|/n^{1.5}$ and this ratio is way larger than 1, indicating explicitly that the conditions are severely violated. Even under such severe violations, when n_0 is relatively large ($n_0 = 100$), the asymptotic p -value formula still give us pretty reasonable critical values. Similar conclusions holds for the weighted edge-count test (Table G.2) and the max-type edge-count test (Table G.3).

When n_0 is small (< 100), the issue that the convergence of the marginal distribution of the test statistic to the normal distribution is slow for t close to 1 or n becomes prominent, causing the critical values obtained from the asymptotic formulas to not be that accurate. While if we adopt the skewness correction, the analytic formulas give us pretty reasonable values even for n_0 as small as 50 and when the conditions on the similarity graph are severely violated (Tables G.2 and G.3).

G.2. Critical Values for k -NNG. Here, we present additional results on checking the critical values obtained from the analytic p -value formulas when the conditions on the similarity graph in the limiting theorem are violated. Rather than using k -MST, here we use k -NNG in constructing the graph, and conclusions are similar to those in drawn from Supplement G.1. Detailed results can be seen in Tables G.4, G.5, and G.6. (The distributions we consider here are the same as those in Supplement G.1: (1) multivariate standard normal with $d = 1000$, (2) multivariate t_5 with $d = 1000$, and (3) multivariate log-normal with $d = 1000$.)

TABLE G.1
Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} S(t)$ based on k -MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
Asy	13.10	13.38	13.70	14.11

	Critical Values				Graph		
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	k	$\frac{\sum_e A_e B_e }{n^{1.5}}$	$d_{\max}$
	Per	Per	Per	Per			
(1)	13.94	14.41	16.35	19.60	1	41.43	60
	13.63	13.90	14.88	17.18	3	1503.06	133
	13.34	13.59	14.60	16.12	5	8571.76	130
	13.25	13.97	14.36	15.79	10	94231.10	241
(2)	14.05	14.98	17.09	20.67	1	170.31	84
	13.73	14.93	16.13	19.15	3	5061.25	199
	13.92	14.34	15.89	18.61	5	48806.98	299
	13.50	14.26	14.75	17.40	10	347843.20	470
(3)	14.75	16.18	18.50	23.47	1	604.01	135
	14.10	15.14	16.88	20.49	3	56072.77	251
	14.54	16.14	18.37	22.13	5	303373.20	707
	13.78	14.62	15.59	18.26	10	496194.32	768

TABLE G.2
Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} Z_w(t)$ based on k -MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
Asy	2.98	3.02	3.08	3.14

	Critical Values				Graph		
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	k	$\frac{\sum_e A_e B_e }{n^{1.5}}$	$d_{\max}$
	Skew (Per)	Skew (Per)	Skew (Per)	Skew (Per)			
(1)	3.05 (3.08)	3.12 (3.16)	3.22 (3.31)	3.39 (3.59)	1	45.62	70
	3.02 (3.04)	3.08 (3.07)	3.16 (3.21)	3.29 (3.44)	3	1281.37	87
	3.02 (3.00)	3.07 (3.10)	3.15 (3.16)	3.26 (3.36)	5	8590.88	176
	3.01 (3.01)	3.07 (3.05)	3.13 (3.14)	3.23 (3.26)	10	146567.90	274
(2)	3.05 (3.08)	3.12 (3.20)	3.21 (3.33)	3.39 (3.71)	1	150.35	92
	3.02 (3.02)	3.08 (3.12)	3.16 (3.25)	3.29 (3.54)	3	6584.56	229
	3.01 (2.98)	3.07 (3.10)	3.14 (3.20)	3.26 (3.39)	5	39563.47	272
	3.01 (3.03)	3.06 (3.06)	3.13 (3.17)	3.23 (3.33)	10	325389.40	502
(3)	3.04 (3.14)	3.11 (3.29)	3.20 (3.47)	3.37 (3.89)	1	1254.18	155
	3.02 (3.06)	3.08 (3.15)	3.16 (3.32)	3.28 (3.57)	3	69460.57	247
	3.01 (3.06)	3.07 (3.15)	3.14 (3.23)	3.25 (3.43)	5	223214.80	929
	3.01 (3.00)	3.06 (3.13)	3.13 (3.23)	3.22 (3.36)	10	892840.50	936

TABLE G.3

Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} M(t)$ based on k -MST at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
Asy	3.23	3.27	3.32	3.38

	Critical Values				Graph		
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	k	$\frac{\sum_e A_e B_e }{n^{1.5}}$	$d_{\max}$
	Skew (Per)	Skew (Per)	Skew (Per)	Skew (Per)			
(1)	3.33 (3.34)	3.42 (3.45)	3.53 (3.64)	3.74 (4.05)	1	40.71	70
	3.28 (3.29)	3.35 (3.42)	3.45 (3.53)	3.60 (3.80)	3	3652.28	128
	3.26 (3.28)	3.32 (3.36)	3.40 (3.44)	3.53 (3.65)	5	10991.81	132
	3.27 (3.22)	3.33 (3.34)	3.41 (3.43)	3.53 (3.67)	10	167770.80	329
(2)	3.33 (3.34)	3.41 (3.44)	3.53 (3.68)	3.73 (4.13)	1	170.33	92
	3.31 (3.33)	3.39 (3.43)	3.49 (3.56)	3.67 (3.93)	3	6052.21	197
	3.30 (3.29)	3.37 (3.39)	3.47 (3.53)	3.63 (3.86)	5	47567.14	291
	3.27 (3.29)	3.33 (3.38)	3.42 (3.47)	3.55 (3.71)	10	204099.90	435
(3)	3.44 (3.50)	3.55 (3.71)	3.71 (4.15)	3.97 (5.05)	1	1039.34	286
	3.45 (3.39)	3.56 (3.61)	3.71 (3.98)	3.96 (4.39)	3	45800.04	203
	3.33 (3.38)	3.41 (3.49)	3.52 (3.72)	3.70 (3.98)	5	256560.60	438
	3.29 (3.30)	3.36 (3.38)	3.45 (3.55)	3.59 (3.79)	10	847006.30	720

TABLE G.4

Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} S(t)$ based on k -NNG at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
Asy	13.10	13.38	13.70	14.11

	Critical Values				Graph		
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	k	$\frac{\sum_e A_e B_e }{n^{1.5}}$	$d_{\max}$
	Per	Per	Per	Per			
(1)	14.36	15.89	17.33	22.37	1	17.42	84
	13.32	13.94	14.74	16.90	3	1136.27	99
	13.14	13.63	14.53	16.19	5	6463.44	151
	13.07	13.70	14.54	16.11	10	84495.06	292
(2)	14.20	15.01	16.84	21.17	1	43.65	88
	14.09	15.62	16.93	21.79	3	10979.30	300
	13.46	14.12	15.09	16.87	5	38128.88	187
	13.37	14.19	15.04	17.09	10	426703.30	523
(3)	14.70	16.04	18.56	24.18	1	735.27	128
	15.08	16.19	18.40	23.75	3	28824.84	417
	14.25	15.54	17.35	21.83	5	216394.94	726
	13.47	14.46	15.98	18.29	10	915297.70	546

TABLE G.5
Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} Z_w(t)$ based on k -NNG at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
Asy	2.98	3.02	3.08	3.14

	Critical Values				Graph		
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	k	$\frac{\sum_e A_e B_e }{n^{1.5}}$	$d_{\max}$
	Skew (Per)	Skew (Per)	Skew (Per)	Skew (Per)			
(1)	3.05 (3.07)	3.12 (3.21)	3.22 (3.32)	3.40 (3.61)	1	45.97	74
	3.02 (3.03)	3.08 (3.08)	3.16 (3.20)	3.30 (3.40)	3	1044.38	141
	3.02 (3.01)	3.07 (3.06)	3.15 (3.18)	3.27 (3.33)	5	32393.57	143
	3.01 (2.99)	3.06 (3.07)	3.13 (3.12)	3.23 (3.30)	10	88837.96	288
(2)	3.03 (3.13)	3.10 (3.31)	3.19 (3.49)	3.35 (3.89)	1	212.55	376
	3.02 (3.03)	3.08 (3.12)	3.16 (3.24)	3.29 (3.48)	3	14358.34	298
	3.01 (3.05)	3.07 (3.11)	3.14 (3.18)	3.25 (3.39)	5	46460.71	276
	3.01 (3.03)	3.06 (3.08)	3.13 (3.14)	3.23 (3.31)	10	288665.3	579
(3)	3.02 (3.29)	3.09 (3.41)	3.18 (3.58)	3.34 (3.96)	1	1197.74	360
	3.01 (3.10)	3.07 (3.19)	3.14 (3.40)	3.26 (3.70)	3	21535.19	622
	3.01 (3.05)	3.06 (3.13)	3.13 (3.29)	3.23 (3.56)	5	92584.55	700
	3.01 (3.01)	3.06 (3.09)	3.12 (3.19)	3.22 (3.38)	10	1039888.10	980

TABLE G.6
Critical values for the single change-point scan statistic $\max_{n_0 \leq t \leq n_1} M(t)$ based on k -NNG at 0.05 significance level. $n = 1000$.

	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$
Asy	3.23	3.27	3.32	3.38

	Critical Values				Graph		
	$n_0 = 100$	$n_0 = 75$	$n_0 = 50$	$n_0 = 25$	k	$\frac{\sum_e A_e B_e }{n^{1.5}}$	$d_{\max}$
	Skew (Per)	Skew (Per)	Skew (Per)	Skew (Per)			
(1)	3.37 (3.38)	3.46 (3.53)	3.59 (3.75)	3.82 (4.30)	1	14.02	83
	3.27 (3.27)	3.33 (3.34)	3.41 (3.46)	3.56 (3.70)	3	900.37	61
	3.27 (3.27)	3.33 (3.32)	3.41 (3.47)	3.54 (3.68)	5	7402.96	147
	3.26 (3.24)	3.31 (3.32)	3.39 (3.43)	3.50 (3.58)	10	104276.58	243
(2)	3.35 (3.35)	3.44 (3.54)	3.57 (3.77)	3.79 (4.27)	1	166.20	105
	3.32 (3.34)	3.40 (3.48)	3.50 (3.64)	3.69 (4.06)	3	11395.44	263
	3.28 (3.30)	3.35 (3.40)	3.44 (3.51)	3.59 (3.79)	5	30006.11	225
	3.28 (3.29)	3.34 (3.38)	3.42 (3.47)	3.56 (3.71)	10	261741.2	441
(3)	3.39 (3.48)	3.49 (3.66)	3.63 (3.93)	3.87 (4.45)	1	3869.47	172
	3.38 (3.43)	3.47 (3.59)	3.60 (3.72)	3.81 (4.38)	3	52365.02	475
	3.35 (3.38)	3.43 (3.52)	3.55 (3.77)	3.74 (4.10)	5	121563.09	712
	3.31 (3.30)	3.38 (3.40)	3.48 (3.60)	3.64 (3.86)	10	960134.00	843

SUPPLEMENT H: RELATIONSHIP OF THE THREE TEST
STATISTICS AND AN EXTENSION TO THE
MAX-TYPE EDGE COUNT TEST

To have a better understanding of the scan statistics $S(t)$, $Z_w(t)$, and $M(t)$, we plot their rejection regions in Figure 2. The horizontal and vertical axes in the plots are $Z_{\text{diff}}(t)$ and $Z_w(t)$, respectively. When there is only a location difference or the dimension is very low, the alternative will appear in the top of the box, where $Z_w(t)$ is quite large. In that case, $Z_w(t)$ and $M(t)$ have higher power than $S(t)$, with the power on $Z_w(t)$ slightly higher than for $M(t)$. However, when there is a scale change and the dimension is moderate to high, for a practical sample size the alternative would appear where $Z_{\text{diff}}(t)$ is quite large or quite small (the right and left parts of the box). Here both $S(t)$ and $M(t)$ still have high power while $Z_w(t)$ has very limited power. $S(t)$ and $M(t)$ have very similar power except at the boundaries. Since there is no alternative hypothesis that falls into the bottom of the box (when both $R_1(t)$ and $R_2(t)$ are smaller than their expected value), the performances of $S(t)$ and $M(t)$ are quite similar. Therefore, it works for a wide range of alternatives and $S(t)$ and $M(t)$ are preferred unless we are certain that the change is only in the mean.

FIG 2. *Rejection regions (shaded) of $S(t)$, $Z_w(t)$, and $M(t)$ for a value of t . The horizontal and vertical axes are $Z_{\text{diff}}(t)$ and $Z_w(t)$, respectively.*

When there is a change in both location and scale, depending on the the signal of interest, it is useful to consider an extended version of the max-type edge count test:

$$\max_{n_0 \leq t \leq n_1} M_\kappa(t) = \max_{n_0 \leq t \leq n_1} \max(|Z_{\text{diff}}(t)|, \kappa Z_w(t)),$$

where $\kappa \geq 0$. Different choices of κ will lead to slightly different rejection regions. For example, if we are more interested in locational changes, we

could choose a large κ . On the other hand, setting κ small would favor detecting scale change. For illustration, the rejection regions for a few κ values are shown in Figure 3.

FIG 3. *Rejection regions (shaded) of $M_\kappa(t)$ for different values of κ . The horizontal and vertical axes are $Z_{\text{diff}}(t)$ and $Z_w(t)$, respectively.*

Table H.1 presents the number of trials (out of 100) that the test reject the null hypothesis of no change under different alternatives at 0.05 significance level (with the numbers in the parentheses the number of trials where the null is rejected and the estimated change-point is within 20 from the true change-point) under different choices of κ (for $\kappa = \infty$, we set $M_\kappa(t)$ to be $Z_w(t)$). In each simulation run, we generated a sequence of $n = 200$ observations from d -dimensional Gaussian data where $d = 500$ and the change is at the center of the sequence. We consider the change both in mean and variance of the distribution. We use Δ to denote the L_2 distance of the means before and after the change. For change in variance, the first $d/5$ of the diagonal

elements of the covariance matrix change with a multiple of σ . When the mean change dominates the variance change (the first three columns of Table H.1), we see that larger values of κ have higher power. When we make σ smaller so that the variance change is dominating (the last three columns), then smaller values of κ have higher power.

Table H.2 reports the range of κ 's with estimated power at least 95% of the highest estimated power under each scenario. Based on Tables H.1 and H.2, $\kappa = 1$ seems to be a reasonable choice under all scenarios spanning from only mean change to only variance change. We thus recommend $\kappa = 1$ be a default value. If one has some prior information that the mean change is dominating, then one can set κ to be larger to achieve higher power; while if the prior information indicates that the variance change is dominating, then one can set κ smaller to achieve higher power.

TABLE H.1
Multivariate Gaussian data, mean and scale difference, τ at center.

Δ	2.2	2	1.9	1	0.5	0
σ	1	0.96	0.94	0.92	0.91	0.90
$\kappa = 0$	5 (2)	25 (8)	35 (18)	69 (45)	71 (51)	77 (52)
$\kappa = 0.05$	5 (2)	25 (8)	35 (18)	69 (45)	71 (51)	77 (52)
$\kappa = 0.35$	5 (2)	25 (8)	35 (18)	69 (45)	71 (51)	77 (52)
$\kappa = 0.5$	8 (5)	25 (8)	35 (18)	69 (45)	71 (51)	77 (52)
$\kappa = 0.65$	30 (24)	35 (17)	47 (29)	68 (44)	71 (51)	77 (52)
$\kappa = 0.75$	46 (37)	44 (24)	57 (38)	68 (44)	70 (50)	75 (51)
$\kappa = 0.85$	59 (48)	55 (36)	61 (44)	67 (44)	68 (49)	72 (48)
$\kappa = 1$	70 (55)	63 (48)	68 (47)	60 (39)	67 (48)	65 (43)
$\kappa = 1.14$	71 (57)	65 (50)	68 (49)	58 (40)	59 (41)	60 (39)
$\kappa = 1.5$	74 (59)	71 (56)	68 (51)	25 (11)	25 (13)	18 (9)
$\kappa = 2$	74 (59)	71 (56)	69 (52)	21 (9)	13 (3)	3 (1)
$\kappa = 2.5$	74 (59)	71 (56)	69 (52)	21 (9)	13 (3)	3 (1)
$\kappa = 3$	74 (59)	71 (56)	69 (52)	21 (9)	13 (3)	3 (1)
$\kappa = 5$	74 (59)	71 (56)	69 (52)	21 (9)	13 (3)	3 (1)
$\kappa = 10$	74 (59)	71 (56)	69 (52)	21 (9)	13 (3)	3 (1)
$\kappa = 20$	74 (59)	71 (56)	69 (52)	21 (9)	13 (3)	3 (1)
$\kappa = \infty$	74 (59)	71 (59)	69 (52)	21 (9)	13 (3)	3 (1)

TABLE H.2
Range of κ with the estimated power at least 95% of the highest estimated power under each scenario.

Δ	2.2	2	1.9	1	0.5	0
σ	1	0.96	0.94	0.92	0.91	0.90
κ	[1.01, $+\infty$)	[1.17, $+\infty$)	[0.95, $+\infty$)	[0, 0.86]	[0, 0.95]	[0, 0.84]

SUPPLEMENT I: BLOCK PERMUTATION FOR LOCAL DEPENDENCY

Independence is an ideal assumption that may be violated in practice. This section discusses one way to deal with local dependency in the sequence. Instead of using the permutation null, we could do block permutation, i.e., the sequence is divided into blocks of size b and the blocks are then permuted. In this way, the local structure in the sequence is retained. In order to retain the local structure within any block of size b in some block permuted sequences, we alter the number of observations in the first block, i.e., the size of the first block ranges from 1 to b . All following blocks are of size b except for the last block that might be of size less than b . Thus, there are b ways to partition the sequence. To get one block permuted sequence, we first randomly choose one out of the b ways in partitioning the sequence and then randomly permute the blocks.

Then, the standardized two-sample test statistics at t for testing H_0 (1.1) against H_1 (1.2) are defined as:

$$Z_{w,bp}(t) = \frac{R_w(t) - \mathbf{E}_{bp}(R_w(t))}{\sqrt{\mathbf{Var}_{bp}R_w(t)}}; \quad Z_{\text{diff},bp}(t) = \frac{R_{\text{diff}}(t) - \mathbf{E}_{bp}(R_{\text{diff}}(t))}{\sqrt{\mathbf{Var}_{bp}R_{\text{diff}}(t)}},$$

where $\mathbf{E}_{bp}(R_w(t))$, $\mathbf{Var}_{bp}R_w(t)$, $\mathbf{E}_{bp}(R_{\text{diff}}(t))$, $\mathbf{Var}_{bp}R_{\text{diff}}(t)$ are the expectation and variance for $R_w(t)$ and $R_{\text{diff}}(t)$, respectively, under the block permutation. Based on these, we can further define:

$$S_{bp}(t) = Z_{w,bp}^2(t) + Z_{\text{diff},bp}^2(t); \quad M_{bp}(t) = \max(|Z_{\text{diff},bp}(t)|, Z_{w,bp}(t)).$$

The test statistics are now defined as:

$$\max_{n_0 \leq t \leq n_1} Z_{w,bp}(t), \quad \max_{n_0 \leq t \leq n_1} S_{bp}(t), \quad \text{and} \quad \max_{n_0 \leq t \leq n_1} M_{bp}(t).$$

The test statistics for testing H_0 (1.1) against H_2 (1.3) under block permutation are defined in a similar manner. The p -values for the above statistics can be obtained by 10,000 block permutation runs. We illustrate the tests under block permutation on the same dataset as in Section 6. The results under different block sizes (1,2,3) of the new test statistics together with the method in [Chen and Zhang \(2015\)](#) are summarized in Table I.1. Observe that a block size of 1 is equivalent to permutation under independence assumption.

From Table I.1, we see that, as b increases, the p -values becomes larger for all tests, while they are all below 0.05, indicating the sequence is not homogeneous. Also, for each test, the estimations to the location of the

TABLE I.1

Changed interval results and corresponding p -values from 10,000 block permutations (reported in parentheses) for NYC taxi pickups from JFK during the period of Oct. 1 - Dec. 31, 2015.

block size	Z_{bp}	$Z_{w,bp}$	S_{bp}	M_{bp}
1	11/21 - 12/31 (< 0.0001)	12/18 - 12/25 (0.0001)	12/18 - 12/25 (0.0001)	12/18 - 12/25 (0.0001)
2	11/21 - 12/31 (0.0011)	12/18 - 12/25 (0.007)	12/18 - 12/25 (0.007)	12/18 - 12/25 (0.007)
3	11/21 - 12/31 (0.0177)	12/18 - 12/25 (0.0256)	12/18 - 12/25 (0.0269)	12/18 - 12/25 (0.0256)

change-points are all the same for different b 's. The increase in p -values as b increases indicates that the sequence might have dependency structure over time, while the results under independence assumption still give us something reasonable.

REFERENCES

- CHEN, L. H. and SHAO, Q.-M. (2005). Steins method for normal approximation. *An introduction to Steins method* **4** 1–59.
- CHEN, H. and ZHANG, N. (2015). Graph-based change-point detection. *The Annals of Statistics* **43** 139–176.
- SIEGMUND, D. (1988). Approximate tail probabilities for the maxima of some random fields. *The Annals of Probability* 487–501.
- SIEGMUND, D. (1992). Tail approximations for maxima of random elds. *Probability theory* 147158.
- WOODROOFE, M. (1976). Frequentist properties of Bayesian sequential tests. *Biometrika* **63** 101–110.
- WOODROOFE, M. (1978). Large deviations of likelihood ratio statistics with applications to sequential testing. *The Annals of Statistics* 72–84.