Special issue of the 3rd International Conference on Computational and Experimental Science and Engineering (ICCESEN 2016)

Determination of ^{40}K Concentration in Gravel Samples from Konyaaltı Beach, Antalya

S. SEÇKİNER^a, İ. AKKURT^a AND K. GÜNOĞLU^{b,*}^aSüleyman Demirel University, Science and Art Faculty, Department of Physics, Isparta, Turkey^bSüleyman Demirel University, Technical Vocational School, Isparta, Turkey

Terrestrial radioactivity is caused by the radioactive elements, located in various amounts in soil and rocks. The concentration of radioactive elements varies with the geographical and geological structure of the region and with the mineralogical composition of soil and rock. In this study, ^{40}K activity concentrations in gravel samples, collected from Konyaaltı Beach, were measured. The measurement was performed using gamma ray spectrometry at gamma spectrometry laboratory of Süleyman Demirel University.

DOI: [10.12693/APhysPolA.132.1095](https://doi.org/10.12693/APhysPolA.132.1095)

PACS/topics: 23.90.+W, 78.66.SQ

1. Introduction

Natural radioactivity arising from natural sources is the main contribution to the annual dose received by the world's population. This exposure is resulting from radionuclides inherent in the Earth's crust and from cosmic rays. The terrestrial ubiquitous radionuclides belong to the ^{238}U and ^{232}Th series and their decay products, as well as to single decay radionuclides, particularly ^{40}K . Gamma radiation emitted from such naturally occurring radionuclides in all ground formations represents the main part of external exposure of human body [1]. The knowledge of concentrations and distributions of the radionuclides in the materials is of interest since it provides useful information for the monitoring of environmental radioactivity. Natural environmental radioactivity and the associated external exposure due to gamma radiation depend primarily on the geological and geographical conditions, and appear at different levels in the soils of each region in the world [2–4].

Of the three isotopes of naturally occurring potassium, only ^{40}K is unstable and has a half-life of 1.277×10^9 years. 0.0118% of the natural potassium forms the ^{40}K isotope. Because potassium is so abundant in the soil, it causes a significant fraction of the natural gamma radiation dose.

Konyaaltı beach in Antalya is the most preferred beach by native and foreign tourists. People spend most of their time in summer on this beach. Therefore, determination of the level of natural radioactivity of the beach is important for human health. In this study, ^{40}K activity concentrations of some gravel samples collected from Konyaaltı Beach were measured. Konyaaltı Beach in Antalya province, which attracts both the local people and the tourists, is about 3 km west of Antalya and is

Fig. 1. Location of Konyaaltı Beach in Antalya.

about 8 km long. This beach contains a mixture of sand and gravel. Figure 1 shows Konyaaltı Beach.

2. Materials and methods

In this study, seven gravel samples have been collected from Konyaaltı Beach. After collection of samples, they were crushed and dried at 100°C in an oven for about 24 h. The dried samples have been poured in cups, which were then tightly sealed with a thick tape around their necks, to limit any gas escape, and stored for four weeks to achieve secular equilibrium between ^{238}U and its progeny [5].

The radioactivity concentration of ^{40}K in gravel samples were determined using gamma ray spectrometer consisting of a $3'' \times 3''$ NaI(Tl) detector, connected to a multi-channel-analyser. The spectra were analyzed using the MAESTRO32, obtained from ORTEC. The schematic view of the experimental system is shown in Fig. 2.

Before measurements the system had to be calibrated. This was carried out using ^{137}Cs and ^{60}Co radioactive sources, which produce γ -rays with energy of 662, 1173 and 1332 keV, respectively. The γ -ray spectrum obtained from the mentioned source and the related fit are displayed in Fig. 3.

*corresponding author; e-mail: kadirgnoglu@gmail.com

Fig. 2. Schematic view of gamma spectrometer and electronic units.

Fig. 3. The γ -ray energy spectrum obtained from ^{137}Cs and ^{60}Co sources (upper) and related fit channel versus energy (keV).

The measurement was based on recording natural radioactivity quantities of three natural long-lived elements: ^{226}Ra , ^{232}Th and ^{40}K , which produce the photo-peaks at 1760, 2610 and 1461 keV respectively, in the natural γ -ray spectrum [6].

The activities for the natural radionuclides were calculated using the following relation [7]

$$A(\text{Bq/kg}) = \frac{N}{\varepsilon \gamma t m}, \quad (1)$$

where A is the activity of the radionuclide in Bq/kg, N is the net peak area under the most prominent photo peaks, calculated by subtracting the background spectrum from the respective count rate, obtained for the same counting time, ε is the detector efficiency for the specific gamma ray, γ is the absolute transition probability of gamma decay, t is the counting time (s) and m the mass of the sample (kg).

3. Results and discussion

The ^{40}K concentration of some gravel samples, collected from seven different points of Konyaaltı Beach, have been measured and the results are shown in Fig. 4. It can be seen from this figure that the measured ^{40}K concentration has ranged from 421.12 to 3363 Bq/kg.

Fig. 4. The activity concentrations of ^{40}K .

The results were compared with the worldwide average concentration, which is 500 Bq/kg for ^{40}K [8]. Measured concentration of ^{40}K for all samples is lower than this average value. Table I compares the activity concentration of ^{40}K , obtained in this study, with those reported in literature and with the world average value. From the table, the values obtained in this study compare favorably with those from other countries.

TABLE I

Comparison of activity concentrations of ^{40}K from different countries.

Location	Activity concentr. [Bq/kg]	References
	^{40}K	
India	349.6	[9]
China	396.4 ± 75.1	[10]
Iran	$179.5 - 464.5$	[11]
Thailand	523	[12]
Penang, Malaysia	369 ± 17	[13]
Saudi Arabia Gulf	153.8	[14]
Oman	$222.89 - 535.07$	[15]
Saudi Arabia	641.08	[16]
Turkey	350.42 ± 17.52	this work

Acknowledgments

This work has been supported by Suleyman Demirel University Foundation Unit (BAP) with the project number of 4668_YL2_16.

References

- [1] T. Santawamaitre, D. Malain, H.A. Al-Sulaiti, M. Matthews, D.A. Bradley, P.H. Regan, *Nucl. Instr. Meth. Phys. Res. A* **652**, 920 (2010).

- [2] A. Abbady, N.K. Ahmed, A.M. El-Arabi, R. Miche, A.H. El-Kamel, A.G.E. Abbady, *Nucl. Sci. Techniq.* **17**, 118 (2006).
- [3] A.M. El-Arabi, N.K. Ahmed, A.H. El-Kamel, *Gamma spectroscopic analysis of powdered granite samples in some Eastern desert's areas*, in: *Proc. 5th Radiation Conference*, Cairo, Egypt 2000.
- [4] A.I. Abd El-mageed, A.H. El-Kamel, A. Abbady, S. Harb, A.M.M. Youssef, I.I. Saleh, *Radiat. Phys. Chem.* **80**, 710 (2011).
- [5] I. Akkurt, B. Oruncak, K. Gunoglu, *Int. J. Phys. Sci.* **5**, 170 (2010).
- [6] İ. Akkurt, K. Günoğlu, *Sci. Technol. Nucl. Install.* **2014**, 950978 (2014).
- [7] I. Akkurt, B. Mavi, H. Akyıldırım, K. Günoglu, *Int. J. Phys. Sci.* **4**, 403 (2009).
- [8] UNSCEAR, United Nations Scientific Committee on the Effects of Atomic Radiation, *Sources, effects and risks of ionizing radiation. Report to the General Assembly with Annex B: Exposures from Natural Sources of Radiation*, United Nations, New York 2000.
- [9] S.M. Gandhi, R. Ravisankar, A. Rajalakshmi, S. Sivakumar, A. Chandrasekaran, P.D. Anand, *J. Radiat. Res. Appl. Sci.* **7**, 7 (2014).
- [10] Y. Huang, X. Lu, X.W. Ding, T. Feng, *Mar. Pollut. Bull.* **91**, 357 (2015).
- [11] M. Tari, S.A.M. Zarandi, K. Mohammadi, M.R. Zare, *Mar. Pollut. Bull.* **74**, 425 (2013).
- [12] D. Malain, P.H. Regan, D.A. Bradley, M. Matthews, H.A. Al-Sulaiti, T. Santawamaitre, *Appl. Radiat. Isot.* **70**, 1467 (2012).
- [13] B.A. Almayahi, A.A. Tajuddin, M.S. Jaafar, *Radiat. Phys. Chem.* **81**, 1547 (2012).
- [14] H. Al-Ghamdi, A. Al-Muqrin, A. El-Sharkawy, *Mar. Pollut. Bull.* **104**, 29 (2016).
- [15] M.R. Zare, M. Mostajaboddavati, M. Kamali, M.R. Abdi, M.S. Mortazavi, *Mar. Pollut. Bull.* **64**, 1956 (2012).
- [16] H. Al-Trabulsy, A. Khater, F. Habbani, *Radiat. Phys. Chem.* **80**, 343 (2011).