Radioactivity Investigation of Sediment Samples from Beni Haroun Dam Using High-Resolution Gamma-Ray Spectroscopy

G. BOUHILA* AND F. BENRACHI

Mathematical and Subatomic Physics Laboratory (LPMS), Frères Mentouri University, Constantine, Algeria

In this study, the levels of natural and anthropogenic radioactivity in the sediment samples, collected from Beni Haroun Dam, were investigated using gamma-ray spectrometry. The results obtained for the ^{238}U , ^{232}Th series, ^{40}K natural element and ^{137}Cs anthropogenic radionuclide are discussed. To evaluate the radiological hazard of radioactivity in samples, the absorbed dose rate, the annual effective dose, the radium equivalent activity R_{eq} , the external and internal hazard indices H_{ex} and H_{in} were calculated and presented in comparison with the data collected from different areas in the world.

DOI: [10.12693/APhysPolA.132.889](https://doi.org/10.12693/APhysPolA.132.889)

PACS/topics: 29.90.+r, 29.30.-h

1. Introduction

Radiation is present everywhere, even without any human contribution. According to the UNSCEAR report, approximately 86% of the radiation to which humans are exposed, arises from natural radioactivity (^{238}U , ^{232}Th series and ^{40}K) and the remaining 14% are from anthropogenic radioactivity (^{137}Cs) [1, 2].

The activity of natural radionuclides in soil and sediment depends mainly on the type of the rocks from which they originate [3–5]. The anthropogenic radionuclide ^{137}Cs has been produced and released into the environment by human nuclear activity, including nuclear weapons testing, the operation of nuclear power plants, research reactors, and nuclear fuel reprocessing [1, 2].

In order to obtain the information about the present levels of environmental radioactivity, various studies were carried out in many countries [2–16]. In the current work, sediment samples were collected near Beni Haroun Dam. This artificial dam is located at $36^{\circ} 33' 19'' \text{ N}$, $6^{\circ} 16' 11'' \text{ E}$, in North-Eastern Algeria, at about 40 km north of Constantine Province.

The aim of this study is the measurement of the ^{238}U , ^{235}U , ^{232}Th , ^{40}K and ^{137}Cs radionuclide concentrations, to provide background data on natural and anthropogenic radioactive isotopes for the studied region. Additionally, the total absorbed dose rate (ADR), the annual effective dose equivalent (AED), the average radium equivalent activity R_{eq} , the external and internal hazard indices H_{ex} , H_{in} were calculated and compared with the results from the literature.

2. Sampling and measurements

In order to evaluate, the radioactivity levels of natural and anthropogenic radionuclides, surface sediment

samples (0–10 cm) were taken from five locations near Beni Haroun Dam in spring 2016 (Fig. 1). The sediments were dried, pulverized, homogenized, weighed and packed into small cylindrical plastic containers for approximately 30 days (~ 7 half-lives) to reach secular equilibrium between the ^{238}U and ^{232}Th series and their respective progeny, before the measurements.

Fig. 1. Map of location of the sample sites.

The activity concentrations of these samples were measured using a planar BEGe detector (Broad Energy Germanium, model BE3825) which has a relative efficiency of 28% and an energy resolution of 1.72 keV at 1332 keV. All spectra were analysed using the Genie 2000 software (Canberra). Spectrum of each sample was collected for 86400 seconds.

The absolute efficiency calibration of the gamma spectrometer has been carried out using standard source IAEA-384. The counting geometry for the sources and sediment samples were identical. The activity concentrations were averaged from gamma-ray photo peaks at several energies, assuming secular equilibrium in the ^{238}U and ^{232}Th series.

The ^{226}Ra activity was calculated from the gamma lines associated with the short-lived daughters of ^{214}Pb (351.93 keV) and ^{214}Bi (609.31 keV). ^{238}U activity was determined from 63.3 keV line of ^{234}Th . While, ^{232}Th activity was estimated by the peaks of ^{228}Ac (911.20 keV)

*corresponding author; e-mail: bouhilaghania@yahoo.fr

and ^{208}Tl (583.19 keV). The activity concentrations of ^{40}K and ^{137}Cs were measured directly from the peak areas at 1460.80 and 661.62 keV, respectively.

3. Results and discussion

The activity concentrations of radionuclides ^{226}Ra , ^{232}Th , ^{40}K and ^{137}Cs have been measured in sediments from five sampling sites near Beni Haroun Dam. The results, presented in Table I, show that the natural

radionuclide concentrations vary from site to site, which is explained by the difference in geological composition of sediment samples. In all sampling sites, the activity of natural radionuclides ^{226}Ra , ^{232}Th , ^{40}K are lower than the world averages which are 35, 33 and 400 respectively [1]. However, the anthropogenic radionuclide ^{137}Cs was observed in three locations (Fig. 2: SBH-1, SBH-4 and SBH-5), where SBH-5 sample has the greatest value. This region is the estuary of Rhumel River in Beni Haroun Dam.

TABLE I

Activity concentrations of ^{226}Ra , ^{232}Th and ^{40}K and dose assessment of investigated samples.

Location	Concentration			Dose estimation	
	^{226}Ra [Bq kg $^{-1}$]	^{232}Th [Bq kg $^{-1}$]	^{40}K [Bq kg $^{-1}$]	ADR [$\eta\text{Gy h}^{-1}$]	AED [$\mu\text{Sv yr}^{-1}$]
SBH-1	23.75 $\pm$ 1.93	26.49 $\pm$ 4.45	180.52 $\pm$ 21.49	34.83 $\pm$ 3.03	42.71 $\pm$ 3.72
SBH-2	15.97 $\pm$ 1.39	20.14 $\pm$ 3.36	139.05 $\pm$ 19.74	25.59 $\pm$ 2.33	31.38 $\pm$ 2.85
SBH-3	16.34 $\pm$ 1.72	18.25 $\pm$ 1.55	201.74 $\pm$ 15.96	27.15 $\pm$ 1.41	33.30 $\pm$ 1.73
SBH-4	14.90 $\pm$ 0.32	21.47 $\pm$ 0.89	261.00 $\pm$ 2.63	26.04 $\pm$ 0.58	33.93 $\pm$ 0.71
SBH-5	17.3 $\pm$ 0.41	24.80 $\pm$ 0.74	203.00 $\pm$ 3.06	68.40 $\pm$ 1.16	38.90 $\pm$ 0.63
Mean	17.65 $\pm$ 1.93	22.23 $\pm$ 4.45	197.06 $\pm$ 21.49	29.07 $\pm$ 3.03	36.04 $\pm$ 3.72

Fig. 2. ^{137}Cs activities in sediment samples.

The absorbed dose rate in air at 1 m above the ground surface and annual effective dose are calculated using Eqs. (1) and (2), respectively [1]:

$$\text{ADR}(\text{nGy/h}) = 0.462A_{\text{Ra}} + 0.621A_{\text{Th}} + 0.0417A_{\text{K}}, \quad (1)$$

$$\text{AED}(\mu\text{Sv/y}) = \text{ADR}(\text{nGy/h}) \times 365.25 (\text{d y}^{-1})$$

$$\times 24 (\text{h d}^{-1}) \times 0.2 \times 0.7 (\text{Sv/Gy}) \times 10^{-3}, \quad (2)$$

where A_{K} , A_{Th} and A_{Ra} are the specific activities of ^{40}K , ^{238}Th and ^{226}Ra (Bq kg $^{-1}$), respectively.

The obtained ADR and AED values for all samples, listed in Table I, are lower than the world averages of 59 nGy/h and 70 $\mu\text{Sv/y}$ [1], respectively. Besides, the results were evaluated in terms of the radiation hazard by means of the radium equivalent activity (Ra_{eq}) which is calculated through the relation given by [17, 18].

$$\text{Ra}_{\text{eq}} = 0.077A_{\text{K}} + 1.43A_{\text{Th}} + A_{\text{Ra}}. \quad (3)$$

The Ra_{eq} values for investigated locations are shown in Table II. In addition, H_{ex} and H_{in} indices were calculated by means of [17, 19]:

$$H_{\text{ex}} = \frac{A_{\text{Ra}}}{370} + \frac{A_{\text{Th}}}{259} + \frac{A_{\text{K}}}{4810}, \quad (4)$$

$$H_{\text{in}} = \frac{A_{\text{Ra}}}{185} + \frac{A_{\text{Th}}}{259} + \frac{A_{\text{K}}}{4810}. \quad (5)$$

In Table II, the values of H_{ex} and H_{in} indices range from 0.15 to 0.20 and 0.19 to 0.27, respectively. The values of Ra_{eq} , H_{ex} and H_{in} in the study area are below their permissible limits, which indicates that the sediments do not pose any significant radiation hazard.

TABLE II

Radiation hazard indices for investigated samples.

Location	Radiation hazard indices		
	Ra_{eq} [Bq kg $^{-1}$]	H_{ex}	H_{in}
SBH-1	75.54 $\pm$ 6.85	0.20 $\pm$ 0.02	0.27 $\pm$ 0.02
SBH-2	55.48 $\pm$ 5.22	0.15 $\pm$ 0.01	0.19 $\pm$ 0.02
SBH-3	57.97 $\pm$ 3.06	0.16 $\pm$ 0.01	0.20 $\pm$ 0.01
SBH-4	56.54 $\pm$ 1.33	0.15 $\pm$ 0.01	0.19 $\pm$ 0.00
SBH-5	68.40 $\pm$ 1.16	0.18 $\pm$ 0.01	0.23 $\pm$ 0.00
Mean	62.78 $\pm$ 6.85	0.17 $\pm$ 0.02	0.21 $\pm$ 0.02

Table III illustrates the comparison of the results obtained in present study with the activity concentrations of natural radionuclides reported in [12–16] for some African countries. The mean values of ^{226}Ra , ^{232}Th in sediment samples of the present work are higher than those

of the other countries, with the exception of Nigeria and Sudan. The activity of ^{40}K is close to those of Ghana and Egypt, but it is lower than the ones of Nigeria and Sudan.

TABLE III

^{226}Ra , ^{232}Th and ^{40}K concentrations in sediment of study area, in comparison with African countries.

Location	^{226}Ra [Bq kg $^{-1}$]	^{232}Th [Bq kg $^{-1}$]	^{40}K [Bq kg $^{-1}$]	Refs.
Algeria	17.65 $\pm$ 1.93	22.23 $\pm$ 4.45	197.06 $\pm$ 21.49	Pres. study
Ghana	16.02 $\pm$ 9.99	20.31 $\pm$ 16.39	195.01 $\pm$ 116.49	[12]
Nigeria	47.89 $\pm$ 18.67	55.37 $\pm$ 32.74	1023.00 $\pm$ 474.00	[13]
Egypt	16.30 $\pm$ 9.22	12.94 $\pm$ 6.23	200.21 $\pm$ 52.94	[14]
Libya	7.50 $\pm$ 2.50	4.50 $\pm$ 1.30	28.50 $\pm$ 6.70	[15]
Sudan	22.83 $\pm$ 4.03	25.11 $\pm$ 4.96	284.31 $\pm$ 80.45	[16]

4. Conclusions

The present study was carried out to give a baseline reference data about the radioactivity levels from natural radionuclides in sediment samples, collected from some locations around Beni Haroun Dam, evaluated using high resolution gamma-ray spectrometry (HPGe detector). The results indicate that the activity levels of ^{226}Ra , ^{232}Th and ^{40}K natural radionuclides in all samples are lower than the world values reported by UNSCEAR [1]. Presence of anthropogenic radionuclide ^{137}Cs was shown in three locations, with different values. Moreover, the obtained values of absorbed dose rate, annual effective dose, radium equivalent activity, external and internal hazard indices were found to be below the permissible limits.

Acknowledgments

Authors of this article thank the organizers of the 3rd International Conference on Computational and Experimental Science and Engineering 19–24 October 2016, Kemer-Antalya-Turkey, for the organization and the support provided during the conference.

Special thanks owed to Dr J.E. Groetz and C. Ma von from Chrono-Environment Laboratory, Bourgogne Franche-Comté University, Besançon, for their help.

References

- [1] United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR), *Sources and Effects of Ionizing Radiation*, Report to the General Assembly, New York 2000.
- [2] V. Kannan, M.P. Rajan, M.A.R. Iyengar, R. Ramesh, *Appl. Radiat. Isotopes* **57**, 109 (2002).
- [3] N. Zaim, A.B. Tugrul, H. Atlas, B. Buyuk, E. Demir, N. Baydogan, N. Altinsoy, *Acta Phys. Pol. A* **130**, 64 (2016).
- [4] B. Çetin, F. Öner, I. Akkurt, *Acta Phys. Pol. A* **130**, 475 (2016).
- [5] G. Bouhila, F. Benrachi, M. Ramdhane, *IJNuRaSaT* **1**, 7 (2016).
- [6] B. Baggoura, A. Nourredine, M. Benkrid, *Appl. Radiat. Isotopes* **49**, 867 (1998).
- [7] S.A. Khatir, M.M.O. Ahmed, F.A. El-Khang, Y.O. Nigumi, E. Holm, *Radiat. Prot. Dosim.* **71**, 141 (1997).
- [8] B. Mitrović, G. Ajitić, M. Lazić, V. Andrić, N. Krstić, B. Vranješ, M. Vićentijević, *Environ. Pollut.* **215**, 273 (2016).
- [9] T. Rajeshwari, S. Rajesh, B.R. Kerur, S. Anilkumer, N. Krishnan, A.D. Pant, *J. Radioanal. Nucl. Chem.* **300**, 61 (2014).
- [10] A. Coşkun, B. Çetin, *Acta Phys. Pol. A* **130**, 309 (2016).
- [11] M.C. Boz, F. Öner, B. Mavi, *Acta Phys. Pol. A* **128**, B-360 (2015).
- [12] O.K. Adukp, A. Faanu, H. Lawluvi, L. Tettey-Larbi, G. Emi-Reynolds, E.O. Darko, C. Kansaana, D.O. Kpeglo, A.R. Awudu, E.T. Glover, P.A. Amoah, A.O. Efa, *J. Radioanal. Nucl. Chem.* **303**, 1679 (2015).
- [13] M.O. Isinkaye, H.U. Emelue, *J. Radiat. Res. Appl. Sci.* **8**, 459 (2015).
- [14] A. El-Gamal, S. Nasr, A. El-Taher, *Radiat. Meas.* **42**, 457 (2007).
- [15] S.U. El-Kameesy, S. Abd El-Ghany, S.M. El-Minyawi, Z. Miligy, E.M. El-Mabrouk, *Turkish J. Eng. Env. Sci.* **32**, 245 (2008).
- [16] N. Fadöl, I. Salih, H. Idriss, A. Elfaki, A. Sam, *Res. J. Physical Sci.* **3**, 1 (2015).
- [17] J. Beretka, P.J. Mathew, *Health Phys.* **48**, 111 (1985).
- [18] B. Mavi, I. Akkurt, *Rad. Phys. Chem.* **79**, 933 (2010).
- [19] I. Akkurt, *Ann. Nucl. En.* **36**, 1702 (2009).