Proc. XIX International School of Semiconducting Compounds, Jaszowiec 1990

SUBMILIMETER MAGNETOSPECTROSCOPY ON ZnFeSe

R. KREVEK, P. KEMPF, M. VON ORTENBERG

Institut für Halbleiterphysik und Optik, Technische Universität Braunschweig,
Mendelssohnstr. 3, D-3300 Braunschweig, Germany

AND A. TWARDOWSKI

Institute of Experimental Physics, Warsaw University, Hoża 69, 00-681 Warszawa,
Poland

(Received August 8, 1990)

We report FIR laser spectroscopy study of $\text{Zn}_{1-x}\text{Fe}_x\text{Se}$ ($x < 0.06$) Semimagnetic Semiconductor at the temperature range of 2–26 K and magnetic fields up to 18 T.

PACS numbers: 78.30.Fs, 71.70.Ej

One of the central problems of Fe-based Semimagnetic Semiconductor (SMSC) is the information about the energy structure of the Fe^{++} ion ground term (5E , split into a singlet A_1 , a triplet T_1 , a doublet E , a triplet T_2 and a singlet A_2 [1]). A useful tool to study this structure is far infrared (FIR) spectroscopy [1, 2]. Recently, an investigation of Fe-type SMSC has been reported [3, 4], however it dealt with rather low Fe concentrations. In this paper we present complementary results of FIR laser spectroscopy for $\text{Zn}_{1-x}\text{Fe}_x\text{Se}$ ($x < 0.06$) in the temperature range $2 < T < 26$ K and magnetic fields up to 18 T.

In Fig. 1 we display resonant energies observed experimentally as a function of magnetic field. The previously observed [3, 4] transitions $A_1 - T_1$ and $A_1 - E$ are clearly visible also for higher x , although the spectra are broader than those obtained for low x . The experimental energies match well the calculated energy level differences for isolated, i.e. not interacting Fe^{++} ion (Fig. 1). We notice that the line observed at 14.3 cm^{-1} coincides with the lowest transition $A_1 - T_1$. However, this transition is only weakly dependent on the field and therefore should not be observed in laser spectroscopy as a resonance (fixed energy of laser line). In that respect the origin of 14.3 cm^{-1} line is still not clear.

Fig. 1. The experimentally observed resonance energies as a function of magnetic field; the lines present calculated energies of transitions: solid lines – model from [1] with $Dq = 293 \text{ cm}^{-1}$ and $\lambda = -95 \text{ cm}^{-1}$, dotted lines – model from [2] with $Dq = 294 \text{ cm}^{-1}$ and $\lambda = -88 \text{ cm}^{-1}$; parameters were obtained by fitting to the zero-field energies of $A_1 - T_1$, $A_1 - T_2$ transitions as well as zero phonon line of ${}^5E - {}^5T$ transition.

For $\text{Zn}_{0.94}\text{Fe}_{0.06}\text{Se}$ we observed a weak and broad line at 8.2 cm^{-1} (Fig. 1). This line disappears at temperatures $T > 5 \text{ K}$ and is not visible for lower Fe concentrations. We stress that the transition energy is roughly two times smaller than the lowest transition energy expected for isolated Fe ion. A possible candidate for the 8.2 cm^{-1} lines is the transition between the ground and the first excited states of a Fe-Fe pair coupled by exchange interaction. In ZnFeSe , calculated energy of this transition is about $8\text{--}9 \text{ cm}^{-1}$ [5]. An increasing contribution of pair and larger clusters to FIR absorption should be expected for $x > 0.03$ and is indeed exhibited by the changes of FIR absorption spectra shape [6, 3] as well as by the temperature variation of the spectra (Fig. 2). We believe, however, that the available data are too preliminary to establish the nature of 8.2 cm^{-1} line.

Fig. 2. The experimental FIR transmission spectra of $\text{Zn}_{0.94}\text{Fe}_{0.06}\text{Se}$ as a function of magnetic field for various temperatures. The laser energy was 21.2 cm^{-1} . Curves at $9 < T < 26 \text{ K}$ show results for upward and downward sweeps of magnetic field.

References

- [1] G.A. Slack, S. Roberts, J.T. Wallin, *Phys. Rev.* **187**, 511 (1969).
- [2] J.T. Vallin, G.A. Slack, C.C. Bradley, *Phys. Rev. B* **2**, 4406 (1970).
- [3] A.M. Witowski, A. Twardowski, M. Pohlmann, W.J.M. de Jonge, A. Wieck, A. Mycielski, M. Demianiuk, *Solid State Commun.* **70**, 27 (1989).
- [4] M. Hausenblas, L.M. Claessen, A. Wittlin, A. Twardowski, M. von Ortenberg, W.J.M. de Jonge, P. Wyder, *Solid State Commun.* **72**, 253 (1989).
- [5] A. Twardowski, H.J.M. Swagten, T.F.H. v.d. Wetering, W.J.M. de Jonge, *Solid State Commun.* **65**, 235 (1988).
- [6] A. Twardowski, H.J.M. Swagten, W.J.M. de Jonge, in *Proc. 19th Internat. Conf. on the Physics of Semiconductors, Warszawa 1988*, ed. W. Zawadzki, Institute of Physics, Polish Academy of Sciences, Warsaw 1989, p. 1543.