Received on 20 April, 2014; received in revised form, 14 June, 2014; accepted, 19 July, 2014; published 01 November, 2014

ISOLATION OF ACTIVE COMPOUNDS FROM HYDRO ALCOHOLIC EXTRACT OF *CARDIOSPERMUM HALICACABUM* LINN

R. Rajeswari¹ and M. Sriidevi^{*2}

Department of Biochemistry, Vinayaka Missions Medical College¹, Vinayaka Missions University, Ariyanoor - 636301, Tamilnadu, India.

Department of Biotechnology, Faculty of Engineering, Vinayaka Missions Engineering college², Vinayaka Missions University, Ariyanoor - 636301, Tamilnadu, India.

Keywords:

Cardiospermum halicacabum Linn., Isolation, Kaempferol- 3- o- α - l- rhamnoside, Apigenin – 7- o- β - d- glucuronide, Antioxidant.

Correspondence to Author:

M. Sriidevi

Assistant Professor,
Department of Biotechnology,
Vinayaka Missions Engineering
college, Vinayaka Missions University,
Periyaseeragapadi – 636301,
Salem, Tamilnadu, India.

E-mail: drsridevimuruhan@gmail.com

ABSTRACT: The uses of plant medicines are very common in Asia and developing countries. Phytochemicals has been used to treat chronic as well as infectious diseases. *Cardiospermum halicacabum* Linn. (Sapindaceae) is an herbaceous climber, commonly found throughout India, used as vegetable as well as traditional medicine for rheumatism, lumbago and nervous diseases. Many pharmacological studies have been done with various parts of *C. halicacabum* plant to demonstrate its medicinal effect. The present work was aimed to isolate the active compounds from the leaf extract of *C. halicacabum*. The hydro alcoholic leaf extract (80% methanol + 20% distilled water) was subjected to isolation procedures for identification of active compounds. Eleven compounds were got and *in vitro* anti oxidant analysis were done with the isolated compounds. Compound 1 and 8 selected for structural elucidation. The NMR spectrum showed characteristic peaks of compound 1 and 8 and the active phytoconstituents were identified as Kaempferol- 3- o- α - l- rhamnoside and Apigenin – 7- o- β - d- glucuronide by structural elucidation. The isolated compounds showed strong antioxidant potential.

INTRODUCTION: The uses of natural drugs are very common in Asia and developing countries. According to World Health Organization (WHO) more than 80% of the world's population relies on traditional medicine due to their efficiency or due to the costs or side effects of synthetic drugs. Plant medicines have been used to treat chronic as well as infectious diseases¹. Natural products, such as plant extracts either as pure compounds or as standardized extracts, provide unlimited opportunities for new drug discoveries.

Many beneficial biological activity such as anticancer, antimicrobial, antioxidant, antidiarrheal, analgesic²⁻⁶ and wound healing activity were reported. So efficient detection and rapid characterization of natural products play an important role in the phytochemical investigation of crude plant extracts and its therapeutic uses.

Cardiospermum halicacabum (*C. halicacabum*) is an herbaceous climber, widely distributed in tropical and subtropical Asia and Africa, and often found throughout India. Different parts of the plant are used in traditional medicine for the treatment of rheumatism, lumbago, cough, hyperthermia, nervous diseases, stiffness of limbs etc. Many pharmacological studies have been done with various parts of *C. halicacabum* plant to demonstrate its medicinal effect. The ethanolic

QUICK RESPONSE CODE	<p style="text-align: center;">DOI: 10.13040/IJPSR.0975-8232.5(11).4832-37</p> <hr/> <p style="text-align: center;">Article can be accessed online on: www.ijpsr.com</p> <hr/> <p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.5(11).4832-37</p>
---	---

extract of *C. halicacabum* shows anti arthritics activity and it was proven by Ramachandran et.al.⁷. The ethanolic extract of *C. halicacabum* shows anti inflammatory, anti pyretic activity, anti hyperglycaemic effect⁸⁻¹¹. Ethanol extracts also exhibit *in vitro* hydroxyl radical scavenging activities¹². The flavone isolated from *C. halicacabum* was investigated for the antioxidant and hepatoprotective effects in Wistar albino rats¹³. Isolation of anxiolytic principle from ethanolic root extract of *C. halicacabum* shows the presence of Cardiospermin a cyanogenic glucoside present in it¹⁴. Mucilage have been isolated from aerial part of *C. halicacabum* possesses many biological activities¹⁵.

Still there is no report on isolation of active compounds from *C. halicacabum* leaf the present study was undertaken to isolate active compounds from *C. halicacabum* which are responsible for various biological effects. This work will be helpful for the discovery of new drugs in the field of plant medicine.

MATERIALS AND METHODS:

Extraction of Plant Material:

Fresh leaves of *C. halicacabum* was collected from the tropical areas of Mallur Taluk, Salem district, Tamilnadu, India, and thoroughly washed in water, dried at room temperature, and powdered. Dry fine powder was suspended in Hydro alcoholic solvent (80% methanol + 20% distilled water) for cold maceration for 4 days. The extract was filtered using a muslin cloth and concentrated at 40 ± 5 °C.

Chemicals:

All chemicals, solvents and reagents were obtained from E. Merck, HIMEDIA and SD Fine chemicals (India). TLC plates were supplied by Merck, (TLC Silica gel 60 F254).

Qualitative Phytochemical Evaluation:

The crude Hydro alcoholic extracts of the leaves of *C. halicacabum* was subjected to chemical tests for its active constituents according to the Harborne method¹⁶.

DETERMINATION AND ISOLATION OF BIOLOGICALLY ACTIVE COMPOUNDS:

Preparative Thin Layer Chromatography (TLCP)¹⁷⁻²¹:

For Preparative thin layer chromatography (TLCP) separations, glass plates coated (1mm) with silica gel were employed.

Preparation of Thin Layer Chromatography Plates (PTLCP):

25g of silica gel was suspended in 50mL of deionised water and thickened slurry was poured into the glass plates for thickness 1mm and allowed to air dry for 30minutes at 50°C (until they turn white). Then few drops of the hydro alcoholic extract were applied to the bottom of each of the glass plates. After 5 minutes of drying, each plate was placed in the separate glass chamber with solvent system Chloroform: methanol (9:1 v/v) as the mobile phase. The different solvent systems were tried as mobile phase, of which chloroform: methanol (9:1) gave clearly visible spots hence tried with that mobile phase.

Column chromatography (CC):

The extract residue (10 g) was applied onto a silica gel column (200 g, 100cm x 3.5 cm), to isolate the active compound from the crude extracts using hexane as a solvent and the polarity was increased by CHCl₃ then MeOH and fractions (100 mL) each were collected. The obtained fractions were concentrated and monitored by TLC using hexane, chloroform and methanol (97:2:1%) as mobile phase.

Fractions (5-20) considered as Fraction A1 eluted with chloroform-methanol (9.5:0.5). Compound 1 was started to coming out to yield 60 fractions (20 mL each). The solvent evaporated and the fractions were TLC monitored. The R_f value was found to be 0.5. Again, the solvent polarity was increased gradually to elute another spot. Compound 2 was collected to yield 50 fractions (20 mL each).

Fractions (25-36) - Fraction B1 eluted with chloroform-methanol (8:2) was found to contain two spots. Compound 3 was started to coming out to yield 80 fractions (20 mL each). The solvent evaporated and the fractions were TLC monitored. The R_f value was found to be 0.6. Again, the solvent polarity was increased gradually to elute another spot. At 5% addition of methanol, the compound 4 was collected to yield 50 fractions (20 mL each). The solvent evaporated and the fractions

were TLC monitored. The R_f value was found to be 0.6.

From fractions (45-58) C1 compound 5 and 6 were eluted with R_f value 0.6. Fractions (64-76), Fraction D₁ eluted with chloroform-methanol (7:3) were found to have three major spots. Compound 7 was obtained with R_f value 0.7. Compound 8 was obtained by increasing the solvent polarity. At 8% addition of methanol, 50 fractions were obtained each 20 mL with R_f value 0.7. Compound 9 was obtained with R_f value 0.8. Likewise compound 10 and 11 were eluted from Fractions (80-98) E1, and Fractions (105-116) F1 with R_f value 0.8.

The collected fractions were concentrated using a Rotavapour. TLC was used to analyze the active compound, and those with similar chemical components were pooled together. The pure compound was dried in pre-weighed pill vial and its mass (g/ml) determined.

Thin layer chromatography (TLC):

The TLC plates were used to observe the separation of individual compounds as a single spot were trimmed and the position of the origin marked by a straight line. The column fractions of sample was spotted on the origin and put in a lidded tank containing a solvent system. The procedure was followed for other plates with various solvent-solvent ratios (chloroform: methanol in the ratio 6:4, 7:3, 5.5:4.5 and 4:6) until good resolution was noticed. The level of solvent in the tank was about 1cm beneath the origin. The solvent travelled up the plate by capillary action till it reached the solvent front (also marked by a straight line across). The lid was lifted off and the plate is dried before it was viewed by spraying with silver nitrate and iodine vapor and visualized under UV light to identify and confirm the compounds eluted through column chromatography.

Distance travelled by substance

$R_f = \dots\dots\dots$

Distance travelled by solvent front

BIOACTIVITY OF ISOLATED COMPOUNDS:

Total anti-oxidant capacity:

The total antioxidant capacity was determined by phosphormolybdenum method²² and is based on the reduction of Mo (VI) to Mo (V) by the antioxidant

compounds and the formation of a green Mo (V) complex which has the maximal absorption at 695nm. An aliquot of 0.1ml of the sample solution containing a reducing species in DMSO was combined in an eppendorff tube with 1ml of reagent solution (0.6M Sulphuric acid, 28mM sodium phosphate, and 4mM ammonium molybdate). The tubes were capped and incubated in water bath at 95°C for 90min. The samples were cooled to room temperature, and the absorbance of each solution was measured at 695nm. The total antioxidant capacity was expressed as mM equivalent of ascorbic acid²³.

Reducing power Assay (RPA):

A method developed by Oyaizu²⁴, 1986 for reducing power test was used. The above sample including the compounds together with Ascorbic acid solutions were spiked with 2.5ml of phosphate buffer (0.2 M, pH 6.6) and 2.5ml of 1% potassium ferricyanide. The mixture was then kept in a 50°C water-bath for 20min. The resulting solution was then cooled rapidly, spiked with 2.5ml of 10% trichloroacetic acid, and centrifuged at 3000rpm for 10 min. The supernatant (5ml) was then mixed with 5ml of distilled water and 1ml of 0.1% ferric chloride. The absorbance at 700nm was then detected after reaction for 10min. The higher the absorbance represents the stronger the reducing power. The reducing power assay was expressed in terms of Ascorbic acid equivalent per gram of dry weight basis.

DPPH Assay:

The DPPH free radical is reduced to a corresponding hydrazine when it reacts with hydrogen donors²⁵. The DPPH radical is purple in colour and upon reaction with hydrogen donor changes to yellow colour. It is a discoloration assay, which is evaluated by the addition of the antioxidant to a DPPH solution in ethanol or methanol and the decrease in absorbance was measured at 490nm. The assay was carried out in a 96 well microtitre plate. To 200µl of DPPH solution, 10µl of each of the test sample or the standard solution was added separately in wells of the microtitre plate. The final concentration of the test and standard solutions used were 1000, 500, 125 and 31.25 µg/ml. The plates were incubated at 37°C for 30 min and the absorbance of each

solution was measured at 490 nm, using a microplate reader.

RESULTS AND DISCUSSION:

Qualitative phytochemical analysis of Hydro alcoholic extract of *C.halicacabum* showed the presence of bioactive phytochemicals like alkaloids, tannins, flavonoids and terpenoids. Further in bioassay guided isolation eleven compounds were isolated and all the compounds were subjected to antioxidant assays. **Graph 1, 2 and 3** shows the results of total antioxidant capacity assay, reducing power assay and DPPH assay. It was clearly observed that compound 1, 6 and 8 possess maximum antioxidant capacity and minimum IC_{50} value. Compound 1 and 8 were selected for structural elucidation. NMR technique was employed in the identification of structure of compounds. **Graph 4 & 5** shows the NMR results of compound 1 (Kaempferol- 3- O- α - I- rhamnoside, **Figure: 1**) and 8 (Apigenin-7- o- β - d- glucuronide, **Figure: 2**).

GRAPH 1: TOTAL ANTI-OXIDANT ACTIVITY

GRAPH 2: RPA ASSAY

GRAPH 3: DPPH ASSAY

GRAPH 4: NMR RESULT OF COMPOUND 1

GRAPH 5: NMR RESULT OF COMPOUND 1

Figure: 1. Compound- 1 Kaempferol- 3- o- α -l- rhamnoside

Figure: 2. Compound- 8 Apigenin – 7- o- β -d- glucuronide

Kaempferol from *Bryophyllum pinnatum* and Apigenin from *Premna cordifolia* were shown to possess antioxidant activity²⁶ and many other studies showing evidence for the same^{27,28} which supports the current study.

CONCLUSION: In conclusion, in this study we isolated the active compounds from *C. halicacabum* leaf and characterized the same. Since plants contain thousands of constituents with considerably different biological properties any successful investigation of biological material lies in the ability to have as large a number of biological screens as possible and can be effectively used as an excipient in pharmaceutical formulations. Kaempferol- 3- o- α - l- rhamnoside and Apigenin – 7-o- β -d- glucuronide showed strong antioxidant activity. Further, both *in-vivo* and *in-vitro* anti diabetic effect study of these compounds are in progress.

ACKNOWLEDGEMENT: The authors express their gratitude to Ms. V.K. Evanjeline and Mr. Ganesh, M.D, Alpha Omega Hi-tech Laboratory for providing laboratory facilities.

REFERENCES:

1. Bharat Agarwal B and Sahdeo Prasad: Identification of Novel Anti-inflammatory Agents from Ayurvedic Medicine for Prevention of Chronic Diseases. NIH Public Access Author Manuscript 2011; 12(11):1595–1653.
2. Narah M, Kalita Jogen C and Kotoky J: Medicinal plants with potential anticancer activities: A Review. International Research Journal of Pharmacy 2012; 3(6):26-30.
3. Hena JS, Adamu AK, Iortsuun DN and Olonitola OS: Phytochemical screening and antimicrobial effect of the aqueous and methanolic extracts of roots of *Balanites aegyptiaca* on some bacteria species. Science World Journal 2010; 5(2):59-63.
4. Lai ZR, Ho YL, Huang SC, Huang TH, Lai SC, Tsai JC, Wang CY, Huang GJ and Chang YS: Antioxidant, anti-inflammatory and antiproliferative activities of *Kalanchoe gracilis* (L.) DC stem. The American Journal of Chinese Medicine 2011; 39(6):1275-1290
5. Patil VV, Bhangale SC, Chaudhari KP, Kakade RT, Thakare VM, Bonde CG and Patil VR: Evaluation of the antidiarrheal activity of the plant extracts of *Ficus species*. Journal of Chinese Integrative Medicine 2012; 10(3):347-352
6. Uma Shankar M, Murthy PN and Sambit Kumar PT: Analgesic and anti-inflammatory activities of Indian medicinal plant *Ziziphus xylopyrus* stem barks in experimental animal models. Elixir Pharmacy 2012; 44:7265-7270
7. Jeyadevi R and Sivasudha T: Anti-arthritis activity of the Indian leafy vegetable *Cardiospermum halicacabum* in Wistar rats and UPLC–QTOF–MS/MS identification of the putative active phenolic components. Inflammation and Research 2013; 62:115–126
8. Boonmars T, Khunkitti W and Sithithaworn P: *In-vitro* antiparasitic activity of extracts of *Cardiospermum halicacabum* against third-stage larvae of *Strongyloides stercoralis*. Parasitol Research 2005; 95:417-419
9. Asha VV and Pushpangadan P: Antipyretic activity of *Cardiospermum halicacabum*. Indian Journal of Experimental Biology 1999; 37:411-414
10. Makonnen E, Debella E, Zerihun D and Abebe F: Antipyretic properties of the aqueous and ethanol extracts of the leaves of *Ocimum suave* and *Ocimum lamiifolium* in mice. Journal of Ethnopharmacology 2003; 88:85-91
11. Veeramani C, Pushpavalli G and Pugalendi KV: Antihyperglycaemic effect of *Cardiospermum halicacabum* Linn. Leaf extract on STZ-induced diabetic rats. Journal of Applied Biomedicine 2008; 19-26.
12. Sheeba MS and Asha VV: Effect of *Cardiospermum halicacabum* in ethanol induced gastric ulcers in rats. Journal of Ethnopharmacology 2006; 106:105-110
13. Rupesh Kumar M, Kavitha K and Basu SK: Antioxidant and Hepatoprotective Effect of flavanone from *Cardiospermum halicacabum* N. against acetaminophen induced Hepatotoxicity in rats. Journal of Pharmacy Research 2012; 5(1):544-547.
14. Rajesh Kumara G and Murugananthana K: Isolation of anxiolytic principle from ethanolic root extract of *Cardiospermum halicacabum*. Phytomedicine 2011; 18:219–223
15. Fatima Grace X and Latha S: Isolation and characterisation of *Cardiospermum halicacabum* mucilage. International Journal of Green Pharmacy 2011; 314-317
16. Harborne JB: 'Phytochemical Methods'- A guide to modern techniques of plant analysis. Chapman and Hall, New York, Edition 3, 1998:1-150
17. Tomer K, Singh V, Sethiya NK, Singh HP, Kumar M and Chandra D: Isolation and characterization of New Lanosteroid from ethanolic extracts of *Eclipta alba* Linn. Journal of Pharmacy Research 2009; 2(10):1635-1637.
18. Mueen Ahmed KK: Introduction to isolation, identification and estimation of lead compounds from natural products. Textbook of industrial pharmacy. Hiremath SR, Chennai, India, Orient Longman Private Ltd. 2008: 345-379.
19. Hota D: Evaluation of plant extracts. Bioactive medicinal plants, Gene-Tech Books. New Delhi, 2010: 86-87.
20. Patra JK, Gouda S, Sahoo SK and Thatoi HN: Chromatography separation, ¹H NMR analysis and bioautography screening of methanol extract of *Excoecaria agallocha* L. from Bhitarkanika, Orissa, India. Asian Pacific Journal of Tropical Biomedicine 2012; 50-56.
21. Ly TN, Shimoyamada M, Kato K and Yamauchi R: Isolation and characterization of some antioxidative

- compounds from the rhizomes of smaller Galangal (*Alpinia officinarum Hance*). Journal of Agricultural and Food Chemistry 2003; 51:4924-4929.
22. Owen RW, Haubner R, Mier W, Giacosa A, Hull WE, Spiegelhalder B and Bartsch H: Isolation, structure elucidation and antioxidant potential of the major phenolic and flavonoid compounds in brined olive drupes. Food and Chemistry Toxicology 2003; 41:703-717.
 23. Mojca S, Petra K Majda, Aandreja S. Marjana and Knez Z: Phenols, proanthocyanidins, flavones and flavonols in some plant materials and their antioxidant activities. Food Chemistry 2005; 89:191-198.
 24. Oyaizu M: Studies on products of browning reactions: antioxidative activities of products of browning reaction prepared from glucosamine. The Japanese Journal of Nutrition and diabetes 1986; 103:413-419.
 25. Braca A, Tommasi ND, Bari LD, Pizza C, Politi M and Morelli I: Anti oxidant principles from *Bauhinia terapotensis*. Journal of Natural Products 2001; 64:892-895.
 26. Monika Majewska and Michal Skrzycki: Evaluation of antioxidant potential of flavonoids: An *in-vitro* study. Acta Poloniae Pharmaceutica - Drug Research 2011; 8(4):611-615.
 27. Tatsimo SJ and Tamokou Jde D: Antimicrobial and antioxidant activity of kaempferol rhamnoside derivatives from *Bryophyllum pinnatum*. BMC Research notes 2012; 5:158.
 28. Sadasivama K and Kumaresanb R: A comparative DFT study on the antioxidant activity of apigenin and scutellarein flavonoid compounds Molecular Physics. An International Journal at the Interface between Chemistry and Physics 2011; (109):839-852.

How to cite this article:

Rajeswari R and Sridevi M: Isolation of Active Compounds from Hydro Alcoholic Extract of *Cardiospermum Halicacabum* Linn. Int J Pharm Sci Res 2014; 5(11): 4832-37.doi: 10.13040/IJPSR.0975-8232.5 (11).4832-37.

All © 2014 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)