| Monomeric and polymeric species | Number concentrations |
|--|---|
| native functional neuroserpin | [N] |
| latent inactive neuroserpin | [L] |
| intermediate conformer, active for aggregation | [I] |
| polymer made of p monomers | [P] |
| | |
| First order processes: | Rates |
| molecular interconversions among monomeric states | ! |
| activation of native monomers | $\kappa_A[N]$ |
| latentization from native monomers | $\kappa_L[N]$ (2) |
| latentization from intermediates | $\kappa_L^*[I]$ |
| inactivation of intermediates | $\kappa'_A[I]$ |
| latent loss to native monomers | $\kappa_L'[L]$ (3) |
| latent loss to intermediates | $\kappa_L^{\prime*}[L]$ (3) |
| | |
| Second order processes: polymer formation | Rates (4) |
| dimerization from a native and an intermediate monomer | $a_{NI}\kappa_D c[N][I]$ (5) |
| dimerization from two intermediate monomers | $a_{II}\kappa_D c[I][I]$ (5) |
| native monomer addition to polymers of p monomers | $a_{Np}\kappa_M c[N][p] \qquad \qquad (^6)$ |
| intermediate monomer addition to polymers of p monomers | $a_{Ip}\kappa_M c[I][p] \tag{6}$ |
| association of two polymers of i and j monomers | $a_{ij}\kappa_P c[i][j]$ |
| | |
| Other first order processes: depolymerization | Rates |
| native monomer release from polymers of p units | $b_{Np}\kappa_R[p] \tag{7}$ |
| intermediate monomer release from polymers of p units | $b_{Ip}\kappa_R[p]$ (7) |
| fragmentation in two polymers of i and j units | $b_{ij}\kappa_F[i+j]$ |
| (1) The total monomer number concentration is $[1] = [N] + [L] + [I]$ (2) This process has been ruled out by Chiou et al. 2009, Biophys. J. 97,2306. (3) Latent loss can be reasonably neglected since latent conformation is in a deep free energy minimum. (4) c is the total mass concentration. (5) Short notation for dimerization rates: $a_{11} = (a_{NI}[N] + a_{II}[I]) [I][1]^{-2} \kappa_D \kappa_P^{-1}$. (6) Short notation for monomer addition rates: $a_{1p} = (a_{Np}[N] + a_{Ip}[I]) [1]^{-1} \kappa_M \kappa_P^{-1}$. (7) Short notation for monomer release rates: $b_{1p} = (b_{Np} + b_{Ip}) \kappa_R \kappa_F^{-1}$; and for $p = 1$: $b_{1p} = (b_{NN} + b_{NI} + b_{II}) \kappa_R \kappa_F^{-1}$. | |