



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 30, 2020 – 02:21 PM JST

PDB ID : 7D6B
Title : Crystal structure of Oryza sativa Os4BGlu18 monolignol beta-glucosidase with delta-gluconolactone
Deposited on : 2020-09-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	<div> <div>3%</div> <div>96%</div> <div>..</div> </div>
1	B	482	<div> <div>2%</div> <div>96%</div> <div>..</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-glucosidase 18.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	475	Total	C	N	O	S	0	2	0
			3867	2481	654	717	15			
1	B	477	Total	C	N	O	S	0	2	0
			3877	2492	649	720	16			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	24	ALA	-	expression tag	UNP Q7XSK0
A	25	MET	-	expression tag	UNP Q7XSK0
B	24	ALA	-	expression tag	UNP Q7XSK0
B	25	MET	-	expression tag	UNP Q7XSK0

- Molecule 2 is D-glucono-1,5-lactone (three-letter code: LGC) (formula: C₆H₁₀O₆) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	290	Total	O	0	0
			290	290		
5	B	323	Total	O	0	0
			323	323		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-glucosidase 18



- Molecule 1: Beta-glucosidase 18



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.11Å 83.83Å 207.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 29.53 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.9 (30.00-2.10) 98.9 (29.53-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.55 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.145 , 0.177 0.158 , 0.187	Depositor DCC
R_{free} test set	2676 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8478	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, LGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3989	0.68	0/5420
1	B	0.44	0/4003	0.69	2/5441 (0.0%)
All	All	0.44	0/7992	0.69	2/10861 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	185	ARG	NE-CZ-NH1	5.15	122.88	120.30
1	B	313	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3867	0	3660	5	0
1	B	3877	0	3671	7	0
2	A	12	0	10	1	0
2	B	12	0	10	1	0
3	A	54	0	72	0	0
3	B	42	0	56	0	0
4	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	290	0	0	0	0
5	B	323	0	0	0	0
All	All	8478	0	7479	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:GLU:OE2	1:A:323:ARG:NH2	2.35	0.59
1:B:414:GLY:O	1:B:418:TYR:CD1	2.62	0.53
1:B:412:ALA:HB2	1:B:473:PHE:CZ	2.49	0.48
1:B:194:GLU:OE2	2:B:601:LGC:O1	2.31	0.48
1:A:194:GLU:OE2	2:A:601:LGC:O1	2.31	0.48
1:A:412:ALA:HB2	1:A:473:PHE:CZ	2.49	0.48
1:A:263:VAL:HG22	1:A:333:GLY:HA3	1.97	0.47
1:A:103:ARG:HA	1:A:143:PHE:O	2.17	0.44
1:B:414:GLY:O	1:B:418:TYR:CE1	2.70	0.44
1:B:415:GLY:O	1:B:416:ASP:HB2	2.18	0.44
1:B:263:VAL:HG22	1:B:333:GLY:HA3	2.02	0.41
1:B:264:LEU:HB3	1:B:290:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/482 (98%)	458 (96%)	17 (4%)	0	100	100
1	B	477/482 (99%)	463 (97%)	13 (3%)	1 (0%)	47	49
All	All	952/964 (99%)	921 (97%)	30 (3%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	416	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/409 (99%)	402 (100%)	2 (0%)	88	92
1	B	406/409 (99%)	405 (100%)	1 (0%)	93	96
All	All	810/818 (99%)	807 (100%)	3 (0%)	91	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	201	HIS
1	A	257	ARG
1	B	170	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 19 ligands modelled in this entry, 1 is monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GOL	B	604	-	5,5,5	0.29	0	5,5,5	0.63	0
3	GOL	A	602	-	5,5,5	0.43	0	5,5,5	0.60	0
2	LGC	B	601	-	12,12,12	2.15	2 (16%)	15,17,17	1.39	2 (13%)
3	GOL	A	609	-	5,5,5	0.37	0	5,5,5	0.43	0
3	GOL	B	607	-	5,5,5	0.28	0	5,5,5	0.34	0
3	GOL	A	607	-	5,5,5	0.30	0	5,5,5	0.19	0
3	GOL	B	603	-	5,5,5	0.40	0	5,5,5	0.51	0
3	GOL	A	603	-	5,5,5	0.22	0	5,5,5	0.39	0
3	GOL	A	604	-	5,5,5	0.40	0	5,5,5	0.22	0
3	GOL	A	610	-	5,5,5	0.59	0	5,5,5	0.76	0
3	GOL	B	609	-	5,5,5	0.30	0	5,5,5	0.46	0
3	GOL	B	606	-	5,5,5	0.32	0	5,5,5	0.15	0
3	GOL	B	605	-	5,5,5	0.33	0	5,5,5	0.59	0
2	LGC	A	601	-	12,12,12	2.09	2 (16%)	15,17,17	1.42	2 (13%)
3	GOL	A	605	-	5,5,5	0.45	0	5,5,5	0.40	0
3	GOL	A	608	-	5,5,5	0.21	0	5,5,5	0.53	0
3	GOL	B	608	-	5,5,5	0.18	0	5,5,5	0.45	0
3	GOL	A	606	-	5,5,5	0.48	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	604	-	-	2/4/4/4	-
3	GOL	A	602	-	-	2/4/4/4	-
2	LGC	B	601	-	-	0/2/22/22	0/1/1/1
3	GOL	A	609	-	-	2/4/4/4	-
3	GOL	B	607	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	607	-	-	0/4/4/4	-
3	GOL	B	603	-	-	0/4/4/4	-
3	GOL	A	603	-	-	1/4/4/4	-
3	GOL	A	604	-	-	0/4/4/4	-
3	GOL	A	610	-	-	2/4/4/4	-
3	GOL	B	609	-	-	2/4/4/4	-
3	GOL	B	606	-	-	2/4/4/4	-
3	GOL	B	605	-	-	0/4/4/4	-
2	LGC	A	601	-	-	0/2/22/22	0/1/1/1
3	GOL	A	605	-	-	2/4/4/4	-
3	GOL	A	608	-	-	0/4/4/4	-
3	GOL	B	608	-	-	0/4/4/4	-
3	GOL	A	606	-	-	4/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	LGC	O5-C1	6.78	1.44	1.34
2	A	601	LGC	O5-C1	6.53	1.44	1.34
2	A	601	LGC	O5-C5	-2.87	1.42	1.46
2	B	601	LGC	O5-C5	-2.59	1.42	1.46

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	LGC	O5-C1-C2	3.81	124.89	119.20
2	A	601	LGC	O5-C1-O1	-3.70	113.07	118.47
2	B	601	LGC	O5-C1-O1	-3.29	113.67	118.47
2	A	601	LGC	O5-C1-C2	3.17	123.92	119.20

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	604	GOL	O1-C1-C2-C3
3	A	602	GOL	O1-C1-C2-C3
3	B	609	GOL	O1-C1-C2-C3
3	A	610	GOL	C1-C2-C3-O3
3	B	606	GOL	O1-C1-C2-C3
3	A	606	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	A	610	GOL	O2-C2-C3-O3
3	A	609	GOL	C1-C2-C3-O3
3	A	603	GOL	C1-C2-C3-O3
3	A	606	GOL	O1-C1-C2-C3
3	B	609	GOL	O1-C1-C2-O2
3	B	606	GOL	O1-C1-C2-O2
3	A	606	GOL	O2-C2-C3-O3
3	B	604	GOL	O1-C1-C2-O2
3	A	606	GOL	O1-C1-C2-O2
3	A	602	GOL	O1-C1-C2-O2
3	A	605	GOL	O1-C1-C2-O2
3	A	609	GOL	O2-C2-C3-O3
3	A	605	GOL	O2-C2-C3-O3

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	LGC	1	0
2	A	601	LGC	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Ligand LGC B 601







CONFIDENTIAL
















Ligand LGC A 601



CONFIDENTIAL





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

CONFIDENTIAL VALIDATION REPORT

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	475/482 (98%)	-0.15	14 (2%) 51 57	12, 18, 34, 54	0
1	B	477/482 (98%)	-0.21	12 (2%) 57 62	13, 18, 29, 55	0
All	All	952/964 (98%)	-0.18	26 (2%) 54 60	12, 18, 32, 55	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	117	PHE	4.7
1	B	418	TYR	4.0
1	A	370	LEU	3.7
1	B	417	SER	3.4
1	A	368	ASN	3.2
1	A	219	HIS	2.9
1	B	416	ASP	2.9
1	A	416	ASP	2.9
1	A	369	GLY	2.8
1	B	415	GLY	2.8
1	A	417	SER	2.7
1	B	500	ASN	2.6
1	A	415	GLY	2.5
1	B	68	PRO	2.4
1	A	263	VAL	2.4
1	B	419	THR	2.4
1	B	69	GLY	2.3
1	B	422	GLU	2.3
1	B	499	GLN	2.2
1	A	26	ALA	2.2
1	B	368	ASN	2.2
1	A	264	LEU	2.1
1	A	222	ARG	2.1
1	A	116	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	35	SER	2.1
1	B	222	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	605	6/6	0.77	0.25	40,43,47,52	0
3	GOL	A	608	6/6	0.79	0.26	36,38,41,42	0
3	GOL	A	605	6/6	0.80	0.27	40,43,45,47	0
3	GOL	A	610	6/6	0.81	0.20	40,45,47,49	0
3	GOL	A	609	6/6	0.81	0.21	43,45,47,48	0
3	GOL	B	609	6/6	0.82	0.30	48,50,51,52	0
3	GOL	B	603	6/6	0.83	0.19	35,40,42,44	0
3	GOL	B	606	6/6	0.85	0.23	42,46,48,48	0
3	GOL	B	608	6/6	0.86	0.25	23,29,30,34	0
3	GOL	A	602	6/6	0.89	0.22	30,32,33,36	0
3	GOL	A	603	6/6	0.89	0.21	37,41,44,47	0
3	GOL	A	607	6/6	0.90	0.13	29,33,34,35	0
3	GOL	A	606	6/6	0.91	0.18	30,34,37,38	0
3	GOL	B	604	6/6	0.93	0.12	25,26,26,27	0
3	GOL	A	604	6/6	0.95	0.14	19,20,21,21	0
3	GOL	B	607	6/6	0.96	0.12	19,21,22,22	0
2	LGC	B	601	12/12	0.97	0.18	14,15,15,17	0
2	LGC	A	601	12/12	0.98	0.15	12,13,13,15	0
4	ZN	B	602	1/1	1.00	0.04	16,16,16,16	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around GOL A 608:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






CONFIDENTIAL

Electron density around GOL A 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GOL A 610:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GOL A 609:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL B 609:




$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL VALIDATION

Electron density around GOL B 603:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL B 606:




$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL B 608:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL A 602:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL A 603:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL A 607:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL A 606:




$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL VALIDATION

Electron density around GOL B 604:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL A 604:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around GOL B 607:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around LGC B 601:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around LGC A 601:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



CONFIDENTIAL

Electron density around ZN B 602:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**6.5 Other polymers** ⓘ

There are no such residues in this entry.