Arithmetic occult period maps

Jeffrey D. Achter

ABSTRACT

Several natural complex configuration spaces admit surprising uniformizations as arithmetic ball quotients, obtained by identifying each parametrized object with the periods of some auxiliary object. In each case, the theory of canonical models of Shimura varieties gives the ball quotient the structure of a variety over the ring of integers of a cyclotomic field. We show that the (transcendentally defined) period map actually respects these algebraic structures, and thus that occult period maps are arithmetic. As an intermediate tool, we develop an arithmetic theory of lattice-polarized K3 surfaces.

1. Introduction

It occasionally happens that complex varieties of a specified type are parametrized by an arithmetic quotient of a unit ball in a surprising way. We situate this remark by recalling some aspects of the primordial period map. Consider M_g , the moduli space of smooth projective curves of genus $g \geq 2$. Given a smooth projective curve of genus g , the possibilities for its period lattice are naturally parametrized by the quotient of $\mathbb{H}_g$, the Siegel upper-half space of dimension $g(g+1)/2$, by $\mathrm{Sp}_{2g}(\mathbb{Z})$. The classical Torelli theorem asserts that the corresponding map $\tau_{g,\mathbb{C}}: M_g(\mathbb{C}) \rightarrow \mathbb{H}_g/\mathrm{Sp}_{2g}(\mathbb{Z})$ is an inclusion. Even more is true. On one hand, M_g has a natural structure as a moduli space over $\mathbb{Z}$. On the other hand, let A_g be the moduli space of principally polarized abelian varieties of dimension g ; it, too, is a space over $\mathbb{Z}$. Via the identification $A_g(\mathbb{C}) \cong \mathbb{H}_g/\mathrm{Sp}_{2g}(\mathbb{Z})$, we endow the latter with a structure over $\mathbb{Z}$, as well. The key arithmetic fact about the Torelli map is that $\tau_{g,\mathbb{C}}$, a priori a transcendental map, descends to a morphism $\tau_g: M_g \hookrightarrow A_g$ over $\mathbb{Z}$ (for example, [MFK94, §7.4]). Still, $\dim M_g < \dim A_g$ as soon as $g > 3$. This means that many of the arithmetic structures on A_g , such as Hecke operators and modular forms, do not readily make sense for the moduli space of curves.

In the special case where $g = 4$, however, we have the intriguing observation of Kondō [Kon02] that $M_4(\mathbb{C})$ is very close to an arithmetic quotient of $\mathbb{B}^9$, the complex unit 9-dimensional ball. Slightly more precisely, let N_4 denote the (open, dense) locus of nonhyperelliptic curves. Kondō shows that there exist an arithmetic group Γ of automorphisms of $\mathbb{B}^9$ and an open immersion $N_4(\mathbb{C}) \hookrightarrow \mathbb{B}^9/\Gamma$. (He even characterizes the image.) Instead of analyzing the periods of a nonhyperelliptic curve C , the construction of [Kon02] proceeds by constructing an auxiliary variety Z associated with C and analyzing *its* periods. Kudla and Rapoport [KR12]—who call

---

Received 1 May 2019, accepted in final form 28 November 2019.

2010 Mathematics Subject Classification 14J10 (primary), 11G18, 14D05 (secondary).

Keywords: K3 surface, Shimura variety, period map.

This journal is © **Foundation Compositio Mathematica** 2020. This article is distributed with Open Access under the terms of the **Creative Commons Attribution Non-Commercial License**, which permits non-commercial reuse, distribution, and reproduction in any medium, provided that the original work is properly cited. For commercial re-use, please contact the **Foundation Compositio Mathematica**.

This work was partly supported by grants from the Simons Foundation (637075) and the National Security Agency (H98230-14-1-0161, (15/16)-1-0247).

such a period map *occult*, in recognition of its hidden nature—observe that the theory of canonical models of Shimura varieties produces a distinguished algebraic model of $\mathbb{B}^9/\Gamma$ over $\mathbb{Q}(\zeta_3)$. They prove that Kondō’s occult period map actually respects the structures of N_4 and $\mathbb{B}^9/\Gamma$ as varieties over $\mathbb{Q}(\zeta_3)$, and conjecture that it extends to a map of integral canonical models over $\mathbb{Z}[\zeta_3, 1/3]$. (They also note certain stack-theoretic issues, which have since been resolved by Zheng [Zhe20]; see Remark 7.6 below.)

In fact, several different situations are known in which, for some moduli space V of low-complexity varieties, an occult period map yields an open immersion $\tau_{V,\mathbb{C}}: V(\mathbb{C}) \hookrightarrow \mathbb{B}^{\dim V}/\Gamma_V$; see, for instance, [DK07] and [KR12], or even Section 7 below, for examples. In each case known to the author, the theory of integral canonical models of Shimura varieties provides a distinguished model $\mathrm{Sh}_{\Gamma_V}(\mathbb{B}^{\dim V})$ of $\mathbb{B}^{\dim V}/\Gamma_V$ over $\mathbb{Z}[\zeta_n, 1/n]$ for some $n = n(V)$. The goal of the present paper is to show that, in many cases, $\tau_{V,\mathbb{C}}$ descends to a morphism $V \rightarrow \mathrm{Sh}_{\Gamma_V}(\mathbb{B}^{\dim V})$ over $\mathbb{Z}[\zeta_n, 1/2n]$.

Many of the original constructions involve somehow building a K3 surface out of the original variety, and then analyzing the periods of the corresponding K3 surface. Consequently, much of the work of the present paper is in analyzing moduli spaces $R_{L,\underline{\chi}}$ of K3 surfaces polarized by the lattice L and equipped with a suitable action of μ_n . A representative result—the notation is defined later in the paper—is the following.

PROPOSITION. *There are morphisms of stacks over $\mathbb{Z}[\zeta_3, 1/6]$*

$$\begin{array}{ccc} R_{L_4, \underline{\chi}_4}^\circ & \xrightarrow{\kappa_4} & N_4 \\ \downarrow \tau_{L_4, \underline{\chi}_4} & & \\ \mathrm{Sh}^{(L_4, \underline{\chi}_4)} & & \end{array}$$

where κ_4 induces an isomorphism of coarse moduli spaces and $\tau_{L_4, \underline{\chi}_4}$ induces an open immersion $R_{L_4, \underline{\chi}_4}^\circ(\mathbb{C}) \hookrightarrow \mathrm{Sh}^{(L_4, \underline{\chi}_4)}(\mathbb{C})$.

The statement over $\mathbb{C}$ is, essentially, [Kon02, Theorem 1]; taking fibers over $\mathbb{Q}(\zeta_3)$ recovers the descent result [KR12, Theorem 8.1].

2. Notation on lattices

2.1 Lattices

Throughout this paper, a *lattice* is a free $\mathbb{Z}$ -module L of finite rank, equipped with a nondegenerate symmetric bilinear pairing $(\cdot, \cdot)$ (often notationally suppressed). For any nonzero n , we let $L(n)$ denote the lattice with the same underlying group structure as L and with pairing $(\cdot, \cdot)_{L(n)} = n(\cdot, \cdot)_L$. We follow the conventions of [Huy16] for lattices. Lattices used here include

- U the hyperbolic plane, which has rank 2 and pairing $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
- $\langle 1 \rangle$ the lattice of rank 1 and pairing (1)
- E_8 the unique positive definite unimodular lattice of rank 8
- A_n, D_n the (positive) lattice associated with the Dynkin diagrams of type A_n and D_n , respectively (in particular, $A_1 \cong \langle 2 \rangle$)
- L_{K3} the lattice $U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$, of signature $(3^+, 19^-)$
- V the lattice of rank 2 and pairing $\begin{pmatrix} 2 & 1 \\ 1 & -2 \end{pmatrix}$.

The pairing induces an inclusion $L \hookrightarrow L^\vee$; the discriminant of L is the finite abelian group $\text{disc}(L) = L^\vee/L$, and we set $\Delta_L = [L^\vee : L] = \# \text{disc}(L)$. Finally, let

$$d(L) = \gcd(\{d \in \mathbb{N} : \exists \langle 2d \rangle \hookrightarrow L \text{ primitive}\}).$$

For use in Section 7, we record the following elementary facts.

LEMMA 2.1. (a) *If M is a primitive sublattice of L , then $d(L) \mid d(M)$.*

(b) *We have $d(U(n)) = n$, while $d(A_1 \oplus A_1(-1)^{\oplus 2}) = d(V \oplus A_4(-1)) = 1$.*

2.2 Orthogonal groups

With L , we associate the orthogonal group O_L , with connected component of identity the special orthogonal group SO_L . Since we start with an integral model for SO as the automorphism group of the lattice L , we have a natural definition of $SO_L(R)$ for any ring R . In particular, $SO_L(\mathbb{Z}_p)$ is well defined and, by definition, a hyperspecial subgroup of $SO_L(\mathbb{Q}_p)$. (In Section 5.1, a choice of hyperspecial subgroup is necessary for the construction of a canonical integral model of the relevant Shimura variety. In fact, since SO_L is adjoint and split, $SO_L(\mathbb{Q}_p)$ admits a unique $SO_L(\mathbb{Q}_p)$ -conjugacy class of hyperspecial subgroups.)

We will often have cause to work with a lattice L which comes equipped with a primitive embedding $\iota: L \hookrightarrow L_{K3}$. With a slight abuse of notation, we will write $L^\perp$ for the orthogonal complement $\iota(L)^\perp$ of $\iota(L)$ in L_{K3} . Set $O^L = O_{L^\perp}$ and $SO^L = SO_{L^\perp}$. An element of $O^L(\mathbb{Z})$ extends to an element of $O_{L_{K3}}(\mathbb{Z})$ acting trivially on L if and only if it acts trivially on $\text{disc}(L)$ (see, for example, [Huy16, Proposition 14.2.6]). More generally, if R is flat over $\mathbb{Z}$, then an element of $O^L(R)$ extends to an element of $O_L(R)$ acting trivially on $L \otimes R$ if and only if it acts trivially on $\text{disc}(L) \otimes R$; see [Mad16, Lemma 2.6]. The subgroup of admissible orthogonal automorphisms of $L^\perp$ is the group scheme $\tilde{O}^L$ fitting in the short exact sequence

$$1 \longrightarrow \tilde{O}^L \longrightarrow O^L \longrightarrow \text{Aut}(\text{disc}(L)) \longrightarrow 1;$$

on points, we have

$$\begin{aligned} \tilde{O}^L(R) &= \{g|_{L^\perp \otimes R} : g \in O_{L_{K3}}(R)\} \\ &= \ker(O^L(R) \rightarrow \text{Aut}(\text{disc}(L))(R)). \end{aligned}$$

If $\tilde{g} \in \tilde{O}^L(R)$, then there is a (necessarily unique) $g \in O_{L_{K3}}(R)$ such that $g|_{L^\perp} = \tilde{g}$ and $g|_L = \text{id}_L$. In this way, $\tilde{O}^L(R)$ is naturally identified with a subgroup of $O_{L_{K3}}(R)$. The group scheme $\widetilde{SO}^L := \tilde{O}^L \times_{O^L} SO^L$ represents admissible automorphisms of determinant 1.

3. Families of K3 surfaces

3.1 K3 surfaces

Let k be an algebraically closed field. A K3 surface over k is a smooth complete irreducible surface Z/k with trivial canonical bundle $\omega_Z := \Omega_{Z/k}^2 \cong \mathcal{O}_Z$ and such that $H^1(Z, \mathcal{O}_Z) = 0$. Like any smooth complete surface, a K3 surface is projective, and its middle Hodge numbers are $(h^{2,0}, h^{1,1}, h^{0,2}) = (1, 20, 1)$. If $k = \mathbb{C}$, then the Betti cohomology group $H^2(Z, \mathbb{Z})$, endowed with the intersection pairing $(\cdot, \cdot)$, is isomorphic to L_{K3} . The natural pairing

$$\Omega_Z \times \Omega_Z \longrightarrow \omega_Z$$

induces an isomorphism of $\mathcal{T}_Z \cong \Omega_Z$. Since $H^1(Z, \mathcal{O}_Z)$ is trivial, so is $\text{Pic}^0(Z)$, and the Néron–Severi group $\text{NS}(Z)$ coincides with the Picard group $\text{Pic}(Z)$. As for any surface, $\text{NS}(Z)$ is a free finitely generated $\mathbb{Z}$ -module, equipped with a symmetric nondegenerate pairing

$$\text{NS}(Z) \times \text{NS}(Z) \xrightarrow{(\cdot, \cdot)} \mathbb{Z}.$$

The intersection pairing is even, nondegenerate, and of signature $(1, \text{rank}(\text{NS}(Z)) - 1)$.

Following Rizov [Riz06] and successors, we say that a relative K3 surface, or K3 space, over a scheme S is an algebraic space $Z \rightarrow S$ such that each geometric fiber is a K3 surface. If $Z \rightarrow S$ is a relative K3 surface, then $\mathcal{H}_{\text{dR}}^2(Z/S)$ and $\mathcal{H}^{2,0}(Z/S)$ are locally free sheaves on S (see [DI81, Proposition 2.2], [Mad15, § 3.4]) of respective ranks 22 and 1.

3.2 Categories of K3 surfaces

We will study three different sorts of moduli spaces of K3 surfaces.

Classically, one has R_{2d}° , the category of K3 surfaces equipped with a primitive ample polarization of degree $2d$. On points, $\text{R}_{2d}^\circ(S)$ is the category of pairs $(Z \rightarrow S, \lambda)$, where $Z \rightarrow S$ is a K3 space and $\lambda \in \text{Pic}_{Z/S}(S)$ is étale-locally represented by an ample line bundle of self-intersection degree $2d$ which is not a nontrivial tensor power of any other line bundle. This is a subcategory of R_{2d} , the category of K3 surfaces equipped with a primitive quasi-ample polarization of degree $2d$. (A quasi-ample, or pseudo-ample, polarization is étale-locally a line bundle which is big and nef.)

Choose a generator e_0 for $\langle 2d \rangle$. To specify data $(Z \rightarrow S, \lambda)$ is to specify a K3 space $Z \rightarrow S$ and an embedding of lattices $\langle 2d \rangle \hookrightarrow \text{Pic}_{Z/S}(S)$ which takes e_0 to the class of an ample line bundle. (Recall that if Z/k is a K3 surface over an algebraically closed field and if $v \in \text{Pic}_{Z/k}(k)$ satisfies $(v, v) > 0$, then exactly one of v and $-v$ represents the class of an ample line bundle. The choice of a generator for $\langle 2d \rangle$ is equivalent to the choice of a “positive cone” in $\langle 2d \rangle \otimes \mathbb{R} \cong \mathbb{R}$.)

More generally, we consider lattice-polarized K3 surfaces. Let L be a primitive sublattice of L_{K3} of signature $(1, r - 1)$. The set

$$\{v \in L \otimes \mathbb{R} : (v, v) > 0\}$$

has two connected components. Choose one such, V^+ , and, as in [Dol96, § 1] or [DK07, § 10], define an abstract “ample cone” $\mathcal{C}(V^+)$, an open subset of V^+ , and let $L^+ = L \cap \mathcal{C}(V^+)$. We suppress the choice of V^+ (and, thus, L^+) from the notation, and let R_L° be the category of ample L -polarized K3 surfaces. Objects in R_L° are isomorphism classes of pairs $(Z \rightarrow S, \alpha)$, where $Z \rightarrow S$ is a K3 space and $\alpha: L \hookrightarrow \text{Pic}_{Z/S}(S)$ is a primitive embedding of lattices such that $\alpha(L^+)$ contains the class of an ample line bundle. (Since α is a primitive embedding, it is equivalent to ask that $\alpha(\mathcal{C}(V^+))$ contain such a class.) Here, we declare that two such data $(Z_i \rightarrow S, \alpha_i)$ are isomorphic if there is an isomorphism $f: Z_1 \rightarrow Z_2$ such that $f^*\alpha_2 = \alpha_1$. We define R_L , the category of L -polarized K3 surfaces, analogously except that it is only assumed that $\alpha(L^+)$ contains the class of a quasi-ample line bundle.

(Of course, one can also make the definition of an L -polarized K3 surface without keeping track of a positive cone, provided that one is willing to identify α and $-\alpha$. For a Shimura-theoretic justification for this approach, see Remark 6.3 and, ultimately, [Tae17, § 5]. Consequently, the choice of L^+ is suppressed here.)

For a finite group scheme G , let $\text{R}_{L,G}^*$ be the category of tuples $(Z \rightarrow S, \alpha, \rho)$ where $(Z \rightarrow S, \alpha) \in \text{R}_L(S)$ and $\rho: G_S \hookrightarrow \text{Aut}_S(Z \rightarrow S, \alpha)$ is a monomorphism of group schemes. If $\#G$

is invertible on S —equivalently, if the cardinality of G is relatively prime to the characteristic exponent of all residue fields of points of S —then representations of G on $\mathcal{O}_S$ -modules are rigid, and thus $\mathcal{H}_{\mathrm{dR}}^2(Z/S)$ and $\mathcal{H}^{2,0}(Z/S)$ are locally free sheaves of $\mathcal{O}_S[G]$ -modules.

We now specialize to the case $G = \mu_n$ and restrict to the category of schemes over $\mathbb{Z}[1/2\Delta_L n]$. Let χ^ω be a faithful character of μ_n , let χ_0 be the trivial character, and let χ be an arbitrary character; let $m(\chi^\omega) = m_\chi(\chi^\omega)$ and $m(\chi_0) = m_\chi(\chi_0)$ be the multiplicities of, respectively, χ_0 and χ^ω in χ . Let $\mathbf{R}_{L, \mu_n, \chi^\omega, \chi}$ be the open and closed substack of $\mathbf{R}_{L, \mu_n}^*$ parametrizing those $(Z \rightarrow S, \alpha, \rho)$ such that

- μ_n acts on $\mathcal{H}^{2,0}(Z/S)$ via χ^ω ,
- μ_n acts on $\mathcal{H}^2(Z/S)$ via χ , and
- $m_\chi(\chi_0) = \mathrm{rank}(L)$.

In particular, the action of μ_n is purely nonsymplectic, in the sense that no nontrivial section of μ_n fixes a nonzero holomorphic 2-form.

Suppose that S is irreducible and $\bar{s}$ is a geometric point of S . Because $2n$ is invertible on S , representations of μ_n on $\mathcal{O}_S$ -modules are rigid. In particular, the character of the action of μ_n on $\mathcal{H}^2(Z/S)$ is determined by the action on $\mathcal{H}_{\mathrm{dR}}^2(Z_{\bar{s}})$. Moreover, it is equivalent to specify this character in terms of the action of μ_n on $H_{\mathrm{cris}}^2(Z_{\bar{s}})$ or any of the étale cohomology groups $H^2(Z_{\bar{s}}, \mathbb{Q}_\ell)$ (see [KM74, Theorem 2.2]) or $H^2(Z_{\bar{s}}, \mathbb{Z}_\ell)$ (since K3 surfaces have torsion-free cohomology).

We will often use the symbol $\underline{\chi}$ to denote the collection of data $(\mu_n, \chi^\omega, \chi)$, and thus write $\mathbf{R}_{L, \underline{\chi}}$ for $\mathbf{R}_{L, \mu_n, \chi^\omega, \chi}$, etc.

The possibilities for data $(L, \underline{\chi})$ such that $\mathbf{R}_{L, \underline{\chi}}(\mathbb{C})$ is nonempty are reasonably well understood. On one hand, it is not hard to see that a purely nonsymplectic group of automorphisms is finite and cyclic; the possible orders of such a group are also known [MO98]. On the other hand, starting with the work of Nikulin, one has a good classification of primitive sublattices of L_{K3} ; see [Nik79]. In Section 7 we will see a number of explicit examples of naturally occurring families of lattice-polarized K3 surfaces.

3.3 Stacks of K3 surfaces

It turns out that each $\mathbf{R}_{2d}$, $\mathbf{R}_L$, and $\mathbf{R}_{L, \underline{\chi}}$ is a Deligne–Mumford stack. Indeed, Rizov proves that $\mathbf{R}_{2d}^\circ$ is a Deligne–Mumford stack [Riz06, Theorem 4.3.3], and Beauville essentially proves the same of $\mathbf{R}_L$ in [Bea04]. The partial compactification $\mathbf{R}_{2d}$ of $\mathbf{R}_{2d}^\circ$ is also Deligne–Mumford and even smooth over $\mathbb{Z}[1/2d]$, see [Mau14, Proposition 2.1], albeit no longer separated (see, for example, [Huy16, §5.1.4]). Rather than working ab ovo to study $\mathbf{R}_L$ and $\mathbf{R}_{L, G}^*$, we find it expedient to bootstrap from Rizov’s work.

It is convenient to make at the outset a few (arbitrary) choices; the final claims are intrinsic and independent of these choices. Let $e_1, \dots, e_r$ be a $\mathbb{Z}$ -basis for L . Fix some $\lambda \in L^+$.

LEMMA 3.1. *The category $\mathbf{R}_L$ is a stack over $\mathrm{Spec} \mathbb{Z}$.*

Proof. We must show that the diagonal $\mathbf{R}_L \rightarrow \mathbf{R}_L \times \mathbf{R}_L$ is representable and that étale descent in the category $\mathbf{R}_L$ is effective.

For the first claim, it suffices to show that if $(Z_1 \rightarrow S, \alpha_1)$ and $(Z_2 \rightarrow S, \alpha_2)$ are elements of $\mathbf{R}_L(S)$, then

$$\mathrm{Isom}((Z_1, \alpha_1), (Z_2, \alpha_2))$$

is representable by a scheme over S . The functor $\mathbf{Isom}(Z_1, Z_2)$ is represented by a scheme over S . (In fact, it is open in $\mathrm{Hilb}(Z_1 \times Z_2)$.) Pulling back by isomorphisms gives a pairing $\mathbf{Isom}(Z_1, Z_2) \times_S \mathrm{Pic}_{Z_2/S} \rightarrow \mathrm{Pic}_{Z_1/S}$. Consider some i between 1 and r . Pulling back the pairing by the section $\alpha_2(e_i): S \rightarrow \mathrm{Pic}_{Z_2/S}$ induces a morphism $\beta_i: \mathbf{Isom}(Z_1, Z_2) \rightarrow \mathrm{Pic}_{Z_1/S}$. Then

$$\mathbf{Isom}((Z_1, \alpha_1(e_i)), (Z_2, \alpha_2(e_i))) := \mathbf{Isom}(Z_1, Z_2) \times_{\mathrm{Pic}_{Z_1/S}, \alpha_1(e_i)} S$$

is the subscheme of $\mathbf{Isom}(Z_1, Z_2)$ parametrizing those isomorphisms which take $\alpha_2(e_i)$ to $\alpha_1(e_i)$. Insofar as $\mathbf{Isom}((Z_1, \alpha_1), (Z_2, \alpha_2))$ is the fiber product over $\mathbf{Isom}(Z_1, Z_2)$ of the r different schemes $\mathbf{Isom}((Z_1, \alpha_1(e_i)), (Z_2, \alpha_2(e_i)))$, it too is represented by a scheme.

For the second, let $T \rightarrow S$ be étale, and let $(\tilde{Z} \rightarrow T, \tilde{\alpha}) \in \mathbf{R}_L(T)$ be equipped with T/S descent data. In [Riz06, Lemma 4.3.7], the author shows that in the ample case, $(\tilde{Z} \rightarrow T, \alpha(\lambda))$ descends, as a polarized K3 space, to S ; the quasi-polarized case follows from [Mau14, § 2]. Since $\mathrm{Pic}_{Z/S}$ is a sheaf in the étale topology, each $\tilde{\alpha}(e_i)$ descends to Z/S . $\square$

LEMMA 3.2. *The category $\mathbf{R}_L$ is a Deligne–Mumford stack over $\mathrm{Spec} \mathbb{Z}$.*

Proof. Because $\mathbf{R}_{2d(\lambda)}$ is known to be a Deligne–Mumford stack, it suffices to show that the forgetful morphism

$$\begin{aligned} \mathbf{R}_L &\xrightarrow{\phi_\lambda} \mathbf{R}_{2d(\lambda)}, \\ (Z \rightarrow S, \alpha) &\longmapsto (Z \rightarrow S, \alpha(\lambda)) \end{aligned}$$

is relatively representable [LM00, Proposition 4.5(ii)]. Now proceed as in [Bea04]. Let $H_\lambda \subset \mathrm{O}_L(\mathbb{Z})$ be the subgroup which stabilizes λ . Since $\lambda^\perp$ is negative definite, H_λ is finite. Given an S -point $S \rightarrow \mathbf{R}_{2d(\lambda)}$,

$$\mathbf{R}_L \times_{\phi_\lambda, \mathbf{R}_{2d(\lambda)}} S,$$

if nonempty, is a torsor under H_λ , and in particular is representable. $\square$

PROPOSITION 3.3. *The category $\mathbf{R}_L$ is a smooth Deligne–Mumford stack over $\mathrm{Spec} \mathbb{Z}[1/2\Delta_L]$ of relative dimension $20 - r$.*

Proof. Given Lemma 3.2, it suffices to show that the local deformation space of a quasi-ample L -polarized K3 surface is smooth. In characteristic zero, this is asserted in [Dol96, Proposition 2.1], and details are provided in [Bea04, Proposition 1.4]. In positive characteristic, this follows from Deligne and Illusie’s deformation theory; see Proposition 3.8 below. $\square$

Now consider the moduli spaces of lattice-polarized K3 surfaces with group action.

LEMMA 3.4. *Suppose $(Z \rightarrow S, \alpha) \in \mathbf{R}_L(S)$. Then $\mathrm{Aut}(Z \rightarrow S, \alpha)$ is represented by a proper finite group scheme over S .*

Proof. The automorphism functor $\mathrm{Aut}_S(Z)$ is represented by a separated unramified group scheme over S ; see [Riz06, Theorem 3.3.1]. Moreover, $\mathrm{Aut}_S(Z \rightarrow S, \alpha(\lambda))$ is a closed finite subgroup scheme of $\mathrm{Aut}_S(Z)$ (see [Riz06, Proposition 3.3.3] for the polarized case; the extension to quasi-polarizations follows from the proof of [Mau14, Proposition 2.8]). As in the proof of Lemma 3.1, the group scheme $\mathrm{Aut}_S(Z \rightarrow S, \alpha)$ is a sub- S -group scheme of $\mathrm{Aut}_S(Z \rightarrow S, \alpha(\lambda))$. The claimed properness follows from [MM64, Theorem 2]. $\square$

LEMMA 3.5. *The forgetful morphism $\mathbf{R}_{L,G}^* \rightarrow \mathbf{R}_L$ is finite, and $\mathbf{R}_{L,G}^*$ is a Deligne–Mumford stack.*

Proof. First, the forgetful functor $R_{L,G}^* \rightarrow R_L$ is relatively representable. Indeed, for any affine scheme S and any $(Z \rightarrow S, \alpha) \in R_L(S)$, both G_S and $\text{Aut}_S(Z \rightarrow S, \alpha)$ are relatively representable, and thus $\text{Hom}(G_S, \text{Aut}_S(Z \rightarrow S, \alpha))$ is representable, too; and the condition that a homomorphism be injective is open. Therefore, $R_{L,G}^*$ is also a Deligne–Mumford stack. The properness (and finitude) in Lemma 3.4 imply that $R_{L,G}^* \rightarrow R_L$ is proper and quasifinite, thus finite. $\square$

PROPOSITION 3.6. *The category $R_{L,\chi}$ is a smooth Deligne–Mumford stack over $\mathbb{Z}[\zeta_n, 1/6\Delta_L n]$ of relative dimension $m(\chi^\omega) - 1$.*

Proof. All that needs to be checked is the smoothness; this is done in Lemma 3.9. $\square$

3.4 Local calculations

If $Z/\mathbb{C}$ is a complex K3 surface, then (the local Torelli theorem asserts that) the deformation theory of Z is well captured by its Hodge theory. In particular, let $\text{Def}(Z)$ be the deformation functor of Z , with base point s . Then there is a canonical isomorphism

$$T_s \text{Def}(Z) \xrightarrow{\sim} \text{Hom}(H^{2,0}(Z), H^{2,0}(Z)^\perp / H^{2,0}(Z)).$$

There is a parallel deformation theory for K3 surfaces in arbitrary characteristic, which we review here. Let k be an algebraically closed field of characteristic $p > 0$, with ring of Witt vectors $W = W(k)$.

Let Z/k be a K3 surface. The deformation functor $\text{Def}(Z)$ is formally smooth over $\text{Spf } W$ of relative dimension 20; it is pro-represented by a formal scheme noncanonically isomorphic to $\text{Spf } W[[t_1, \dots, t_{20}]]$; see [DI81, Corollary 1.2], [Riz06, Proposition 4.1.1]. Let s be the base point of $\text{Def}(Z)$, corresponding to Z/k itself.

LEMMA 3.7. *There is a canonical isomorphism of k -vector spaces*

$$T_s \text{Def}(Z) \xrightarrow{\sim} \text{Hom}(\text{Fil}^2 H_{\text{dR}}^2(Z/k), \text{Gr}^1 H_{\text{dR}}^2(Z/k)).$$

Sketch of a proof. See [DI81, Proposition 2.4], [NO85, Theorem 5.2], and [Ogu84, Theorem 5.1]. Briefly, let A be a nilpotent extension of k with a divided power structure. The intersection pairing $(\cdot, \cdot)$ extends to a pairing on the crystal $H_{\text{cris}}^2(Z)$. To give a deformation of Z to A is to lift $\text{Fil}^2 H_{\text{cris}}^2(Z)(k)$ to an isotropic direct summand of $H_{\text{cris}}^2(Z)(A)$. Now use the fact [DI81, equation (2.3.7)] that the orthogonal complement to $\text{Fil}^2 H_{\text{cris}}^2(Z)(k)$ is $\text{Fil}^1 H_{\text{cris}}^2(Z)(k)$. $\square$

If $2d$ is invertible in k and $(Z, \lambda) \in R_{2d}(k)$, then $\text{Def}(Z, \lambda) \subset \text{Def}(Z)$ is pro-representable and formally smooth of dimension 19 over $\text{Spf } W$; see [DI81, Proposition 1.5 and Theorem 1.6], [Mad15, § 3.8], and [Riz06, Lemma 4.1.3]. More generally, we have the following.

PROPOSITION 3.8. *Let L be a lattice of rank r and discriminant Δ_L , and suppose that Δ_L is invertible in k . Let $(Z/k, \alpha) \in R_L(k)$ be an L -polarized K3 surface. Then $\text{Def}(Z, \alpha)$ is pro-representable and formally smooth of dimension $20 - r$ over $\text{Spf } W$.*

Proof. As in Section 3.3, let $e_1, \dots, e_r$ be a $\mathbb{Z}$ -basis for L , and let $\mathcal{L}_i = \alpha(e_i) \in \text{Pic}(Z)$. Then $\text{Def}(Z, \alpha) = \text{Def}(Z, \{\mathcal{L}_1, \dots, \mathcal{L}_r\})$ is the largest formal subscheme of $\text{Def}(Z)$ to which each of the line bundles $\mathcal{L}_i$ extends. Thus, $\text{Def}(Z, \alpha)$ is the scheme-theoretic intersection of the $\text{Def}(Z, \mathcal{L}_i)$. Now, each $\text{Def}(Z, \mathcal{L}_i)$ is the vanishing locus in $\text{Def}(Z)$ of a single function f_i ; see [DI81, Proposition 1.5]. So (any component of) $\text{Def}(Z, \alpha)$ has codimension at most r in $\text{Def}(Z)$, and it suffices to show that the dimension of the tangent space of $\text{Def}(Z, \alpha)$ at the base point is exactly $20 - r$.

Since Z is smooth and proper, there is a crystalline Chern class map $c_1: \mathrm{NS}(Z) \rightarrow H_{\mathrm{cris}}^2(Z/W)$. Moreover, since the crystalline cohomology of Z is torsion-free and the Hodge to de Rham spectral sequence for $H^\bullet(Z/W)$ degenerates at E_1 , the Chern class map yields an inclusion $\bar{c}_1: \mathrm{NS}(Z)/p\mathrm{NS}(Z) \hookrightarrow H_{\mathrm{cris}}^2(Z/k) \cong H_{\mathrm{dR}}^2(Z/k)$; see [DI81, Remark 3.5].

Let $\bar{c}_1(\alpha) \subset H_{\mathrm{dR}}^2(Z/k)$ be the span of $\bar{c}_1(\alpha(e_1)), \dots, \bar{c}_1(\alpha(e_r))$; it is actually a subspace of $\mathrm{Fil}^1 H_{\mathrm{dR}}^2(Z/k)$ (see, for example, [Mad15, §3.4]).

Since Δ_L is invertible in k , the line bundles $\mathcal{L}_1, \dots, \mathcal{L}_r$ are linearly independent in the quotient $\mathrm{NS}(Z)/p\mathrm{NS}(Z)$, and thus $\dim \bar{c}_1(\alpha) = r$. Now use the fact (see, for example, [Mad15, Theorem 3.8(3)], modeled after [Ogu84, Theorem 5.1(2)]) that a line bundle $\mathcal{L}$ extends to a given deformation $\tilde{Z}/A$ if and only if $c_1(\mathcal{L})$ is orthogonal to the corresponding lift $\mathrm{Fil}^2 H_{\mathrm{cris}}^2(\tilde{Z})(A)$. Under the isomorphism of Lemma 3.7, we see that the tangent space $T_s \mathrm{Def}(Z, \alpha)$ corresponds to homomorphisms from $\mathrm{Fil}^2 H_{\mathrm{dR}}^2(Z/k)$ into the orthogonal complement of $\bar{c}_1(\mathcal{L}_1), \dots, \bar{c}_1(\mathcal{L}_r)$ in $\mathrm{Gr}^1 H_{\mathrm{dR}}^2(Z/k)$. Because these Chern classes are linearly independent over k and $(\cdot, \cdot)$ is nondegenerate, the codimension of $T_s \mathrm{Def}(Z, \alpha)$ in $T_s \mathrm{Def}(Z)$ is r . $\square$

We now suppose that data $\underline{\chi} = (\mu_n, \chi^\omega, \chi)$ is chosen such that $R_{L, \underline{\chi}}$ is nonempty and further assume that n is invertible in k .

LEMMA 3.9. *Suppose $(Z, \alpha, \rho) \in R_{L, \underline{\chi}}(k)$ and $\mathrm{char}(k) \nmid 2\Delta_L n$. The (equicharacteristic) tangent space to $R_{L, \underline{\chi}}$ at (Z, α, ρ) has dimension $m(\chi^\omega) - 1$.*

Proof. By the crystalline local Torelli theorem ([BO83, Remark 3.23] and [Jan17, Lemma 3.1]; see also [NO85, Theorem 5.3 and Remark 5.3.1] and [Ogu84, Theorem 5.1(2)] for characteristic at least 5), it suffices to identify the sublocus of $\mathrm{Def}(Z, \alpha)$ to which the G -action on $H_{\mathrm{cris}}^2(Z)$ extends. Thus, let $\bar{c}_1(\alpha)^\perp$ be the orthogonal complement of $\bar{c}_1(\alpha)$, and consider the inclusions of formal deformation spaces $\mathrm{Def}(Z, \alpha, \rho) \subset \mathrm{Def}(Z, \alpha) \subset \mathrm{Def}(Z)$. Computing equicharacteristic tangent spaces at the base point s , we have

$$\begin{array}{ccc} T_s \mathrm{Def}(Z) & \xleftarrow{\sim} & \mathrm{Hom}(\mathrm{Fil}^2 H_{\mathrm{dR}}^2(Z/k), \mathrm{Gr}^1 H_{\mathrm{dR}}^2(Z/k)) \\ \uparrow & & \uparrow \\ T_s \mathrm{Def}(Z, \alpha) & \xleftarrow{\sim} & \mathrm{Hom}(\mathrm{Fil}^2 H_{\mathrm{dR}}^2(Z/k), c_1(\alpha)^\perp / \mathrm{Fil}^2 H_{\mathrm{dR}}^2(Z/k)) \\ \uparrow & & \uparrow \\ T_s \mathrm{Def}(Z, \alpha, \rho) & \xleftarrow{\sim} & \mathrm{Hom}_G(\mathrm{Fil}^2 H_{\mathrm{dR}}^2(Z/k), c_1(\alpha)^\perp / \mathrm{Fil}^2 H_{\mathrm{dR}}^2(Z/k)). \end{array}$$

By the definition of $R_{L, \underline{\chi}}$, the χ -eigenspace of $\mathrm{Fil}^1 H_{\mathrm{dR}}^2(Z/k)$ is fully contained in $c_1(\alpha)^\perp$; the result now follows. $\square$

4. Period maps for complex K3 surfaces

4.1 Period maps

The global complex Torelli theorem for K3 surfaces asserts that the isomorphism class of a K3 surface $Z/\mathbb{C}$ is determined by the isomorphism class of $H^2(Z, \mathbb{Z})$ as a polarized Hodge structure. Via Hodge theory, one thus obtains a good global understanding of the moduli of complex K3 surfaces, as follows.

Let Z be a marked K3 surface, that is, a K3 surface $Z/\mathbb{C}$ equipped with an isometry

$\phi: H^2(Z, \mathbb{Z}) \xrightarrow{\sim} L_{K3}$. The image of $\phi_{\mathbb{C}}(H^{2,0}(Z))$ determines an element of the period domain

$$\mathbb{X}_{L_{K3}} = \{[\sigma] \in \mathbb{P}(L_{K3} \otimes \mathbb{C}) : (\sigma, \sigma) = 0, (\sigma, \bar{\sigma}) > 0\},$$

and the isomorphism class of Z is determined by the class of $\phi_{\mathbb{C}}(H^{2,0}(Z))$ in $O(L_{K3}) \backslash \mathbb{X}_{L_{K3}}$. (We recall a useful description of $\mathbb{X}_{L_{K3}}$ below in Section 4.2.) We remind the reader that the action of the orthogonal group $O(L_{K3})$ on $\mathbb{X}_{L_{K3}}$ is not properly discontinuous, and thus the quotient space $O(L_{K3}) \backslash \mathbb{X}_{L_{K3}}$ is not even Hausdorff.

Now suppose that Z is equipped with a polarization λ of degree $2d$. Recall that we have fixed an embedding $\iota: \langle 2d \rangle \hookrightarrow L_{K3}$. A marking ϕ of Z induces an identification of the primitive cohomology $P_{\lambda}^2(Z, \mathbb{Z})$ with $\langle 2d \rangle^{\perp} \subset L_{K3}$, and thus $\phi_{\mathbb{C}}(H^{2,0}(Z))$ lies in

$$\mathbb{X}^{(2d)} := \mathbb{X}_{\langle 2d \rangle^{\perp}} = \{[\sigma] \in \mathbb{P}(\langle 2d \rangle^{\perp} \otimes \mathbb{C}) : (\sigma, \sigma) = 0, (\sigma, \bar{\sigma}) > 0\} \subset \mathbb{X}_{L_{K3}}.$$

Recall (Section 2) that $\tilde{O}^{(2d)}(\mathbb{Z})$ consists of those orthogonal automorphisms of $\langle 2d \rangle^{\perp}$ which admit an extension to L_{K3} fixing $\langle 2d \rangle$. We thus have a natural inclusion

$$\tilde{O}^{(2d)}(\mathbb{Z}) \backslash \mathbb{X}^{(2d)} \hookrightarrow O_{L_{K3}}(\mathbb{Z}) \backslash \mathbb{X}_{L_{K3}}.$$

The strong Torelli theorem for polarized K3 surfaces implies that there is an open immersion

$$\mathbb{R}_{2d, \mathbb{C}} \xrightarrow{\tau_{2d, \mathbb{C}}} \tilde{O}^{(2d)}(\mathbb{Z}) \backslash \mathbb{X}^{(2d)},$$

see for example [Huy16, Theorem 6.3.4], which extends to an isomorphism of coarse moduli spaces $\mathbb{R}_{2d, \mathbb{C}} \rightarrow \tilde{O}^{(2d)}(\mathbb{Z}) \backslash \mathbb{X}^{(2d)}$; see [Huy16, Remark 6.4.5].

More generally, for a primitive sublattice $L \subset L_{K3}$ of signature $(1, r-1)$, we set

$$\mathbb{X}^L = \mathbb{X}_{L^{\perp}} = \{[\sigma] \in \mathbb{P}(L^{\perp} \otimes \mathbb{C}) : (\sigma, \sigma) = 0, (\sigma, \bar{\sigma}) > 0\}$$

and obtain an open immersion

$$\mathbb{R}_{L, \mathbb{C}} \xrightarrow{\tau_{L, \mathbb{C}}} \tilde{O}^L(\mathbb{Z}) \backslash \mathbb{X}^L;$$

see [Dol96, § 3] and [DK07, § 11] for more details.

Finally, fix data $\underline{\chi} = (\mu_n, \chi^{\omega}, \chi)$, and suppose $(Z, \alpha, \rho) \in \mathbb{R}_{L, \underline{\chi}}(\mathbb{C})$. A choice of marking ϕ on Z induces an action of μ_n on L_{K3} with character χ . The period point $\phi_{\mathbb{C}}(H^{2,0}(Z))$ then lies in

$$\mathbb{X}^{L, \underline{\chi}} = \{[\sigma] \in \mathbb{P}((L_{\mathbb{C}}^{\perp})(\chi^{\omega})) : (\sigma, \sigma) = 0, (\sigma, \bar{\sigma}) > 0\},$$

where we single out an eigenspace for the action of μ_n on $L^{\perp}$ by

$$L_{\mathbb{C}}^{\perp}(\chi^{\omega}) = \{v \in L^{\perp} \otimes \mathbb{C} : \forall \zeta \in \mu_n(\mathbb{C}), \zeta v = \chi^{\omega}(\zeta)v\}.$$

Let $O^{L, \underline{\chi}}$ be the group of automorphisms of $L^{\perp}$ which commute with the action of μ_n ; if R is a ring over $\mathbb{Z}[\zeta_n]$, then

$$O^{L, \underline{\chi}}(R) = \{g \in O^L(R) : \forall \zeta \in \mu_n(R), g\zeta v = \zeta gv\}.$$

Let $\tilde{O}^{L, \underline{\chi}}$ be the subgroup of admissible automorphisms of $L^{\perp}$. Then we again have an open immersion

$$\mathbb{R}_{L, \underline{\chi}, \mathbb{C}} \hookrightarrow \tilde{O}^{L, \underline{\chi}}(\mathbb{Z}) \backslash \mathbb{X}^{L, \underline{\chi}}.$$

Recall that $m_{\underline{\chi}}(\chi^{\omega}) = m(\chi^{\omega}) = \dim L_{\mathbb{C}}^{\perp}(\chi^{\omega})$ is the multiplicity of the faithful character χ^{ω} in the representation χ and that $L \subset L_{K3}$ is the module of μ_n -invariants.

If $n \geq 3$, then χ^ω is imaginary, and

$$\mathbb{X}^{L, \underline{\chi}} \cong \mathbb{B}^{m_{\underline{\chi}}(\chi^\omega) - 1},$$

the complex unit ball of dimension $m_{\underline{\chi}}(\chi^\omega) - 1$.

If $n = 2$, then χ^ω is real, $L_{\mathbb{C}}^\perp(\chi^\omega) = L_{\mathbb{R}}^\perp(\chi^\omega) \otimes \mathbb{C}$, and $\mathbb{X}^{L, \underline{\chi}}$ is a type IV Hermitian symmetric space of dimension $m_{\underline{\chi}}(\chi^\omega) - 1$.

4.2 Period spaces

Since L has signature $(1, r - 1)$, the orthogonal complement $L^\perp$ has signature $(2, 19 - (r - 1))$. It is traditional in the K3 literature to describe the relevant period space as

$$\mathbb{X}^L = \mathbb{X}_{L^\perp} \cong \frac{\mathrm{O}^L(\mathbb{R})}{\mathrm{SO}_2(\mathbb{R}) \times \mathrm{O}_{20-r}(\mathbb{R})}.$$

To facilitate comparison with the Shimura variety literature, we prefer to recall that the special orthogonal group $\mathrm{SO}^L(\mathbb{R})$ already acts transitively on $\mathbb{X}^L$, and we in fact have

$$\mathbb{X}^L \cong \frac{\mathrm{SO}^L(\mathbb{R})}{\mathrm{SO}_2(\mathbb{R}) \times \mathrm{SO}_{20-r}(\mathbb{R})}.$$

It is perhaps worth noting that the special orthogonal group of a definite form is connected, while $\mathrm{SO}_{2,20-r}(\mathbb{R})$ has two topological components, indexed by the two classes of the spinor norm. In particular, $\mathbb{X}^L$ consists of two connected components, say $\mathbb{X}^{L+}$ and $\mathbb{X}^{L-}$; these components are stabilized by the component $\mathrm{SO}^L(\mathbb{R})^+ \subset \mathrm{SO}^L(\mathbb{R})$ of elements with trivial spinor norm.

Let $\Gamma \subset \mathrm{O}^L(\mathbb{R})$ be any arithmetic group. Then Γ has finite covolume and, in particular, meets every topological component of $\mathrm{O}^L(\mathbb{R})$. We have isomorphisms of complex-analytic spaces

$$\Gamma \backslash \mathbb{X}^L \cong (\Gamma \cap \mathrm{O}^L(\mathbb{R})^+) \backslash \mathbb{X}^{L+} \cong (\Gamma \cap \mathrm{SO}^L(\mathbb{R})) \backslash \mathbb{X}^L.$$

In particular, the period map is an open immersion

$$\mathrm{R}_{L, \mathbb{C}} \xrightarrow{\tau_L} \widetilde{\mathrm{SO}}^L(\mathbb{Z}) \backslash \mathbb{X}^L.$$

5. Shimura varieties

5.1 Integral canonical models

We review some basic concepts concerning Shimura varieties, referring the reader to [Del71] for foundational material, [Del79] for canonical models, [Kis10] for integral canonical models, and [Tae17] for stack-theoretic issues. All Shimura data are assumed to be of abelian type, so that the cited references suffice.

Let $(G, \mathbb{X})$ be a Shimura datum, consisting of a reductive group $G/\mathbb{Q}$ and a conjugacy class $\mathbb{X}$ of homomorphisms $\mathbf{R}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m \rightarrow G_{\mathbb{R}}$ of $\mathbb{R}$ -groups, subject to the usual axioms. Further assume that $(G, \mathbb{X})$ is of abelian type.

Let $\mathbb{K} \subset G(\mathbb{A}_f)$ be a neat compact open subgroup of the finite adelic points. The holomorphic analytic quotient stack $\mathrm{Sh}_{\mathbb{K}}^{\mathrm{an}}[G, \mathbb{X}] := [G(\mathbb{Q}) \backslash (\mathbb{X} \times G(\mathbb{A}_f) / \mathbb{K})]$ is represented by the analytification of a smooth complex quasiprojective variety $\mathrm{Sh}_{\mathbb{K}}(G, \mathbb{X})$. The variety $\mathrm{Sh}_{\mathbb{K}}(G, \mathbb{X})$ and the stack $\mathrm{Sh}_{\mathbb{K}}[G, \mathbb{X}]$ both descend to the reflex field $E(G, \mathbb{X})$.

More generally, let $\mathbb{K} \subset G(\mathbb{A}_f)$ be an open compact subgroup, and let $\mathbb{K}_0 \subset \mathbb{K}$ be a neat subgroup of finite index. Define the corresponding Shimura stack by $\mathrm{Sh}_{\mathbb{K}} = [\mathrm{Sh}_{\mathbb{K}_0}[G, \mathbb{X}]/(\mathbb{K}/\mathbb{K}_0)$; it is independent of the choice of $\mathbb{K}_0$.

Fix a prime p , let v be a prime of $E(G, \mathbb{X})$ lying over p , and let $\mathbb{K}_p \subset G(\mathbb{Q}_p)$ be hyperspecial. Then the pro-variety

$$\mathrm{Sh}_{\mathbb{K}_p} := \varprojlim_{\mathbb{K}^p \subset G(\mathbb{A}_f^p)} \mathrm{Sh}_{\mathbb{K}_p \mathbb{K}^p}(G, \mathbb{X})$$

admits an extension to $\mathcal{O}_{E(G, \mathbb{X}), v}$ as a pro-scheme with continuous $G(\mathbb{A}_f^p)$ -action, which we continue to denote by $\mathrm{Sh}_{\mathbb{K}_p}$. What makes this model the *integral canonical model* is the following extension property: if T is any regular formally smooth (pro-)scheme over $\mathcal{O}_{E(G, \mathbb{X}), v}$, then any morphism $T_{E(G, \mathbb{X})} \rightarrow \mathrm{Sh}_{\mathbb{K}_p}$ extends to all of T ; see, for example, [Kis10, § 2.3.7].

Consequently, for any $\mathbb{K} \subset G(\mathbb{A}_f)$ hyperspecial at p , the stack $\mathrm{Sh}_{\mathbb{K}}[G, \mathbb{X}]$ extends canonically to a smooth Deligne–Mumford stack over $\mathcal{O}_{E(G, \mathbb{X}), v}$. (If necessary, one can start with the canonical integral model of $\mathrm{Sh}_{\mathbb{K}_0}[G, \mathbb{X}]$ for some neat compact open subgroup $\mathbb{K}_0 \subset \mathbb{K}$ and then pass to the quotient by the action of $\mathbb{K}/\mathbb{K}_0$.)

In fact, let $\mathbb{K} \subset G(\mathbb{A}_f)$ be an open compact subgroup, and let $M = M(\mathbb{K})$ be the (finite) product of all primes p such that the component $\mathbb{K}_p$ is *not* hyperspecial. Using [Lov17, Theorem 2.2.1], we see that $\mathrm{Sh}_{\mathbb{K}}[G, \mathbb{X}]$ admits a canonical integral model over $\mathcal{O}_{E(G, \mathbb{X})}[1/M]$.

A morphism $f: (G_1, \mathbb{X}_1) \rightarrow (G_2, \mathbb{X}_2)$ of Shimura data is a morphism $G_1 \rightarrow G_2$ of algebraic groups which induces a morphism $\mathbb{X}_1 \rightarrow \mathbb{X}_2$. For future use, we collect some standard functorialities for morphisms of Shimura varieties.

LEMMA 5.1. *Let $f: (G_1, \mathbb{X}_1) \rightarrow (G_2, \mathbb{X}_2)$ be a morphism of Shimura data. Let $\mathbb{K}_1 \subset G_1(\mathbb{A}_f)$ and $\mathbb{K}_2 \subset G_2(\mathbb{A}_f)$ be compact open subgroups such that $f(\mathbb{K}_1) \subseteq f(\mathbb{K}_2)$.*

- (a) *Then f induces a morphism $f_{\mathbb{K}_1, \mathbb{K}_2}: \mathrm{Sh}_{\mathbb{K}_1}[G_1, \mathbb{X}_1] \rightarrow \mathrm{Sh}_{\mathbb{K}_2}[G_2, \mathbb{X}_2]$ of Shimura stacks over E .*
- (b) *If $\mathbb{K}_1$ and $\mathbb{K}_2$ are hyperspecial at all $p \nmid M$, then $f_{\mathbb{K}_1, \mathbb{K}_2}$ extends to a morphism of Shimura stacks over $\mathcal{O}_E[1/M]$.*
- (c) *If $f: G_1 \rightarrow G_2$ is injective, then $f_{\mathbb{K}_1, \mathbb{K}_2}$ is a closed morphism of Shimura stacks. If $\mathbb{K}_2$ is a sufficiently small compact open subgroup of $G_2(\mathbb{A}_f)$ which contains $\mathbb{K}_1$, then the generic fiber of $f_{\mathbb{K}_1, \mathbb{K}_2}$ is a closed embedding.*

Proof. Part (a) and (the generic fiber of) part (c) are due to Deligne [Del71, Proposition 1.15]; see also [Mil05, Theorem 5.16 and Remark 13.8]. The extension to integral models follows from the extension property and the smoothness of the integral model of $\varprojlim_{\mathbb{K}_1} \mathrm{Sh}_{\mathbb{K}_1}[G_1, \mathbb{X}_1]$. $\square$

5.2 Orthogonal Shimura varieties

Fix a nondegenerate lattice L of signature $(2, n_-)$, and let $G_L = \mathrm{SO}_{L \otimes \mathbb{Q}}$ be the associated special orthogonal group. Let $\mathbb{X}_L$ be the corresponding Hermitian symmetric space (Section 4.2).

Inside $G_L(\mathbb{A}_f)$, we single out the admissible integral automorphisms:

$$\mathbb{K}_L := \ker G_L(\widehat{\mathbb{Z}}) \rightarrow \mathrm{Aut}(\mathrm{disc}(L))(\widehat{\mathbb{Z}}).$$

The local component $\mathbb{K}_{L, p}$ at p is hyperspecial if $p \nmid \Delta_L$. Consequently, we have an integral canonical model

$$\mathrm{Sh}_L := \mathrm{Sh}_{\mathbb{K}_L}[G_L, \mathbb{X}_L]$$

over $\mathbb{Z}[1/2\Delta_L]$. (By inverting 2, we sidestep the intricacies of orthogonal groups and Shimura varieties in even characteristic.)

Note that $\mathbb{K} \cap \mathrm{SO}_L(\mathbb{R})^+ = \widetilde{\mathrm{SO}}_L(\mathbb{Z})$, and thus [Del79, 2.1.2]

$$\mathrm{Sh}_{L,\mathbb{C}} \cong \mathrm{SO}_L(\mathbb{Z})^+ \backslash \mathbb{X}_L^+ \cong \widetilde{\mathrm{SO}}_L(\mathbb{Z}) \backslash \mathbb{X}_L. \quad (5.2.1)$$

If $\mathbb{K} \subset \mathbb{K}_L \subset G_L(\mathbb{A}_f)$ is any compact open subgroup, then there is a surjection $\mathrm{Sh}_{\mathbb{K}}[G_L, \mathbb{X}_L] \rightarrow \mathrm{Sh}_L$ of stacks over $\mathbb{Z}[1/2\Delta_L N(\mathbb{K})]$. In particular, let $\mathbb{K}_{L,N} = \ker(\mathbb{K}_L \rightarrow G_L(\mathbb{Z}/N))$, and let $\mathrm{Sh}_{L,N} = \mathrm{Sh}_{\mathbb{K}_{L,N}}[G_L, \mathbb{X}_L]$, a stack over $\mathbb{Z}[1/2\Delta_L N]$. There is a surjection $\mathrm{Sh}_{L,N} \rightarrow \mathrm{Sh}_L$ with each geometric fiber a torsor under $\mathbb{K}_L/\mathbb{K}_{L,N} \cong G_L(\mathbb{Z}/N)$.

Now fix a primitive embedding of lattices $L_1 \hookrightarrow L_2$ with respective signatures $(2, n_{1,-})$ and $(2, n_{2,-})$.

LEMMA 5.2. *There is a closed morphism*

$$\mathrm{Sh}_{L_1} \xrightarrow{\psi_{L_1, L_2}} \mathrm{Sh}_{L_2}$$

of Shimura stacks over $\mathbb{Z}[1/(2\Delta_{L_1}\Delta_{L_2})]$ whose generic fiber is a closed embedding.

Proof. The chosen embedding gives an inclusion $G_{L_1} \rightarrow G_{L_2}$ of groups over $\mathbb{Q}$, which induces $\mathbb{X}_{L_1} \hookrightarrow \mathbb{X}_{L_2}$. Because of the admissibility condition, we have an inclusion $\mathbb{K}_{L_1} \hookrightarrow \mathbb{K}_{L_2}$, whence (Lemma 5.1) a morphism $\psi_{L_1, L_2}: \mathrm{Sh}_{L_1} \rightarrow \mathrm{Sh}_{L_2}$ over $\mathbb{Z}[1/(2\Delta_{L_1}\Delta_{L_2})]$. To verify that $\psi_{L_1, L_2, \mathbb{Q}}$ is a closed embedding, it suffices to check that $\psi_{L_1, L_2, \mathbb{C}}$ is an inclusion. This last claim follows from the description (5.2.1) and the fact that $\widetilde{\mathrm{SO}}_{L_1}(\mathbb{Z})$ consists of those orthogonal transformations of determinant 1 which lift to automorphisms of L_2 . $\square$

Similarly, for each positive integer N , there is a closed morphism $\mathrm{Sh}_{L_1, N} \rightarrow \mathrm{Sh}_{L_2, N}$ whose generic fiber is a closed embedding.

5.3 Unitary Shimura varieties

Let K be a quadratic imaginary field. Let L be a free $\mathcal{O}_K$ -module of rank r , equipped with a nondegenerate Hermitian form $h(\cdot, \cdot)$ of signature $(1, r-1)$. Attached to this is a Shimura datum $(G_{\mathcal{O}_K, L}, \mathbb{X}_{\mathcal{O}_K, L})$, where $G_{\mathcal{O}_K, L} = \mathrm{U}(L, h)$ is the group of $\mathcal{O}_K$ -linear automorphisms of L which preserve h and $\mathbb{X}_{\mathcal{O}_K, L} \cong \mathbb{B}^{r-1}$, the unit complex ball of dimension $r-1$. Let $\mathbb{K}_{\mathcal{O}_K, L}$ be the stabilizer in $G_{\mathcal{O}_K, L}(\mathbb{A}_f)$ of L . Let $\mathrm{Sh}_{\mathcal{O}_K, L} = \mathrm{Sh}_{\mathbb{K}_{\mathcal{O}_K, L}}[G_{\mathcal{O}_K, L}, \mathbb{X}_{\mathcal{O}_K, L}]$; it is the moduli space of abelian varieties of dimension r equipped with an action by $\mathcal{O}_K$ of signature $(1, r-1)$ and a polarization λ with $\ker(\lambda) \cong \mathrm{disc}(L)$. (More precisely, the relevant Shimura datum is $(\mathrm{U}(L \otimes \mathbb{Q}, h), \mathbb{X}_{(L \otimes \mathbb{Q}, h)})$; the choice of the lattice L inside the $\mathbb{Q}$ -vector space $L \otimes \mathbb{Q}$ defines the integral structure on $G_{\mathcal{O}_K, L}(\mathbb{A}^f)$.)

More generally, suppose that K is a CM field, with maximal totally real subfield K^+ , and again let L be a free $\mathcal{O}_K$ -module of rank r , equipped with a nondegenerate Hermitian form h . The archimedean signature of (L, h) is determined by data

$$\{(m_\sigma, n_\sigma)\}_{\sigma: K^+ \hookrightarrow \mathbb{R}}. \quad (5.3.1)$$

Let $G_{\mathcal{O}_K, L} = \mathrm{U}(L, h)$; the associated Hermitian symmetric domain $\mathbb{X}_{\mathcal{O}_K, L}$ has dimension $\sum m_\sigma n_\sigma$. If there exists some σ_0 such that $(m_{\sigma_0}, n_{\sigma_0}) = (1, r-1)$, and if $m_\sigma n_\sigma = 0$ for $\sigma \neq \sigma_0$, then we again have $\mathbb{X}_{\mathcal{O}_K, L} \cong \mathbb{B}^{r-1}$. Again, let $\mathrm{Sh}_{\mathcal{O}_K, L} = \mathrm{Sh}_{\mathbb{K}_{\mathcal{O}_K, L}}[G_{\mathcal{O}_K, L}, \mathbb{X}_{\mathcal{O}_K, L}]$.

In the applications pursued here, it turns out that either L is unimodular, or there is some prime p which is totally ramified in $\mathcal{O}_K$; the ring $\mathcal{O}_K$ acts on $\mathrm{disc}(L)$ through its quotient $\mathbb{F}_p$,

and $\text{disc}(L)$, as an abelian group, is isomorphic to $(\mathbb{Z}/p)^{2[(r-1)/2]}$. This shows up in the analysis by Kudla and Rapoport [KR12] of occult period maps. The only impact on the present study is that it shapes the structure of the polarization in the moduli-theoretic interpretation of $\text{Sh}_{\mathcal{O}_K, L}$.

In any event, the Shimura stack $\text{Sh}_{\mathcal{O}_K, L}$ admits a smooth integral model over $\mathcal{O}_K[1/\Delta(K)\Delta_L]$.

5.4 Shimura varieties and K3 surfaces

Let $L \hookrightarrow L_{K3}$ be a primitive sublattice of signature $(1, r-1)$. Consistent with earlier notation, we set

$$\text{Sh}^L = \text{Sh}_{\mathbb{K}^L} [G^L, \mathbb{X}^L] = \text{Sh}_{\mathbb{K}_{L^\perp}} [G_{L^\perp}, \mathbb{X}_{L^\perp}].$$

Now let $\underline{\chi} = (\mu_n, \chi^\omega, \chi)$ determine an action of μ_n on $L^\perp$ as in Section 3.2. Let $E(\underline{\chi}) = \mathbb{Q}(\zeta_n)$.

- If $n \geq 3$, let $\text{Sh}^{(L, \underline{\chi})} = \text{Sh}_{\mathcal{O}_{E(\underline{\chi})}, L^\perp}$; then $\text{Sh}^{(L, \underline{\chi})}(\mathbb{C})$ is an arithmetic quotient of a complex ball.
- If $n = 2$, let $\text{Sh}^{(L, \underline{\chi})} = \text{Sh}_{L^\perp}$; then $\text{Sh}^{(L, \underline{\chi})}(\mathbb{C})$ is an arithmetic quotient of a Hermitian symmetric space of type IV.

LEMMA 5.3. *The periods of a structured K3 surface determine holomorphic open immersions*

$$\begin{aligned} R_{L, \mathbb{C}} &\xrightarrow{\tau_{L, \mathbb{C}}} \text{Sh}_{\mathbb{C}}^L, \\ R_{(L, \underline{\chi}), \mathbb{C}} &\xrightarrow{\tau_{(L, \underline{\chi}), \mathbb{C}}} \text{Sh}^{(L, \underline{\chi})}. \end{aligned}$$

Proof. The period domain of a family of structured K3 surfaces is computed in, for example, [Dol96, DK07]. The interpretation in terms of Shimura varieties is standard and is drawn out (in some cases) in, for instance, [Kud13, Mad15, Riz06]. $\square$

Remark 5.4. In the case of a datum $(L, \underline{\chi})$, the complement of the image of $\tau_{(L, \underline{\chi})}$, when known, is often a ball quotient in its own right; see, for example, [Kon02] for a representative example. Kudla and Rapoport, in several cases, interpret this complement as a “special cycle.” In particular, this complement is itself a Shimura variety. The author conjectures that this structure of the complement holds integrally, as well. In the special case of cubic surfaces, this is worked out in [Ach14]; for now, it seems that the general case remains open.

6. Integral period maps

With the notation established above, the Torelli theorem for complex K3 surfaces asserts that there is an open immersion

$$R_{2d, \mathbb{C}} \xhookrightarrow{\tau_{2d, \mathbb{C}}} \text{Sh}_{\mathbb{C}}^{\langle 2d \rangle}$$

of stacks over $\mathbb{C}$. In fact, it is known that this map preserves arithmetic.

PROPOSITION 6.1. *The period map $\tau_{2d, \mathbb{C}}$ descends to a morphism $\tau_{2d}: R_{2d} \rightarrow \text{Sh}^{\langle 2d \rangle}$ of stacks over $\mathbb{Q}$.*

Proof. Rizov has proved this for R_{2d}° , using an analogue of CM theory for K3 surfaces; see [Riz10, Theorem, p. 14] and [Tae17, Theorem 5]. The statement for R_{2d} follows from descent relative to $\text{Spec } \mathbb{C} \rightarrow \text{Spec } \mathbb{Q}$ since R_{2d}° is dense in R_{2d} . $\square$

Using Proposition 6.1 as a starting point, we will show that other period maps also descend to a natural field of definition and extend integrally. We start with an interlude on level structures, so that we can work with quasiprojective schemes and verify descent in an elementary fashion.

6.1 Level structures

It is possible to define the notion of a lattice-polarized K3 surface with $\mathbb{K}$ level structure for an essentially arbitrary open subgroup of $G^L(\mathbb{Z})$; but we will content ourselves here with a more limited notion which is adequate for our purposes. (See [Tae17, § 5.2] and [Riz10] for more details in the case $L = \langle 2d \rangle$.)

Fix an integer $N > 2$ which is relatively prime to $2p\Delta_L$. Then $\mathrm{SO}^L(\mathbb{Z}/N)$ is admissible; any automorphism of $L^\perp \otimes \mathbb{Z}/N$ lifts uniquely to $L_{K3} \otimes \mathbb{Z}/N$ as an element which fixes L .

If $(Z \rightarrow S, \alpha) \in \mathbf{R}_L(S)$, a full level N structure on $(Z \rightarrow S, \alpha)$ is an isomorphism of formed spaces $\beta: L_{K3} \otimes \mathbb{Z}/N \xrightarrow{\sim} R^2 f_* \mathbb{Z}/N(1)$ such that the following diagram commutes:

$$\begin{array}{ccc} L_{K3} \otimes \mathbb{Z}/N & \xrightarrow[\sim]{\beta} & R^2 f_* \mathbb{Z}/N(1) \\ \uparrow & & \uparrow c_{1,N} \\ L & \xrightarrow{\alpha} & \mathrm{Pic}_{Z/S}(S), \end{array}$$

where the right-hand vertical map is the Chern class and the left-hand map is induced by the fixed inclusion $L \hookrightarrow L_{K3}$.

Since $N > 2$, the stack $\mathbf{R}_{L,N}$ is representable by a smooth quasiprojective scheme over $\mathbb{Z}_{(p)}$ (see, for example, [Riz10, Corollary 2.4.3] for the case $L = \langle 2d \rangle$). Moreover, because of the admissibility condition, $\mathbf{R}_{L,N} \rightarrow \mathbf{R}_L$ is Galois, with covering group isomorphic to $\{g \in \mathrm{SO}_{L_{K3}}(\mathbb{Z}/N) : g|_{L^\perp \otimes \mathbb{Z}/N} = \mathrm{id}\}$.

As before, given L , choose a primitive embedding of lattices $\langle 2d \rangle \hookrightarrow L$. The forgetful maps yield a Cartesian diagram

$$\begin{array}{ccc} \mathbf{R}_{L,N} & \hookrightarrow & \mathbf{R}_{\langle 2d \rangle, N} \cong \mathbf{R}_{2d, N} \\ \downarrow & & \downarrow \\ \mathbf{R}_L & \hookrightarrow & \mathbf{R}_{\langle 2d \rangle}, \end{array}$$

where the horizontal arrows are closed immersions and the vertical arrows are quotients by suitable subgroups of $\mathrm{SO}_{L_{K3}}(\mathbb{Z}/N)$.

6.2 Descent to the reflex field

LEMMA 6.2. (a) *Let $L \hookrightarrow L_{K3}$ be a primitive lattice of signature $(1, r-1)$. Then the complex period map $\tau_{L, \mathbb{C}}$ descends to a morphism $\tau_L: \mathbf{R}_L \rightarrow \mathrm{Sh}^L$ of stacks over $\mathbb{Q}$.*

(b) *Let $(L, \underline{\chi})$ be as in Section 3.2. Then the complex period map $\tau_{(L, \underline{\chi}), \mathbb{C}}$ descends to a morphism $\tau_{(L, \underline{\chi})}$ of stacks over $E(\underline{\chi})$.*

Proof. We address part (a) in detail. Fix some $N > 2$. Since $\mathbf{R}_L = [\mathbf{R}_{L,N}/G^L(\mathbb{Z}/N)]$ and $\mathrm{Sh}^L = [\mathrm{Sh}_N^L/G^L(\mathbb{Z}/N)]$, it suffices to show that the complex period map with level N structure, $\tau_{L,N, \mathbb{C}}: \mathbf{R}_{L,N, \mathbb{C}} \rightarrow \mathrm{Sh}_{N, \mathbb{C}}^L$, descends to $\mathbb{Q}$. Choose a primitive embedding $\langle 2d \rangle \hookrightarrow L$. We have a commuting diagram of universally injective morphisms of complex reduced quasiprojective

varieties

$$\begin{array}{ccc}
 R_{L,N,\mathbb{C}} & \xrightarrow{\tau_{L,N,\mathbb{C}}} & \mathrm{Sh}_{N,\mathbb{C}}^L \\
 \downarrow \phi_{L,2d,\mathbb{C}} & & \downarrow \psi_{G^L,G^{(2d)}} \\
 R_{2d,N,\mathbb{C}} & \xrightarrow{\quad} & \mathrm{Sh}_{N,\mathbb{C}}^{(2d)}.
 \end{array} \tag{6.2.1}$$

Since $R_{L,N,\mathbb{C}} \rightarrow R_{2d,N,\mathbb{C}}$ and $R_{2d,N,\mathbb{C}} \rightarrow \mathrm{Sh}_{N,\mathbb{C}}^{(2d)}$ descend to $\mathbb{Q}$, so does $\psi_{G^L,G^{(2d)}} \circ \tau_{L,N,\mathbb{C}}$. Since $\psi_{G^L,G^{(2d)}}$ is universally injective (Lemma 5.2), the morphism $\tau_{L,N,\mathbb{C}}$ is $\mathrm{Aut}(\mathbb{C}/\mathbb{Q})$ -equivariant on $\mathbb{C}$ -points and thus descends to $\mathbb{Q}$ as well.

The proof of part (b) is exactly the same, except that the role of $G^L(\mathbb{Z}/N)$ is now played by the finite unitary group $G^{(L,\underline{\chi})}(\mathbb{Z}/N)$ and (6.2.1) is replaced by

$$\begin{array}{ccc}
 R_{(L,\underline{\chi}),N,\mathbb{C}} & \xrightarrow{\tau_{(L,\underline{\chi}),N,\mathbb{C}}} & \mathrm{Sh}_{N,\mathbb{C}}^{(L,\underline{\chi})} \\
 \downarrow \phi_{(L,\underline{\chi}),L,\mathbb{C}} & & \downarrow \psi_{G^{(L,\underline{\chi})},G^L} \\
 R_{L,N,\mathbb{C}} & \xrightarrow{\quad} & \mathrm{Sh}_{N,\mathbb{C}}^L.
 \end{array} \quad \square$$

6.3 Integral extension

Granting the existence of integral canonical models of Shimura varieties, it is not hard to see that τ_{2d} extends to a morphism of stacks over $\mathbb{Z}[1/6d]$; this is achieved in [Riz06, Theorem 4.3.3]. We refer to [Mad15, Corollary 5.15] for the difficult extension of this work to $\mathbb{Z}[1/2]$.

Remark 6.3. In fact, in [Mad15], the proof naturally gives rise to a period map for a trivial double cover of R_{2d} . Taelman has observed [Tae17] that by viewing the period map as measuring the primitive cohomology *twisted by the determinant*, the need for a double cover is eliminated.

Following Taelman's analysis [Tae17, §5], let $\mathbb{K}_L^* = \{\gamma \in \tilde{O}_L(\widehat{\mathbb{Z}}) : \det(\gamma) \in \{\pm 1\}\}$. Then $\mathbb{K}_{L^\perp}$ acts on $L^\perp$ via the determinant; Taelman defines, for instance, a period map

$$R_{2d}(\mathbb{C}) \rightarrow [\mathrm{SO}^{(2d)}(\mathbb{C}) \backslash (\mathbb{X}^{(2d)} \times \mathrm{SO}^{2d}(\mathbb{A}_f) / (\mathbb{K}^{(2d)})^*].$$

The target space is isomorphic, as an analytic space, to our $\mathrm{Sh}_{\mathbb{K}^{(2d)}}[G^{(2d)}, \mathbb{X}^{(2d)}]$. However, this target naturally identifies a polarized Hodge structure (H, s) with $(H, -s)$. In this way, the effect of a choice of generator of $\langle 2d \rangle$ is erased.

More generally, by following Taelman's formulation, we can suppress the choice of a “positive light cone” in the definition of R_L in Section 3.2; two L -polarizations which agree up to sign are identified by the action of $\mathbb{K}_L^*$ through its determinant.

We now secure analogous results for other period maps.

LEMMA 6.4. (a) *Let $L \hookrightarrow L_{K3}$ be a primitive lattice of signature $(1, r-1)$. Then the period map extends to a morphism $\tau_L : R_L \rightarrow \mathrm{Sh}^L$ of stacks over $\mathbb{Z}[1/2\Delta(L)]$.*

(b) *Let $(L, \underline{\chi})$ be as in Section 3.2. Then the period map extends to a morphism $\tau_{(L,\underline{\chi})} : R_{(L,\underline{\chi})} \rightarrow \mathrm{Sh}^{(L,\underline{\chi})}$ of stacks over $\mathcal{O}_{E(\underline{\chi})}[1/2\Delta(L)]$.*

Proof. Since Sh^L is separated, it suffices to show that for a fixed $p \nmid 2\Delta(L)$, the morphism τ_L extends to $\mathbb{Z}_{(p)}$. Let $N \geq 3$ be a natural number relatively prime to p . Since $R_{L,N}$ is smooth

(Proposition 3.8), the extension property of the integral canonical model implies that the morphism τ_L extends to $\mathbb{Z}_{(p)}$.

The proof of part (b) is the same, except that the necessary smoothness is secured in Lemma 3.9. $\square$

Remark 6.5. The generic fiber of the morphism $\mathrm{Sh}^L \hookrightarrow \mathrm{Sh}^{\langle 2d \rangle}$ is a closed immersion (Lemma 5.2). If it were known that $\psi_{L^\perp, \langle 2d \rangle^\perp}$ is a closed immersion of Shimura stacks, one could give an elementary proof of Lemma 6.4, as follows. Suppose $p \nmid \Delta(L)d(L)$; choose d with $p \nmid d$ such that there exists a primitive embedding $\langle 2d \rangle \hookrightarrow L$. We start with a diagram as in (6.2.1), where all objects are defined over $\mathbb{Z}_{(p)}$, except that τ_L is only known to be defined over $\mathbb{Q}$:

$$\begin{array}{ccc} \mathrm{R}_{L,N} & \xrightarrow{\tau_L} & \mathrm{Sh}_N^L \\ \downarrow \phi & & \downarrow \psi \\ \mathrm{R}_{2d,N} & \xrightarrow{\tau_{2d}} & \mathrm{Sh}_N^{\langle 2d \rangle} \end{array}$$

We know that ϕ is a closed immersion and τ_{2d} is an open immersion, and thus the composition $\mathrm{R}_{L,N} \rightarrow \mathrm{Sh}_N^{\langle 2d \rangle}$ is a locally closed immersion. All schemes involved are Noetherian and $\mathrm{R}_{L,N}$ is reduced, so the image of $\mathrm{R}_{L,N}$ is an open subscheme of a closed subscheme of $\mathrm{Sh}_N^{\langle 2d \rangle}$; see [Sta17, Tag 03DQ]. We are operating under the hypothesis that ψ is a closed immersion (Lemma 5.1). Since Sh_N^L is reduced, ψ maps Sh_N^L isomorphically onto its image, a closed subscheme of $\mathrm{Sh}_N^{\langle 2d \rangle}$.

We have observed (Section 5.4) that $\tau_{2d} \circ \phi$ maps the characteristic zero fiber $\mathrm{R}_{L,N,\mathbb{Q}}$ into $\psi(\mathrm{Sh}_{N,\mathbb{Q}}^L)$ inside $\mathrm{Sh}_N^{\langle 2d \rangle}$. Since $\psi(\mathrm{Sh}_N^L)$ is closed, $\mathrm{R}_{L,N,\mathbb{Q}}$ is dense in $\mathrm{R}_{L,N}$ (by flatness over $\mathbb{Z}_{(p)}$; see Proposition 3.3), and $\tau_{2d} \circ \phi(\mathrm{R}_{L,N})$ is locally closed, it follows that $\tau_{2d} \circ \phi(\mathrm{R}_{L,N}) \subseteq \psi(\mathrm{Sh}_N^L)$.

In particular, $\tau_{2d} \circ \phi$ factors through a locally closed immersion $\tau_L: \mathrm{R}_{L,N} \rightarrow \mathrm{Sh}_N^L$. We again invoke the fact that for Noetherian schemes, a locally closed immersion factors as an open immersion followed by a closed immersion. The fact that $\dim \mathrm{R}_{L,N} = \dim \mathrm{Sh}_N^L$ (and the reducedness of $\mathrm{R}_{L,N}$) now implies that, in any such factorization, the closed immersion must be the identity map, and therefore τ_{2d} is an open immersion.

7. From complete intersections to K3 surfaces

Thanks especially to works of Kondō, we know that sometimes one can associate a structured K3 surface with certain types of complete intersection varieties. Some of these constructions are reviewed here, with an eye toward making sense of these associations in families and ultimately explaining the arithmetic origin of Kondō's analytic ball-quotient maps.

In an attempt to minimize repetition in the statement of our main results, we make the following definition.

DEFINITION 7.1. We say that $(\mathrm{R}^\circ, \mathrm{N}, \mathrm{S}, \kappa, \tau)$ satisfies $(\dagger)$ over $\mathcal{O}$ if there is a diagram

$$\begin{array}{ccc} \mathrm{R} & \xrightarrow{\kappa} & \mathrm{N} \\ \downarrow \tau & & \\ \mathrm{S} & & \end{array} \quad (\dagger)$$

of stacks over $\mathcal{O}$, where κ induces an isomorphism on coarse moduli spaces and τ induces an open immersion $\mathrm{R}^\circ(\mathbb{C}) \hookrightarrow \mathrm{S}(\mathbb{C})$.

We should note that in many of the examples studied here (Sections 7.2, 7.3, and 7.7), Kudla and Rapoport have already shown that a transcendently defined occult period map descends to a natural cyclotomic field of definition [KR12, § 9]. Their method of proof goes back (at least) to Deligne [Del72, Theorem 2.12]. Roughly speaking, one shows that a monodromy representation is so large that a certain abelian scheme admits no automorphisms and thus descends. This strategy presumably also dispatches Section 7.6, perhaps with [DM86] providing the necessary monodromy calculation. The applications in Sections 7.4 and 7.5 do not literally fit within the framework of unitary Shimura varieties attached to quadratic imaginary fields, which may explain their omission from [KR12].

7.1 Stacks of varieties with group action

In Kondō's constructions, the original variety is encoded in the fixed locus of the group action on the K3 surface. If a group scheme G/S acts on a scheme Z/S , one can define Z^G , the fixed point stack [Rom05, Proposition 2.5].

LEMMA 7.2. *Suppose that $Z \rightarrow S$ is a K3 space and $G \subset \mathrm{Aut}_{Z/S}(S)$ is a nontrivial finite cyclic group.*

- (a) *The fixed locus $Z^G \rightarrow S$ is a scheme.*
- (b) *If $s \in S$, then Z_s^G is smooth and has at most one component of dimension 1 and genus at least 2.*
- (c) *If S is irreducible with generic point η and if $C_\eta \subset Z_\eta^G$ is a curve of genus at least 2, then the closure C of C_η in Z^G is a smooth proper relative curve over S .*

Proof. Since $Z \rightarrow S$ is an algebraic space, so is Z^G (see [Rom05, Remark 3.4(ii)]); since all components of all fibers are smooth (see below) of dimension at most 1, the fixed locus Z^G is actually a scheme.

The smoothness assertion of part (b) is proved in [AST11, Lemma 2.2] (in characteristic zero) and [Keu14, Proposition 1.4] (in positive characteristic). The fact that there is at most one curve of general type is also in [AST11, Lemma 2.2]. (While the statement is only claimed for complex K3 surfaces, the argument relies on nothing more than the Hodge index theorem.)

Part (c) follows from the upper semicontinuity, on Z^G , of the function $z \mapsto \dim(Z_{\varpi(z)}^G)$; see [GD64, IV, Theorem 13.1.3]. $\square$

We will occasionally have cause to work with the stacks of smooth relative uniform cyclic covers of projective spaces, as in [AV04]. Recall that if $X \rightarrow S$ is a smooth scheme, then a smooth relative uniform cyclic cover of degree n consists of a morphism $f: Y \rightarrow X$ which commutes with an action of μ_n on Y such that the branch divisor of f is smooth over S and, Zariski-locally on X , the scheme Y is μ_n -equivariantly isomorphic to $\mathcal{O}_Y(U)[y]/(y^n - h)$. With a slight adjustment of the notation of [AV04], let $\mathrm{H}(n, m, d)$ be the stack of smooth relative uniform cyclic covers $f: Y \rightarrow P \rightarrow S$ of degree n , where $P \rightarrow S$ is a Brauer–Severi scheme of dimension m and the branch divisor of f has degree d . Thus, for example, $\mathrm{H}(2, 1, 2g + 2)$ is the moduli stack of hyperelliptic curves of genus g . (A Brauer–Severi scheme $P \rightarrow S$ of dimension m is an S -scheme which, étale-locally on S , is isomorphic to the projective space of dimension m .)

In the special case where $m = 1$, let $\tilde{\mathrm{H}}(n, 1, d)$ be the stack of smooth relative uniform cyclic covers of Brauer–Severi curves equipped with a labeling of the branch locus; there is a forgetful map $\tilde{\mathrm{H}}(n, 1, d) \rightarrow \mathrm{H}(n, 1, d)$, with fiber a torsor under the symmetric group S_d on d letters. (Since

a Brauer–Severi scheme with a section is trivial, the underlying scheme of an object in $\tilde{H}(n, 1, d)$ is actually a family of projective lines, rather than merely étale-locally a family of projective lines.)

Let $\tilde{M}_{0,d}$ be the moduli space of d distinct labeled points in $\mathbb{P}^1$. By sending a labeled branched cover of the projective line to its branch locus, we obtain a morphism $\tilde{H}(n, 1, d) \rightarrow \tilde{M}_{0,d}$. In fact, this morphism is the rigidification along μ_n ; it factors as $\tilde{H}(n, 1, d) \rightarrow \tilde{H}(n, 1, d) // \mu_n \xrightarrow{\sim} \tilde{M}_{0,d}$ and, in particular, induces an isomorphism on coarse moduli spaces. This morphism is S_d -equivariant, and we have $H(n, 1, d) \rightarrow M_{0,d}$.

Below, we will often have a morphism $\alpha: S \rightarrow T$ of smooth stacks. Then each stack is normal and, in particular, has a normal coarse moduli space. If α induces a bijection on geometric points, then (by Zariski’s main theorem) α induces an isomorphism of coarse moduli spaces.

7.2 Curves of genus 4

Here we follow [Kon02]. The argument given here is also a prototype for the remainder of this section.

Let C/k be a smooth projective nonhyperelliptic curve of genus 4 with no vanishing theta constants, over an algebraically closed field in which 6 is invertible. Its canonical model is the (complete) intersection in $\mathbb{P}^3$ of quadric and cubic surfaces Q and S . Let $\varpi: Z \rightarrow Q$ be the triple cover of Q branched along C ; then Z comes equipped with an action by μ_3 . Let M_1 and M_2 be smooth lines on Q which represent the two rulings, and let $N_i = \varpi^{-1}M_i$. Then each N_i is an elliptic curve, and the two of them pair as $(N_1, N_2) = 3$. Moreover, N_1 and N_2 span a primitive lattice of $\text{Pic}(Z)$ isomorphic to $L_4 := U(3)$. Let $L_4 \hookrightarrow L_{K3}$ be a primitive embedding; the orthogonal complement of this copy of $U(3)$ is $L_4^\perp \cong U(3) \oplus U \oplus E_8(-1)^{\oplus 2}$; see [Kon02, § 2, p. 386]. In the notation of Section 2, we have $d(L_4) = 3$, and so there is a closed immersion $R_L \hookrightarrow R_{(6)}$ of smooth stacks over $\mathbb{Z}[1/6]$.

The action of μ_3 on Z is nonsymplectic, in the sense that χ^ω , the character of the representation of μ_3 by which μ_3 acts on $H^0(Z, \Omega^2)$, is faithful. Kondō explicitly writes down a certain representation ρ of μ_3 on $L_4^\perp$. (Of course, $L_4^\perp$ is free over $\mathbb{Z}[\zeta_3]$, in accordance with [MO98, Lemma 1.1].) Let χ_4 be the character of $\rho \oplus \rho_{\text{triv}}^{\oplus 2}$. In the case where $k = \mathbb{C}$, Kondō shows that Z is an element of $R_{L_4, \chi_4}(\mathbb{C})$. Let N_4 be the moduli space of nonhyperelliptic curves of genus 4. It is not hard to extend the work in [Kon02] to show the following.

LEMMA 7.3. *There is a morphism $\kappa_4: R_{L_4, \chi_4}^\circ \rightarrow N_4$ of stacks over $\mathbb{Z}[\zeta_3, 1/6]$ which is a bijection on geometric points.*

Proof. Suppose $(Z \rightarrow S, \alpha, \rho) \in R_{L_4, \chi_4}^\circ(S)$. In particular, $Z \rightarrow S$ is an algebraic space. Let $B = Z^{\mu_3} \rightarrow S$ be the scheme of fixed points (Lemma 7.2).

We will show that every fiber of $B \rightarrow S$ is a smooth projective nonhyperelliptic curve of genus 4. Then B is a scheme over S , and the sought-for functor $\kappa_4: R_{L_4, \chi_4}^\circ \rightarrow N_4$ is then given by $(Z \rightarrow S, \alpha, \rho) \mapsto (Z^{\mu_3} \rightarrow S)$.

So, let s be a point of S . If s has residue characteristic zero, the proof of [Kon02, Theorem 1] shows that B_s is a smooth projective nonhyperelliptic curve of genus 4.

If s has positive characteristic $p \geq 5$, since R_{L_4, χ_4}° is smooth over $\mathbb{Z}[\zeta_3, 1/6]$, the point s lifts to characteristic zero. More precisely, there exist a mixed-characteristic discrete valuation ring A , with general and special fibers η and $\circ$, and a point $P \in R_{L_4, \chi_4}^\circ(\text{Spec } A)$ with $P_\circ = s$. The

characteristic zero result for B_η , combined with the specialization argument of Lemma 7.2(c), shows that there is a (necessarily unique) smooth projective curve C_s of genus 4 in B_s .

Moreover, the quotient Z_s/μ_4 is a quadric (cone) [Kon02, §2, Proof of Claim, p. 389], and C_s maps isomorphically onto its image in the quotient. Insofar as C_s is a genus 4 curve lying on a quadric surface in $\mathbb{P}^3$, it is not hyperelliptic.

This defines the morphism $R_{L_4, \chi_4}^\circ \rightarrow N_4$. Now let k be an algebraically closed field in which 6 is invertible. The construction at the beginning of this subsection—modified to take a minimal resolution, if necessary, to account for the impact of vanishing theta characteristics—gives a set-theoretic section to $R_{L_4, \chi_4}^\circ(k) \rightarrow N_4(k)$. $\square$

We can finally explain the arithmetic origin of Kondō's observation that $N_4(\mathbb{C})$ is an arithmetic ball quotient.

PROPOSITION 7.4. *The tuple $(R_{L_4, \chi_4}^\circ, N_4, \text{Sh}^{(L_4, \chi_4)}, \kappa_4, \tau_{L_4, \chi_4})$ satisfies $(\dagger)$ over $\mathbb{Z}[\zeta_3, 1/6]$.*

Proof. This simply summarizes the foregoing. Consider κ_4 from Lemma 7.3. Since it yields a bijection on geometric points, it induces an isomorphism of coarse moduli spaces. For τ_{L_4, χ_4} , Kudla and Rapoport [KR12, Theorem 8.1] interpret Kondō's isomorphism [Kon02, Theorem 1] as a morphism $R_{L_4, \chi_4, \mathbb{C}}^\circ \rightarrow \text{Sh}_{\mathbb{C}}^{(L_4, \chi_4)}$ (see Section 5.4). Then Lemma 6.4 shows that this map descends and spreads to $\mathbb{Z}[\zeta_3, 1/6]$. $\square$

Remark 7.5. In characteristic zero, Kudla and Rapoport use a transcendental construction, and the fact that $\text{Sh}^{(L_4, \chi_4)}$ is a moduli space for abelian varieties with action by $\mathbb{Z}[\zeta_3]$, to interpret Kondō's construction as a morphism of stacks $N_{4, \mathbb{C}} \rightarrow \text{Sh}_{\mathbb{C}}^{(L_4, \chi_r)}$. They then use a monodromy argument [KR12, Proof of Lemma 9.1] to show that this map descends to a morphism of stacks over $\mathbb{Q}(\zeta_3)$, and conjecture that it extends to a morphism over $\mathbb{Z}[\zeta_3]$.

Remark 7.6. Let $R_{L_4, \chi_4}^\circ // \mu_3$ be the rigidification of R_{L_4, χ_4}° along μ_3 ([ACV03]; see also [Rom05, §5]). Then $R_{L_4, \chi_4}^\circ // \mu_3$ has the same coarse moduli space as R_{L_4, χ_4}° , and the morphism of Proposition 7.4 factors as

$$R_{L_4, \chi_4}^\circ \longrightarrow R_{L_4, \chi_4}^\circ // \mu_3 \longrightarrow N_4.$$

In this notation, Kudla and Rapoport conjecture [KR12, Remark 7.2] that the second map is an isomorphism of stacks. In fact, this has recently been resolved using transcendental means by Zheng [Zhe20, Proposition 7.9], who goes on to show that, at least over $\mathbb{C}$, there is an open immersion of orbifolds $N_{4, \mathbb{C}} \rightarrow \text{Sh}_{\mathbb{C}}^{(L_4, \chi_4)}$. Zheng also proves analogous statements in the situations of Sections 7.3 and 7.7 below.

7.3 Curves of genus 3

Kondō has given [Kon00] a similar characterization of $N_3(\mathbb{C})$, the set of complex nonhyperelliptic curves of genus 3. This construction also descends to arithmetic geometry, as follows.

Let C/k be a smooth projective nonhyperelliptic curve of genus 3 over an algebraically closed field in which 2 is invertible. Its canonical model is a smooth plane quartic curve; let $\varpi: Z \rightarrow \mathbb{P}^2$ be the cyclic quartic cover ramified along C . Then Z is a K3 surface, and inside $\text{Pic}(Z)$ is a lattice $L_3 \cong A_1 \oplus A_1(-1)^{\oplus 7}$; see [Kon00, Section 2, p. 222]. (Briefly, Z is a double cover of Y , itself a double cover of $\mathbb{P}^2$ branched along C . Then Y is a del Pezzo surface of degree 2 and thus may also be obtained as the blowup of a projective plane at seven points. The seven copies of

$A_1(-1)$ in $\text{Pic}(Z)$ are obtained from lifts to Z of the seven exceptional divisors; the remaining element of modulus 2 is the pullback of the class of a line on the projective plane.) Then L_3 embeds primitively into L_{K3} , with orthogonal complement $L_3^\perp \cong U(2)^{\oplus 2} \oplus D_8(-1) \oplus A_1(-1)^{\oplus 2}$ and $d(L_3) = 2$.

By construction, Z comes equipped with an action by μ_4 . Then μ_4 acts on the space of holomorphic 2-forms via a faithful character, χ^ω . The action ρ of μ_4 on $L_3^\perp$ is given explicitly in [Kon00, § 2], and we let χ_4 be the character of $\rho \oplus \rho_{\text{triv}}^{\oplus 8}$.

PROPOSITION 7.7. *There is a morphism $\kappa_3: R_{L_3, \chi_3}^\circ \rightarrow N_3$ such that $(R_{L_3, \chi_3}^\circ, N_3, \text{Sh}^{(L_3, \chi_3)}, \kappa_3, \tau_{L_3, \chi_3})$ satisfies $(\dagger)$ over $\mathbb{Z}[\sqrt{-1}, 1/2]$.*

Proof. As in Lemma 7.3, the map κ_3 is given by $(Z \rightarrow S, \alpha, \rho) \mapsto (Z^{\mu_4} \rightarrow S)$; see [Kon00, Section 2, p. 225] and the étale Lefschetz fixed point theorem [SGA5, III.(4.11.3)] provide the necessary geometric input to show that Z^{μ_4} is a relative nonhyperelliptic curve of genus 3. The fact that κ_3 gives a bijection on geometric points is established by the construction at the beginning of this section. The asserted behavior of τ_{L_3, χ_3} is a special case of Lemma 6.4. $\square$

Remark 7.8. See [KR12, § 7] for earlier results over $\mathbb{Q}(\sqrt{-1})$.

7.4 Curves of genus 6

Following [AK11], let N_6 denote the moduli stack (over $\mathbb{Z}[1/2]$) of non-special curves of genus 6. (Thus, a curve is represented by a point in N_6 if it is smooth, projective, and irreducible of genus 6, and neither hyperelliptic, trigonal, bielliptic, nor smooth quintic and planar.)

In fact, let C/k be a non-special curve over an algebraically closed field. The canonical embedding of C is a quadric section of a unique quintic del Pezzo surface Y in $\mathbb{P}^5$. Let Z be the double cover of Y branched along C . (If C has fewer than five distinct linear systems of type g_6^2 , then one must actually take the minimal resolution of this cover.) Then Z is a K3 surface with an action by $\mu_2 = \{\pm 1\}$, and this action fixes a lattice in $\text{Pic}(Z)$ isomorphic to $L_6 := A_1 \oplus A_1(-1)^4$; see [AK11, § 2.1]. (Note that $d(L_6) = 2$.)

PROPOSITION 7.9. *There is a morphism $\kappa_6: R_{L_6, \chi_6}^\circ \rightarrow N_6$ such that $(R_{L_6, \chi_6}^\circ, N_6, \text{Sh}^{(L_6, \chi_6)}, \kappa_6, \tau_{L_6, \chi_6})$ satisfies $(\dagger)$ over $\mathbb{Z}[1/6]$.*

Proof. As before, κ_6 is given by sending $(Z \rightarrow S, \alpha, \rho) \in R_{L_6, \chi_6}^\circ(S)$ to its fixed locus $Z^{\mu_2} \rightarrow S$ (see [AK11, Proof of Theorem 2.7, p. 1452] for the argument, valid in any characteristic, that each geometric fiber $Z_s^{\mu_2}$ is a non-special curve of genus 6). The construction described above gives a set-theoretic section on geometric points, and τ_{L_6, χ_6} is supplied by Lemma 6.4. $\square$

7.5 Five points on a line

Kondō has also explained how, in favorable cases, one can associate a structured K3 surface with certain configuration spaces of points.

For instance, as in [Kon07], consider the moduli space $\tilde{M}_{0,5}$ of five distinct ordered points in $\mathbb{P}^1$. (In fact, our discussion extends to the case of *stable* configurations of points.)

Fix an embedding $\beta: \mathbb{P}^1 \hookrightarrow \mathbb{P}^2$ as a coordinate line, and let $Q_\infty \in \mathbb{P}^2$ denote a point “at infinity” which is not contained in $\beta(\mathbb{P}^1)$.

Initially, let k be an algebraically closed field, and let $(P_1, \dots, P_5) \in \tilde{M}_{0,5}(k)$ be an ordered 5-tuple of distinct points. Following [Kon07, §§ 3.1 and 3.2], let C be the cyclic degree 5 cover

of $\mathbb{P}^1$ ramified exactly at $P_1, \dots, P_5$. It naturally admits a model as a plane curve inside $\mathbb{P}^2$, intersecting $\beta(\mathbb{P}^1)$ exactly at $Q_i := \beta(P_i)$ for $i = 1, \dots, 5$. Let L_i denote the line connecting Q_i and Q_∞ , and let $E_0 = \beta(\mathbb{P}^1)$.

Next, let X be the minimal resolution of the double cover of $\mathbb{P}^2$ branched along the sextic plane curve $C + E_0$, with covering involution τ . The degree 5 automorphism of C also induces a degree 5 automorphism σ of X . Moreover, because of our construction, we can identify certain divisor (classes) on X which are fixed by τ .

Indeed, for $1 \leq i \leq 5$, there is an exceptional curve E_i of the minimal resolution of singularities corresponding to $Q_i \in C \cap E_0$. Moreover, the inverse image of L_i in X is the union of two smooth rational curves $F_{i,-}$ and $F_{i,+}$, which pass, respectively, through the two points R_- and R_+ of X lying over Q_∞ . The involution τ exchanges $F_{i,-}$ and $F_{i,+}$, and σ stabilizes each of $F_{i,-}$ and $F_{i,+}$.

Finally, there is a sixteenth tautological cycle on X , namely, the inverse image $E_{0,X}$ of E_0 in X . It is stable under σ and τ .

Let L_5 be the lattice generated by these sixteen divisors, equipped with the intersection pairing. Then $L_5 \cong V \oplus A_4(-1) \oplus A_4(-1)$, see [Kon07, Lemma 4.2], and $d(L_5) = 2$. The labeling of the original points, combined with the labeling of the points in X over Q_∞ , yields an inclusion $L_5 \hookrightarrow \text{Pic}(X)$.

There is an embedding $L_5 \hookrightarrow L_{K3}$. The orthogonal complement of L_5 is computed in [Kon07, § 4.3], and a structure ρ of $L_5^\perp$ as a μ_5 -representation is described in [Kon07, § 5.2]. Let χ_5 be the character of $\rho \oplus \rho_{\text{triv}}^{\oplus 10}$. Then X is represented by a k -point of R_{L_5, χ_5}° . Conversely, we have the following.

LEMMA 7.10. *There is a morphism $\kappa_5: R_{L_5, \chi_5}^\circ \rightarrow \tilde{M}_{0,5}$ such that $(R_{L_5, \chi_5}^\circ, \tilde{M}_{0,5}, \text{Sh}^{(L_5, \chi_5)}, \kappa_5, \tau_{L_5, \chi_5})$ satisfies $(\dagger)$ over $\mathbb{Z}[\zeta_5, 1/10]$.*

Proof. As usual, it suffices to describe κ_5 . The previous construction gives the desired inverse on geometric points, and thus we have an induced isomorphism of coarse moduli spaces.

Suppose $(Z \rightarrow S, \iota, \alpha) \in R_{L_5, \chi_5}^\circ(S)$. Then there is also an involution $\beta \in \text{Aut}_{Z/S}(S)$. (In characteristic zero, this is described in the last paragraph of the proof of [Kon07, Lemma 5.7]; in positive characteristic, this then follows from a specialization argument.) The fixed locus Z^β is the disjoint union of a curve $C \rightarrow S$ with each fiber smooth and projective of genus 6 (use loc. cit. and Lemma 7.2(c)) and a relative rational curve. Moreover, the action of μ_5 on Z restricts to an action of μ_5 on C . The lattice polarization, in particular the numbering of the cycles $E_1, \dots, E_5$, labels the fixed sections $C^{\mu_5} \rightarrow S$. The quotient curve $C/\mu_5 \rightarrow S$ has fibers of genus zero, and the sought-for configuration is $(C^{\mu_5} \subset C/\mu_5) \in \tilde{M}_{0,5}(S)$. $\square$

In this case, diagram $(\dagger)$ is part of a larger diagram of moduli stacks. By its construction, the map $\kappa_5: R_{L_5, \chi_5}^\circ \rightarrow \tilde{M}_{0,5}$ factors through $H(5, 1, 5)$. Now if $(C_S \rightarrow \mathbb{P}_S^1) \in H(5, 1, 5)(S)$, then $\text{Pic}_{C/S}^0$ has an action by $\mathbb{Z}[\zeta_5]$, of signature (5.3.1) equal to $\Sigma = \{(2, 1), (0, 3)\}$. Inside A_6 , we have $A_{\mathbb{Z}[\zeta_5], \Sigma}$, the locus of principally polarized abelian 6-folds with an action by $\mathbb{Z}[\zeta_5]$ of signature Σ . Consider the classical Torelli map $\tau_6: M_6 \rightarrow A_6$. The image of the restriction of τ_6 to $H(5, 1, 5)$ is open in $A_{\mathbb{Z}[\zeta_5], \Sigma}$.

Of course, $A_{\mathbb{Z}[\zeta_5], \Sigma}$ is a Shimura variety in its own right. The complex-analytic uniformization of $A_{\mathbb{Z}[\zeta_5], \Sigma}$ is worked out in detail in [Shi64, Case (5)]. Let $G = G_{\mathbb{Z}[\zeta_5], L_5^\perp}$, and let $\mathbb{X}_G$ be the corresponding Hermitian symmetric domain; it is isomorphic to the unit 2-ball $\mathbb{B}^2$. There is a compact open subgroup $\mathbb{K}_0 \subset G(\mathbb{A}_f)$ such that $A_{\mathbb{Z}[\zeta_5], \Sigma} \cong \text{Sh}_{\mathbb{K}_0}[G, \mathbb{X}_G]$.

(Briefly, let $M = \mathbb{Z}[\zeta_5]^{\oplus 3}$, endowed with the Hermitian form h represented by $\text{diag}(1, 1, \frac{1}{2}(1 - \sqrt{5}))$. The unitary group of (M, h) is an integral form of G , and $\mathbb{K}_0$ is the stabilizer of the lattice M . Conversely, $\mathbb{K}^{(L_5, \chi_5)}$ can be recovered from $\mathbb{K}_0$ as those group elements which act trivially on the discriminant group of L .)

Then $\mathbb{K}_0 \supset \mathbb{K}^{(L_5, \chi_5)} := \mathbb{K}_{\mathbb{Z}[\zeta_5], L_5^\perp}$, with quotient group $\mathbb{K}_0/\mathbb{K}^{(L_5, \chi_5)} \cong \text{O}(\text{disc}(L^\perp)) \cong \text{O}_3(\mathbb{F}_5) \cong \{\pm 1\} \times S_5$.

We summarize this discussion as follows.

PROPOSITION 7.11. *There is a diagram of stacks over $\mathbb{Z}[\zeta_5, 1/10]$*

$$\begin{array}{ccccc}
 & & \kappa_5 & & \\
 & \nearrow & & \searrow & \\
 R_{L_5, \chi_5}^\circ & \xrightarrow{\quad} & \tilde{H}(5, 1, 5) & \xrightarrow{\quad} & \tilde{M}_{0,5} \\
 \downarrow \tau_{L_5, \chi_5} & & \downarrow [S_5] & & \downarrow [S_5] \\
 & & H(5, 1, 5) & \xrightarrow{\quad} & M_{0,5} \\
 & & \downarrow [\mu_2] & & \\
 \text{Sh}^{(L_5, \chi_5)} & \xrightarrow{[\mathbb{K}_0/\mathbb{K}^{(L_5, \chi_5)}]} & A_{\mathbb{Z}[\zeta_5], \Sigma} & &
 \end{array}$$

where an arrow is labeled $[\Gamma]$ if it is a quotient by the finite group Γ ; the given factorization of κ_5 is, on coarse moduli spaces, a composition of isomorphisms; and $\tau_{L_5, \chi_5, \mathbb{C}}$ is an open immersion.

Proof. Since the canonical models of both $\text{Sh}^{(L_5, \chi_5)}$ and $A_{\mathbb{Z}[\zeta_5], \Sigma}$ receive maps from R_{L_5, χ_5} over $\mathbb{Z}[\zeta_5, 1/10]$ with dense image, it suffices to observe that the quotient map $\text{Sh}^{(L_5, \chi_5)} \rightarrow A_{\mathbb{Z}[\zeta_5], \Sigma}$ is defined on the canonical models. $\square$

7.6 Six points on a line

In [DK07, § 12], Dolgachev and Kondō show that the configuration space of six labeled points on the (complex) projective line is an arithmetic quotient of $\mathbb{B}^4$.

The pointwise construction in [DK07, § 12] works over an arbitrary algebraically closed field k . Let $(P_1, \dots, P_6) \in \tilde{M}_{0,6}$ be an ordered 6-tuple of distinct points. Let C be the cyclic degree 3 cover of $\mathbb{P}^1$ ramified exactly at the P_i , and let Z' be the cyclic triple cover of the ambient weighted projective space $\mathbb{P}(1, 1, 2)$ ramified along C . (Explicitly, let $f(X_0, X_1)$ be a homogeneous form of degree 6 vanishing at the P_i ; then Z' is given by the equation $X_3^3 + X_2^3 + f(X_0, X_1) = 0$ in $\mathbb{P}(1, 1, 2, 2)$.) Then Z' comes with an action by $\mu_3 \times \mu_3$; we single out the action of μ_3 on Z' via the diagonal embedding $\mu_3 \hookrightarrow \mu_3 \times \mu_3 \hookrightarrow \text{Aut}_k(Z')$. The variety Z' has three ordinary nodes; its minimal resolution Z is a K3 surface, and the μ_3 -action lifts to Z . One finds that, by construction, $\text{Pic}_{Z/k}(k)$ comes equipped with a primitive inclusion of the lattice $L'_6 := U \oplus E_6(-1) \oplus A_2(-1)^{\oplus 3}$, with orthogonal complement $A_2(1) \oplus A_2(-1)^{\oplus 3}$, and that $d(L'_6) = 2$. As a $\mathbb{Z}[\zeta_3]$ -module, $(L'_6)^\perp$ is free of rank 4 and comes equipped with a Hermitian form of signature $(3, 1)$. Let ρ be the corresponding μ_3 -representation, and let χ'_6 be the character of $\rho \oplus \rho_{\text{triv}}^{\oplus 14}$. As usual, we have $(\dagger)$ for $R_{(L'_6, \chi'_6)}^\circ$, $\tilde{M}_{0,6}$, and $\text{Sh}^{(L'_6, \chi'_6)}$ over $\mathbb{Z}[\zeta_3, 1/6]$.

Alternatively, we could use the strategy of Section 7.5 and compute the periods of C directly. If $(C \rightarrow S \rightarrow \mathbb{P}_S^1) \in H(3, 1, 6)$, then C/S is a family of curves of genus 4 and $\text{Pic}_{C/S}^0$ has an action by $\mathbb{Z}[\zeta_3]$ of signature $(1, 3)$. (This is case (2) of [Shi64].) The moduli space $A_{\mathbb{Z}[\zeta_3], (1, 3)}$ of principally polarized abelian fourfolds with action by $\mathbb{Z}[\zeta_3]$ of signature $(1, 3)$ is isomorphic

to $\mathrm{Sh}_{\mathbb{K}_0}[G, \mathbb{X}_G]$, where $G = G_{\mathbb{Z}[\zeta_3], (L'_6)^\perp}$ and $\mathbb{K}_0$ is the stabilizer of the lattice $(L'_6)^\perp$. There is a surjection $\mathrm{Sh}^{(L'_6, \mathbb{X}'_6)} \rightarrow \mathrm{Sh}_{\mathbb{K}_0}[G, \mathbb{X}_G]$ with covering map $\mathbb{K}_0/\mathbb{K}^{(L'_6, \mathbb{X}'_6)} \cong \mathrm{O}(\mathrm{disc}((L'_6)^\perp)) \cong \mu_2 \times S_6$ (it seems that, in the third displayed equation of [DK07, § 12, second example, p. 93], the authors may have neglected to account for the discriminant kernel), and we obtain the following.

PROPOSITION 7.12. *There is a diagram of stacks over $\mathbb{Z}[\zeta_3, 1/6]$*

$$\begin{array}{ccccc}
 & & \kappa'_6 & & \\
 & \xrightarrow{\quad} & \widetilde{H}(3, 1, 6) & \xrightarrow{\quad} & \widetilde{M}_{0,6} \\
 R_{L'_6, \mathbb{X}'_6}^\circ & \xrightarrow{\quad} & \downarrow [S_6] & & \downarrow [S_6] \\
 & \xrightarrow{\quad} & H(3, 1, 6) & \xrightarrow{\quad} & M_{0,6} \\
 & \xrightarrow{\quad} & \downarrow [\mu_2] & & \\
 & \xrightarrow{\quad} & A_{\mathbb{Z}[\zeta_3], (1,3)} & & \\
 \downarrow \tau_{L'_6, \mathbb{X}'_6} & & & & \\
 \mathrm{Sh}^{(L'_6, \mathbb{X}'_6)} & \xrightarrow{[\mathbb{K}_0/\mathbb{K}^{(L'_6, \mathbb{X}'_6)}]} & A_{\mathbb{Z}[\zeta_3], (1,3)} & &
 \end{array}$$

where an arrow is labeled $[\Gamma]$ if it is a quotient by the finite group Γ ; the given factorization of κ'_6 is, on coarse moduli spaces, a composition of isomorphisms; and $\tau_{L'_6, \mathbb{X}'_6, \mathbb{C}}$ is an open immersion.

7.7 Cubic surfaces

Let Cub_2 be the moduli space of cubic surfaces. If $V/\mathbb{C}$ is a complex cubic surface, then by associating either a cubic threefold [ACT02] or a K3 surface [DvGK05] with it and measuring its periods, one obtains an open immersion $\mathrm{Cub}_2(\mathbb{C}) \hookrightarrow \mathbb{B}^4/\Gamma$. The arithmetic nature of this map is explored in [Ach14]. Unfortunately, there is a stack-theoretic mistake there. While Cub_2 and $H(3, 3, 3)$ have isomorphic coarse moduli spaces, they are not literally the same stack; $H(3, 3, 3) \rightarrow \mathrm{Cub}_2$ is the rigidification along the μ_3 -action. We take the present opportunity to correct this oversight and recast the main result in the framework developed here.

Let Cub_3 be the moduli space of smooth projective cubic *threefolds*. If $T \in \mathrm{Cub}_3(\mathbb{C})$, then its intermediate Jacobian is a principally polarized abelian fivefold. This gives a period map $\mathrm{Cub}_3(\mathbb{C}) \rightarrow A_5(\mathbb{C})$, which is known to be an embedding. By using either monodromy considerations [Del72, Theorem 2.12] or the arithmetic nature of intermediate Jacobians [ACV20, Theorem 6.1], one can show that this period map descends to a morphism $\mathrm{Cub}_3 \rightarrow A_5$ of stacks over $\mathbb{Q}$. Using the algebro-geometric construction of the intermediate Jacobian as a Prym variety, one can actually spread this out to achieve a morphism $\mathrm{Cub}_3 \rightarrow A_5$ of stacks over $\mathbb{Z}[1/2]$; see [Ach14, Corollary 3.5].

Now, points of $H(3, 3, 3)$ correspond to cyclic triple covers of $\mathbb{P}^3$ branched along a cubic surface; as such, they are smooth projective threefolds in their own right. The μ_3 -action on the threefold induces an action of $\mathbb{Z}[\zeta_3]$ on the corresponding intermediate Jacobian, with signature $(1, 4)$. Ultimately, one obtains the following.

PROPOSITION 7.13. *There is a diagram of stacks over $\mathbb{Z}[\zeta_3, 1/6]$*

$$\begin{array}{ccccc}
 \mathrm{Cub}_3 & \hookleftarrow & H(3, 3, 3) & \xrightarrow{\kappa} & \mathrm{Cub}_2 \\
 \downarrow & & \downarrow \tau & & \\
 A_5 & \hookleftarrow & A_{\mathbb{Z}[\zeta_3], (1,4)} & &
 \end{array}$$

in which τ is an open immersion and κ induces an isomorphism of coarse moduli spaces.

ACKNOWLEDGEMENTS

My interest in this circle of ideas was sparked when S. Casalaina-Martin told me about the remarkable identification of the moduli of complex cubic surfaces as a ball quotient [ACT02, DvGK05]. It's a pleasure to thank him for continued inspiration and insights. I thank M. Rapoport, K. Madapusi Pera, and the referee for helpful comments, and I. Dolgachev for suggesting that this paper should address the quasi-polarized situation.

REFERENCES

- Ach14 J. D. Achter, *Arithmetic Torelli maps for cubic surfaces and threefolds*, Trans. Amer. Math. Soc. **366** (2014), no. 11, 5749–5769; doi:10.1090/S0002-9947-2014-05978-3.
- ACV20 J. D. Achter, S. Casalaina-Martin, and C. Vial, *Distinguished models of intermediate Jacobians*, J. Inst. Math. Jussieu **19** (2020), no. 3, 891–918; doi:10.1017/S1474748018000245.
- ACT02 D. Allcock, J. A. Carlson, and D. Toledo, *The complex hyperbolic geometry of the moduli space of cubic surfaces*, J. Algebraic Geom. **11** (2002), no. 4, 659–724; doi:10.1090/S1056-3911-02-00314-4.
- ACV03 D. Abramovich, A. Corti, and A. Vistoli, *Twisted bundles and admissible covers*, Comm. Algebra **31** (2003), no. 8, 3547–3618; doi:10.1081/AGB-120022434.
- AK11 M. Artebani and S. Kondō, *The moduli of curves of genus six and K3 surfaces*, Trans. Amer. Math. Soc. **363** (2011), no. 3, 1445–1462; doi:10.1090/S0002-9947-2010-05126-8.
- AST11 M. Artebani, A. Sarti, and S. Taki, *K3 surfaces with non-symplectic automorphisms of prime order (with an appendix by Shigeyuki Kondō)*, Math. Z. **268** (2011), no. 1-2, 507–533; doi:10.1007/s00209-010-0681-x.
- AV04 A. Arsie and A. Vistoli, *Stacks of cyclic covers of projective spaces*, Compos. Math. **140** (2004), no. 3, 647–666; doi:10.1112/S0010437X03000253.
- Bea04 A. Beauville, *Fano threefolds and K3 surfaces*, The Fano Conference (University Torino, Turin, 2004), 175–184.
- BO83 P. Berthelot and A. Ogus, *F-isocrystals and de Rham cohomology. I*, Invent. Math. **72** (1983), no. 2, 159–199; doi:10.1007/BF01389319.
- Del71 P. Deligne, *Travaux de Shimura*, Séminaire Bourbaki, 23ème année (1970/71), Exp. No. 389, Lecture Notes in Math., vol. 244 (Springer, Berlin, 1971), 123–165; doi:10.1007/BFb0058700.
- Del72 ———, *Les intersections complètes de niveau de Hodge un*, Invent. Math. **15** (1972), 237–250; doi:10.1007/BF01404128.
- Del79 ———, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, Automorphic Forms, Representations and L-Functions (Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII (Amer. Math. Soc., Providence, RI, 1979), 247–289.
- DI81 P. Deligne and L. Illusie, *Relèvement des surfaces K3 en caractéristique nulle*, Algebraic Surfaces (Orsay, 1976–78), Lecture Notes in Math., vol. 868 (Springer, Berlin – New York, 1981), 58–79; doi:10.1007/BFb0090646.
- DK07 I. V. Dolgachev and S. Kondō, *Moduli of K3 surfaces and complex ball quotients*, in *Arithmetic and geometry around hypergeometric functions*, Progr. Math., vol. 260 (Birkhäuser, Basel, 2007), 43–100; doi:10.1007/978-3-7643-8284-1_3.
- DM86 P. Deligne and G. D. Mostow, *Monodromy of hypergeometric functions and nonlattice integral monodromy*, Publ. Math. Inst. Hautes Études Sci. **63** (1986), 5–89; doi:10.1007/BF02831622.
- Dol96 I. V. Dolgachev, *Mirror symmetry for lattice polarized K3 surfaces*, J. Math. Sci. **81** (1996), no. 3, 2599–2630; doi:10.1007/BF02362332.

- DvGK05 I. Dolgachev, B. van Geemen, and S. Kondō, *A complex ball uniformization of the moduli space of cubic surfaces via periods of K3 surfaces*, J. reine angew. Math. **588** (2005), 99–148; doi:10.1515/crll.2005.2005.588.99.
- GD64 A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I*, Publ. Math. Inst. Hautes Études Sci. **20** (1964), 5–251; doi:10.1007/BF02684747.
- Huy16 D. Huybrechts, *Lectures on K3 surfaces*, Cambridge Stud. Adv. Math., vol. 158 (Cambridge Univ. Press, Cambridge, 2016); doi:10.1017/CB09781316594193.
- Jan17 J. Jang, *A lifting of an automorphism of a K3 surface over odd characteristic*, Int. Math. Res. Not. **2017** (2017), no. 6, 1787–1804; doi:10.1093/imrn/rnw071.
- Keu14 J. Keum, *K3 surfaces with an order 60 automorphism and a characterization of supersingular K3 surfaces with Artin invariant 1*, Math. Res. Lett. **21** (2014), no. 3, 509–520; doi:10.4310/MRL.2014.v21.n3.a8.
- Kis10 M. Kisin, *Integral models for Shimura varieties of abelian type*, J. Amer. Math. Soc. **23** (2010), no. 4, 967–1012; doi:10.1090/S0894-0347-10-00667-3.
- KM74 N. M. Katz and W. Messing, *Some consequences of the Riemann hypothesis for varieties over finite fields*, Invent. Math. **23** (1974), 73–77; doi:10.1007/BF01405203.
- Kon00 S. Kondō, *A complex hyperbolic structure for the moduli space of curves of genus three*, J. reine angew. Math. **525** (2000), 219–232; doi:10.1515/crll.2000.069.
- Kon02 ———, *The moduli space of curves of genus 4 and Deligne–Mostow’s complex reflection groups*, Algebraic Geometry 2000, Azumino (Hotaka), Adv. Stud. Pure Math., vol. 36 (Math. Soc. Japan, Tokyo, 2002), 383–400; doi:10.2969/aspm/03610383.
- Kon07 ———, *The moduli space of 5 points on $\mathbb{P}^1$ and K3 surfaces*, in *Arithmetic and Geometry around Hypergeometric Functions*, Progr. Math., vol. 260 (Birkhäuser, Basel, 2007), 189–206; doi:10.1007/978-3-7643-8284-1_7.
- KR12 S. Kudla and M. Rapoport, *On occult period maps*, Pacific J. Math. **260** (2012), no. 2, 565–581; doi:10.2140/pjm.2012.260.565.
- Kud13 S. Kudla, *A note about special cycles on moduli spaces of K3 surfaces*, in *Arithmetic and Geometry of K3 Surfaces and Calabi–Yau Threefolds*, Fields Inst. Commun., vol. 67 (Springer, New York, 2013), 411–427; doi:10.1007/978-1-4614-6403-7_14.
- LM00 G. Laumon and L. Moret-Bailly, *Champs algébriques*, Ergeb. Math. Grenzgeb. (3), vol. 39 (Springer-Verlag, Berlin, 2000); doi:10.1007/978-3-540-24899-6.
- Lov17 T. Lovering, *Integral canonical models for automorphic vector bundles of abelian type*, Algebra Number Theory **11** (2017), no. 8, 1837–1890; doi:10.2140/ant.2017.11.1837.
- Mad15 K. Madapusi Pera, *The Tate conjecture for K3 surfaces in odd characteristic*, Invent. Math. **201** (2015), no. 2, 625–668; doi:10.1007/s00222-014-0557-5.
- Mad16 ———, *Integral canonical models for spin Shimura varieties*, Compos. Math. **152** (2016), no. 4, 769–824; doi:10.1112/S0010437X1500740X.
- Mau14 D. Maulik, *Supersingular K3 surfaces for large primes (with an appendix by Andrew Snowden)*, Duke Math. J. **163** (2014), no. 13, 2357–2425; doi:10.1215/00127094-2804783.
- MFK94 D. Mumford, J. Fogarty, and F. Kirwan, *Geometric invariant theory*, 3rd ed., Ergebn. Math. Grenzgeb. (2), vol. 34 (Springer-Verlag, Berlin, 1994); doi:10.1007/978-3-642-57916-5.
- Mil05 J. S. Milne, *Introduction to Shimura varieties*, in *Harmonic Analysis, the Trace Formula, and Shimura Varieties*, Clay Math. Proc., vol. 4 (Amer. Math. Soc., Providence, RI, 2005), 265–378.
- MM64 T. Matsusaka and D. Mumford, *Two fundamental theorems on deformations of polarized varieties*, Amer. J. Math. **86** (1964), 668–684; doi:10.2307/2373030.
- MO98 N. Machida and K. Oguiso, *On K3 surfaces admitting finite non-symplectic group actions*, J. Math. Sci. Univ. Tokyo **5** (1998), no. 2, 273–297.

- Nik79 V. V. Nikulin, *Integer symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), no. 1, 111–177.
- NO85 N. Nygaard and A. Ogus, *Tate’s conjecture for K3 surfaces of finite height*, Ann. of Math. **122** (1985), no. 3, 461–507; doi:10.2307/1971327.
- Ogu84 A. Ogus, *Periods of integrals in characteristic p* , Proceedings of the International Congress of Mathematicians, Vols. 1, 2 (Warsaw, 1983) (PWN, Warsaw, 1984), 753–762.
- Riz06 J. Rizov, *Moduli stacks of polarized K3 surfaces in mixed characteristic*, Serdica Math. J. **32** (2006), no. 2–3, 131–178.
- Riz10 ———, *Kuga–Satake abelian varieties of K3 surfaces in mixed characteristic*, J. reine angew. Math. **648** (2010), 13–67; doi:10.1515/CRELLE.2010.078.
- Rom05 M. Romagny, *Group actions on stacks and applications*, Michigan Math. J. **53** (2005), no. 1, 209–236; doi:10.1307/mmj/1114021093.
- SGA5 L. Illusie (ed.), *Séminaire de Géométrie Algébrique du Bois-Marie, 1965–66 (SGA5), Cohomologie l -adique et Fonctions L* , dirigé par A. Grothendieck avec la collaboration de I. Bucur, C. Houzel, L. Illusie, J.-P. Jouanolou et J.-P. Serre, Lecture Notes in Math., vol. 589 (Springer-Verlag, Berlin – New York, 1977); doi:10.1007/BFb0096802
- Shi64 G. Shimura, *On purely transcendental fields automorphic functions of several variable*, Osaka Math. J. **1** (1964), no. 1, 1–14.
- Sta17 The Stacks Project Authors, *The Stacks Project*, version 2017, available at <http://stacks.math.columbia.edu>.
- Tae17 L. Taelman, *Complex multiplication and Shimura stacks*, Motives and Complex Multiplication (Ascona, 2016), Progr. Math., to appear, arXiv:1707.01236.
- Zhe20 Z. Zheng, *Orbifold aspects of certain occult period maps*, Nagoya Math. J., published online on 27 November 2019, doi:10.1017/nmj.2019.36, to appear in print.

Jeffrey D. Achter j.achter@colostate.edu

Department of Mathematics, Colorado State University, Fort Collins, CO 80523, USA