Homological mirror symmetry for Milnor fibers of simple singularities

Yankı Lekili and Kazushi Ueda

ABSTRACT

We prove homological mirror symmetry for Milnor fibers of simple singularities in dimensions greater than 1, which are among the log Fano cases of Conjecture 1.5 of the authors’ manuscript “Homological mirror symmetry for Milnor fibers via moduli of A_∞ -structures” [arXiv: 1806.04345]. The proof is based on a relation between matrix factorizations and Calabi–Yau completions. As an application, we give an explicit computation of the Hochschild cohomology group of the derived n -preprojective algebra of a Dynkin quiver for any $n \geq 1$, and the symplectic cohomology group of the Milnor fiber of any simple singularity in any dimension greater than 1.

1. Introduction

A *simple singularity* is an isolated hypersurface singularity of modality 0. Arnold classified such singularities; up to right equivalence, they are given by one of the following:

$$\begin{aligned} A_\ell: x_1^{\ell+1} + x_2^2 + \cdots + x_{n+1}^2 &= 0, \quad \ell = 1, 2, \dots, \\ D_\ell: x_1^{\ell-1} + x_1 x_2^2 + x_3^2 + \cdots + x_{n+1}^2 &= 0, \quad \ell = 4, 5, \dots, \\ E_6: x_1^4 + x_2^3 + x_3^2 + \cdots + x_{n+1}^2 &= 0, \\ E_7: x_1^3 + x_1 x_2^3 + x_3^2 + \cdots + x_{n+1}^2 &= 0, \\ E_8: x_1^5 + x_2^3 + x_3^2 + \cdots + x_{n+1}^2 &= 0. \end{aligned} \tag{1.1}$$

In the case $n = 2$, simple surface singularities have many other characterizations, such as Kleinian singularities, rational double points, or canonical singularities, to name a few.

Let $\tilde{w}$ be one of these defining polynomials, which we think of as a holomorphic function on $\mathbb{C}^{n+1}$, and equip $\tilde{w}^{-1}(1)$ with the Liouville structure induced from the standard one on $\mathbb{C}^{n+1}$. This is the Liouville completion of the *Milnor fiber*, which is the Liouville domain obtained by intersecting $\tilde{w}^{-1}(1)$ with a ball. Let $\mathcal{W}(\tilde{w}^{-1}(1))$ denote the idempotent-complete derived wrapped Fukaya category of $\tilde{w}^{-1}(1)$.

For $n \geq 2$, since $\tilde{w}^{-1}(1)$ is not a log Calabi–Yau manifold but a log Fano manifold, its mirror is not a manifold but a *Landau–Ginzburg model*, by which we mean a pair of a stack and a section

Received 1 May 2020, accepted in final form 11 February 2021.

2020 *Mathematics Subject Classification* 53D37.

Keywords: homological mirror symmetry, simple singularity, symplectic cohomology.

This journal is © **Foundation Compositio Mathematica** 2021. This article is distributed with Open Access under the terms of the **Creative Commons Attribution Non-Commercial License**, which permits non-commercial reuse, distribution, and reproduction in any medium, provided that the original work is properly cited. For commercial re-use, please contact the **Foundation Compositio Mathematica**.

Y. L. is partially supported by the Royal Society URF\R\180024. K. U. is partially supported by Grant-in-Aid for Scientific Research (15KT0105, 16K13743, 16H03930).

of a line bundle on it. One way to obtain a Landau–Ginzburg mirror of a log Fano manifold is to first remove a divisor to make it log Calabi–Yau, then find its mirror, which is another log Calabi–Yau manifold, and finally add a potential to this mirror [Aur07, Aur09]. This produces a Landau–Ginzburg mirror whose underlying manifold is of the same dimension as the original manifold. When the singularity is toric (that is, a simple surface singularity of type A), there is a standard choice for the divisor to remove, and the resulting mirror is the Landau–Ginzburg model consisting of a complement of a toric divisor in the minimal resolution of the singularity of the same type and a monomial function on it (see, for example, [AAK16, Section 9.2]). The choice of the divisor is not unique in general, and there are multiple mirrors for a given Milnor fiber.

In this paper, we consider an alternative mirror of the Milnor fiber of a simple singularity based on the transposition of invertible polynomials introduced in [BH93, BH95]. A weighted homogeneous polynomial $\mathbf{w} \in \mathbb{C}[x_1, \dots, x_{n+1}]$ with an isolated critical point at the origin is *invertible* if there is an integer matrix $A = (a_{ij})_{i,j=1}^{n+1}$ with non-zero determinant such that

$$\mathbf{w} = \sum_{i=1}^{n+1} \prod_{j=1}^{n+1} x_j^{a_{ij}}.$$

The *transpose* of $\mathbf{w}$ is defined as

$$\check{\mathbf{w}} = \sum_{i=1}^{n+1} \prod_{j=1}^{n+1} x_j^{a_{ji}},$$

whose exponent matrix $\check{A}$ is the transpose matrix of A . The group

$$\Gamma_{\mathbf{w}} := \{(t_0, t_1, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+2} \mid t_1^{a_{1,1}} \dots t_{n+1}^{a_{1,n+1}} = \dots = t_1^{a_{n+1,1}} \dots t_{n+1}^{a_{n+1,n+1}} = t_0 t_1 \dots t_{n+1}\}$$

acts naturally on $\mathbb{A}^{n+2} := \text{Spec } \mathbb{C}[x_0, \dots, x_{n+1}]$. Let $\text{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w} + x_0 \dots x_{n+1})$ denote the idempotent completion of the dg category of $\Gamma_{\mathbf{w}}$ -equivariant coherent matrix factorizations of $\mathbf{w} + x_0 \dots x_{n+1}$ on $\mathbb{A}^{n+2}$ in the sense of [EP15]. Conjecture 1.1 below is given in [LU18, Conjecture 1.5].

CONJECTURE 1.1. For any invertible polynomial $\mathbf{w}$, one has a quasi-equivalence

$$\text{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w} + x_0 \dots x_{n+1}) \simeq \mathcal{W}(\check{\mathbf{w}}^{-1}(1)).$$

In other words, the Landau–Ginzburg model $([\mathbb{A}^{n+2}/\Gamma_{\mathbf{w}}], \mathbf{w} + x_0 \dots x_{n+1})$ is mirror to the Liouville manifold $\check{\mathbf{w}}^{-1}(1)$. The main result of this paper is the following.

THEOREM 1.2. Conjecture 1.1 holds for $n \geq 2$ and $\check{\mathbf{w}}$ one of the defining polynomials of simple singularities appearing in (1.1).

The proof of Theorem 1.2 consists of four steps. The first step is the quasi-equivalence

$$\text{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w} + x_0 \dots x_{n+1}) \simeq \text{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w}), \quad (1.2)$$

which comes from the fact that $\mathbf{w} + x_0 \dots x_{n+1}$ is right equivalent to $\mathbf{w}$ by a formal change of variables, which holds if $n \geq 2$ and $\mathbf{w}$ defines a simple singularity.

The second step is the quasi-equivalence

$$\text{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w}) \simeq \Pi_n(\text{mf}(\mathbb{A}^{n+1}, \Gamma_{\mathbf{w}}, \mathbf{w})), \quad (1.3)$$

where Π_n denotes the n -Calabi–Yau completion in the sense of [Kel11]. This holds for any invertible polynomial $\mathbf{w}$ and any $n \geq 0$.

The third step is the quasi-equivalence

$$\mathrm{mf}(\mathbb{A}^{n+1}, \Gamma_{\mathbf{w}}, \mathbf{w}) \simeq \mathrm{perf} A_Q \quad (1.4)$$

with the dg category $\mathrm{perf} A_Q$ of perfect dg modules over the path algebra A_Q of a Dynkin quiver Q (with any orientation) of the corresponding type. For type A, this is proved in [Tak05, Theorem 3.1] for $n = 0$, and the $n \geq 1$ case follows either from the $n = 0$ case and the Knörrer periodicity [Knö87], or as a special case of [FU11, Theorem 1.2]. For type D, this follows from [FU13, Section 4]. For type E, this follows either by combining [HS20, Theorem 1] and [Sei01, Proposition 3.4] or by finding a suitable mutation of a generator appearing in [HS20, Theorem 2]. Note that [KST07, Theorem 3.1] gives a result close to (1.4), which is not exactly the same since the grading group is different.

The last step is

$$\mathcal{W}(\check{w}^{-1}(1)) \simeq \Pi_n(\mathrm{perf} A_Q), \quad (1.5)$$

which holds if $n \geq 2$ and $\check{w}$ defines a simple singularity. As discussed in Section 3, the proof of (1.5) in [EL17b] depends on the computation of the Hochschild cohomology of $\Pi_n(\mathrm{perf} A_Q)$, which was missing for type E cases in [EL17b] and is done in Section 5.

For $n = 1$ not covered by Theorem 1.2, a quasi-equivalence of the full subcategory $\mathcal{F}(\check{w}^{-1}(0))$ of $\mathcal{W}(\check{w}^{-1}(0))$ consisting of (direct summands of bounded complexes of) compact Lagrangians and a category $\mathrm{perf} Z_{\mathbf{w}}$ equivalent to the full subcategory of $\mathrm{mf}(\mathbb{A}^3, \Gamma_{\mathbf{w}}, \mathbf{w} + x_0 x_1 x_2)$ consisting of homologically finite objects (that is, those X satisfying $\dim \bigoplus_{i \in \mathbb{Z}} \mathrm{Ext}^i(X, Y) < \infty$ for any object Y) is given in [Hab20, Theorem 1.1].

As an application of (1.3), we compute the Hochschild cohomology group of the n -Calabi–Yau completion $\Pi_n(A_Q)$, also known as the *derived n -preprojective algebra*, of the path algebra A_Q of any Dynkin quiver Q for any $n \geq 1$. It is possible to compute the Hochschild homology along the same lines.

The zeroth cohomology of the derived 2-preprojective algebra is the *preprojective algebra*. The Hochschild homology and cohomology of the preprojective algebra of the path algebra of a Dynkin quiver is calculated in [ES98a, ES98b, ES98c, EE07]. Even the calculus structure in the sense of [GDT90, TT00] (which includes the Batalin–Vilkovisky structure and is known to be derived invariant [AK19]) is calculated in [Eu10], and it is an interesting problem to do the same for the derived n -preprojective algebra. Note that the preprojective algebra and the derived 2-preprojective algebra of a Dynkin quiver are very different. The derived 2-preprojective algebra of a Dynkin quiver is a smooth dg algebra, which has cohomology in every negative cohomological degree and moreover is not formal. In contrast, the preprojective algebra is always concentrated in cohomological degree 0 by definition, and the global dimension is infinite for a Dynkin quiver.

It follows from [Gan12, Theorem 1.1] combined with [CDGG17, Theorem 1.4], which builds on [Gan12, Gao17], that the closed-open map of any Weinstein manifold from the symplectic cohomology to the Hochschild cohomology of the wrapped Fukaya category is an isomorphism:

$$\mathrm{SH}^*(M) \xrightarrow{\sim} \mathrm{HH}^*(\mathcal{W}(M)).$$

Hence, by Theorem 1.2, we see that the symplectic cohomology of the Milnor fiber $\check{w}^{-1}(1)$ of a simple singularity for $n \geq 2$ is isomorphic to $\mathrm{HH}^*(\Pi_n(A_Q))$. This enables us to give an explicit computation of the symplectic cohomology of Milnor fibers of all simple singularities in a uniform way. Previous partial results computing symplectic cohomology for Milnor fibers of simple singularities appeared in [EL17b] for A_ℓ and D_ℓ in complex dimension 2 and in [KvK16, Ueb16]

for various versions of symplectic cohomology for certain higher-dimensional A_ℓ -Milnor fibers for which an associated Morse–Bott spectral sequence yields computations. Our computation also shows that $\mathrm{HH}^*(\Pi_1(A_Q))$ is not isomorphic to $\mathrm{SH}^*(\tilde{w}^{-1}(0))$ given in [Hab20, Section 3.3], which is consistent with the failure of (1.5) for $n = 1$.

This paper is organized as follows: In Section 2, we collect basic definitions and results on Calabi–Yau completions and trivial extension algebras. In Section 3, we recall the description of the wrapped Fukaya category of the Milnor fiber of a simple singularity for $n \geq 2$ in terms of the n -Calabi–Yau completion of a Dynkin quiver of the corresponding type. In Section 4, we prove (1.2) and (1.3). The computation of Hochschild cohomologies of the derived preprojective algebras of Dynkin quivers are given in Section 5.

2. Calabi–Yau completions and trivial extension algebras

The n -Calabi–Yau completion (or the derived n -preprojective algebra) of a dg category $\mathcal{A}$ is defined in [Kel11, Section 4.1] as the tensor algebra

$$\Pi_n(\mathcal{A}) := T_{\mathcal{A}}(\theta) := \mathcal{A} \oplus \theta \oplus \theta \otimes_{\mathcal{A}} \theta \oplus \cdots,$$

where the $\mathcal{A}$ -bimodule $\theta := \Theta[n-1]$ is a shift of the inverse dualizing complex

$$\Theta := \mathrm{hom}_{\mathcal{A}^e}(\mathcal{A}, \mathcal{A}^e).$$

A dg algebra is regarded as a dg category with one object. The Morita invariance of the Calabi–Yau completion shown in [Kel11, Proposition 4.2] implies that Calabi–Yau completion commutes with the operation of taking the dg category of perfect dg modules:

$$\Pi_n(\mathrm{perf}\mathcal{A}) \simeq \mathrm{perf}(\Pi_n \mathcal{A}).$$

The Ginzburg dg algebra $\mathcal{G}_Q^n$ of a quiver Q (without potential) is a model of the n -Calabi–Yau completion $\Pi_n(A_Q)$ of the path algebra A_Q , defined in [Kel11, Section 6.2] after [Gin06] as the path algebra of the graded quiver $\overline{Q}$ with same vertices as Q and arrows consisting of

- the original arrows $g \in Q_1$ in degree 1,
- the opposite arrows g^* for each arrow $g \in Q_1$ in degree $1 - n$, and
- loops h_v at each vertex $v \in Q_0$ in degree $1 - n$,

equipped with the differential d given by

$$dg = dg^* = 0 \quad \text{and} \quad dh = \sum_{g \in Q_1} g^*g - gg^*,$$

where $h = \sum_{v \in Q_0} h_v$.

The degree n trivial extension algebra of a finite-dimensional algebra A is defined as $A \oplus A^\vee[-n]$ equipped with the multiplication $(a, f) \cdot (b, g) = (ab, ag + fb)$, where $A^\vee$ is the dual of A as a vector space.

The degree n trivial extension algebra B_Q^n of the path algebra A_Q of a Dynkin quiver Q is the (derived) Koszul dual of $\mathcal{G}_Q^n$ in the sense that

$$\mathrm{hom}_{\mathcal{G}_Q^n}(\mathbf{k}_{\mathcal{G}}, \mathbf{k}_{\mathcal{G}}) \simeq B_Q^n, \quad \mathrm{hom}_{(B_Q^n)^{\mathrm{op}}}(\mathbf{k}_B, \mathbf{k}_B) \simeq (\mathcal{G}_Q^n)^{\mathrm{op}}, \quad (2.1)$$

where $\mathbf{k}_{\mathcal{G}} := \bigoplus_{v \in Q_0} S_v$ is the direct sum of simple left $\mathcal{G}_Q^n$ -modules S_v associated with vertices $v \in Q_0$, and similarly for $\mathbf{k}_B$ (see, for example, [EL17b, Theorem 23, Corollary 25]).

This Koszul duality implies an isomorphism

$$\mathrm{HH}^*(\mathcal{G}_Q^n) \cong \mathrm{HH}^*(B_Q^n) \quad (2.2)$$

of Hochschild cohomologies (see, for example, [FMT05, Theorem 1] and [Her19, Theorem 3.4]).

3. The wrapped Fukaya category of the Milnor fiber of a simple singularity

Let $\tilde{w}$ be one of the defining polynomials of a simple singularity and $M^n = \tilde{w}^{-1}(1)$ be the Milnor fiber, which we view as a Weinstein manifold where the Weinstein structure is induced by restriction from the ambient $\mathbb{C}^{n+1}$. It is well known that this Weinstein manifold is symplectomorphic (in fact, Weinstein homotopic) to the plumbing X_Q of cotangent bundles of spheres T^*S^n according to the Dynkin diagram Q corresponding to the simple singularity. One way to see this is to verify it directly for $n = 1$ and then use the fact that in higher dimensions, the Milnor fiber is obtained by stabilization—increasing the dimension corresponds to the suspension of the Lefschetz fibration [Sei10]. See also [Abo11] for an explicit construction of a symplectic structure on plumbings. This stabilization point of view also enables one to describe M via Legendrian surgery. Namely, M is obtained by attaching critical handles to a Legendrian link Λ_Q^{n-1} on $\partial\mathbb{D}^n$ whose components are unknotted Legendrian spheres S^{n-1} which are clasped together (as in Hopf link) according to the Dynkin diagram Q . The direct sum of co-cores to the critical handles (that is, cotangent fibers away from the plumbing region) form a generating object of the wrapped Fukaya category by the main theorem in [CDGG17], and the surgery formula of [BEE12, Ekh19] allows one to explicitly compute the endomorphism algebra of this generator as the Chekanov–Eliashberg algebra $\mathrm{CE}^*(\Lambda_Q^{n-1})$.

This Chekanov–Eliashberg algebra was computed directly in the case $n = 2$ in the paper [EL17b], and the resulting dg algebra was shown to be quasi-isomorphic to the derived multiplicative preprojective algebra of the corresponding Dynkin type. Moreover, working over $\mathbb{C}$, for $Q = A_\ell$ or D_ℓ , it was shown in [EL17b, Theorem 13] that the derived multiplicative preprojective algebra of Dynkin type Q is quasi-isomorphic to the Ginzburg algebra $\mathcal{G}_Q^2$, also known as the derived (additive) preprojective algebra of Dynkin type Q . It was conjectured in op. cit. that the same result holds for $Q = E_6, E_7, E_8$, and this is indeed so. The key ingredient for the proof of [EL17b, Theorem 13] to go through that was missing in the case $Q = E_6, E_7, E_8$ was the computation that

$$\mathrm{HH}^2(\mathcal{G}_Q^2)^s = 0 \quad \text{for } s < 0,$$

but this follows from computations given in Section 5 below.

For $n \geq 3$, one can do a direct computation in an analogous way, but we can also deduce this by the Koszul duality result given in [EL17a, Theorem 58], which shows that $\mathrm{CE}^*(\Lambda_Q^{n-1})$ is the (derived) Koszul dual of the endomorphism algebra of the union of the core spheres of the plumbing. Notice that for $n \geq 3$, the polynomial $\tilde{w}$ is suspended at least twice; thus, the formality of the endomorphism algebra of vanishing cycles in the compact Fukaya category of $\tilde{w}^{-1}(1)$ follows automatically by [Sei10, Proposition 4.4] (the formality of the A_∞ -algebra $\mathcal{A}$ and an $\mathcal{A}$ -bimodule $\mathcal{B}/\mathcal{A}$ in Seidel’s notation is obvious in the case at hand, since Γ is a tree and one can shift the objects to put all morphisms in degree 0). Putting it all together, we conclude that $\mathrm{CE}^*(\Lambda_Q^{n-1})$ is Koszul dual to the degree n trivial extension algebra B_Q^n of the path algebra A_Q of a Dynkin quiver of the corresponding type (see also [Li19] for another example).

As a result of these computations, for $n \geq 2$ we have a quasi-isomorphism

$$\mathrm{CE}^*(\Lambda_Q^{n-1}) \simeq \mathcal{G}_Q^n \quad (3.1)$$

over $\mathbb{C}$, which implies a quasi-equivalence

$$\mathcal{W}(\check{w}^{-1}(1)) \simeq \mathrm{perf}\Pi_n(A_Q) \quad (3.2)$$

between the wrapped Fukaya category of $\check{w}^{-1}(1)$ and the dg category of perfect modules over $\Pi_n(A_Q)$.

Remark 3.1. Note from [Sei01, Proposition 3.4] that A_Q is derived equivalent to the Fukaya–Seidel category $\mathcal{F}(\check{w})$ of the LG-model $\check{w}: \mathbb{C}^{n+1} \rightarrow \mathbb{C}$. Thus (3.2) shows that $\mathcal{W}(\check{w}^{-1}(1))$ is the Calabi–Yau completion of $\mathcal{F}(\check{w})$ for $n \geq 2$. Although this relationship between $\mathcal{F}(\check{w})$ and $\mathcal{W}(\check{w}^{-1}(1))$ is not true in general, we expect it to hold when $\check{w}$ is a double suspension of an invertible polynomial whose Milnor fiber is a log Fano manifold, since one has

$$\begin{aligned} \mathbf{w}(x_1, \dots, x_{n-1}) + x_n^2 + x_{n+1}^2 + x_0 \cdots x_{n+1} \\ = \mathbf{w}(x_1, \dots, x_{n-1}) + \left(\sqrt{1 - \frac{1}{4}(x_0 \cdots x_{n-1})^2 x_n} \right)^2 + \left(x_{n+1} + \frac{1}{2}x_0 \cdots x_n \right)^2 \end{aligned}$$

in $\mathbf{k}[[x_0, \dots, x_{n+1}]]$.

Remark 3.2. The isomorphism (3.1) remains true for $n \geq 3$ over an arbitrary commutative ring, but for $n = 2$ we have to require that 2 be invertible for type D_ℓ, E_6, E_7, E_8 , that 3 be invertible for type E_6, E_7, E_8 , and that 5 be invertible for type E_8 . Otherwise, $\mathrm{CE}^*(\Lambda_Q)$ is quasi-isomorphic to the derived multiplicative preprojective algebra (see [EL19]), which is not quasi-isomorphic to the derived (additive) preprojective algebra $\Pi_n(A_Q)$.

4. Matrix factorizations and Calabi–Yau completions

Let Γ be a subgroup of $(\mathbb{G}_m)^{n+1}$ acting diagonally on $\mathbb{A}^{n+1} := \mathrm{Spec} \mathbb{C}[x_1, \dots, x_{n+1}]$. Assume that Γ is a finite extension of the multiplicative group $\mathbb{G}_m$, so that the group $\mathrm{Char}(\Gamma) := \mathrm{Hom}(\Gamma, \mathbb{G}_m)$ of characters of Γ is an extension of a finite group by $\mathbb{Z}$. The coordinate ring $\mathbb{C}[x_1, \dots, x_{n+1}]$ has a $\mathrm{Char}(\Gamma)$ -grading coming from the Γ -action on $\mathbb{A}^{n+1}$, and we set $\chi_i := \deg x_i$ for $i \in \{1, \dots, n+1\}$. Let $\mathbf{w} \in \mathbb{C}[x_1, \dots, x_{n+1}]_\chi$ be a homogeneous element of degree $\chi \in \mathrm{Char}(\Gamma)$. Assume that $\mathbf{w}$ has an isolated critical point at the origin, so that the structure sheaf $\mathcal{O}_0$ of the origin split-generates $\mathrm{mf}(\mathbb{A}^{n+1}, \mathbf{w})$ by [KMVdB11, Proposition A.2] (see also [Orl11, Dyc11]). Let $R \subset \mathrm{Char}(\Gamma)$ be a set of representatives of the group $\mathrm{Char}(\Gamma)/(\chi)$, which we assume to be finite. Then $\mathcal{E} := \bigoplus_{\rho \in R} \mathcal{O}_0(\rho)$ split-generates $\mathrm{mf}(\mathbb{A}^{n+1}, \Gamma, \mathbf{w})$ since the autoequivalence $M \mapsto M(\chi)$ of $\mathrm{mf}(\mathbb{A}^{n+1}, \Gamma, \mathbf{w})$ shifting the Γ -weight by χ is isomorphic to the functor $M \mapsto M[2]$ shifting the cohomological grading by 2.

The n -Calabi–Yau completion of the dg Yoneda algebra $\mathcal{A} := \mathrm{hom}(\mathcal{E}, \mathcal{E})$ is given by

$$\Pi_n(\mathcal{A}) := \mathcal{A} \oplus \theta \oplus \theta \otimes_{\mathcal{A}} \theta \oplus \cdots \simeq \bigoplus_{i=0}^{\infty} \mathrm{hom}(\mathcal{E}, \theta^i(\mathcal{E})),$$

where $\theta = \Theta[n-1]$ as in Section 2, and we abuse notation and use the same symbol for an autoequivalence and its graph bimodule. Since Θ is the graph of the inverse Serre functor $\mathbb{S}^{-1}$, we have

$$\theta = \mathbb{S}^{-1}[n-1]. \quad (4.1)$$

Now, as in [LU18, Section 2], we introduce another variable x_0 of degree $\chi_0 := \chi - (\chi_1 + \cdots + \chi_{n+1})$ and consider the polynomial ring $\mathbb{C}[x_0, x_1, \dots, x_{n+1}]$ in $n+2$ variables, which naturally contains $\mathbb{C}[x_1, \dots, x_{n+1}]$ as a subring. One has

$$\mathrm{mf}(\mathbb{A}^{n+2}, \mathbf{w}) \simeq \mathrm{mf}(\mathbb{A}^1, 0) \otimes \mathrm{mf}(\mathbb{A}^{n+1}, \mathbf{w}),$$

for example, by the ungraded ($G = H = 1$) version of [BFK14, Lemma 3.52] with $v = 0$; note that $\mathrm{mf}(\mathbb{A}^1, 0)$ is obtained from $\mathrm{coh} \mathbb{A}^1$ by collapsing the cohomological grading to $\mathbb{Z}/2\mathbb{Z}$, and the tensor product of split-generators of $\mathrm{mf}(\mathbb{A}^1, 0)$ and $\mathrm{mf}(\mathbb{A}^{n+1}, \mathbf{w})$ gives a split-generator of $\mathrm{mf}(\mathbb{A}^{n+2}, \mathbf{w})$ since the critical locus of $\mathbf{w}$ as a function on $\mathbb{A}^{n+2}$ is the product of $\mathbb{A}^1$ times that as a function on $\mathbb{A}^{n+1}$.

As shown in [IT13, Theorem 2.5], whose proof carries over directly to Γ -graded cases, graded Auslander–Reiten duality [AR87] implies that

$$\mathbb{S} := (\chi_0)[n-1] \tag{4.2}$$

is a Serre functor on $\mathrm{mf}(\mathbb{A}^{n+1}, \Gamma, \mathbf{w})$. It follows from (4.1) and (4.2) that

$$\theta \simeq (-\chi_0). \tag{4.3}$$

Let $\mathcal{F}$ be the generator of $\mathrm{mf}(\mathbb{A}^{n+2}, \Gamma, \mathbf{w})$ obtained from the tensor product of the generator $\mathcal{E}$ of $\mathrm{mf}(\mathbb{A}^{n+1}, \Gamma, \mathbf{w})$ and the generator $\mathbb{C}[x_0]$ of $\mathrm{coh} \mathbb{A}^1$. If we write both of the forgetful functors $\mathrm{mf}(\mathbb{A}^{n+1}, \Gamma, \mathbf{w}) \rightarrow \mathrm{mf}(\mathbb{A}^{n+1}, \mathbf{w})$ and $\mathrm{mf}(\mathbb{A}^{n+2}, \Gamma, \mathbf{w}) \rightarrow \mathrm{mf}(\mathbb{A}^{n+2}, \mathbf{w})$ as $(\bullet)$, then one has

$$\mathrm{hom}(\overline{\mathcal{F}}, \overline{\mathcal{F}}) \simeq \mathrm{hom}(\overline{\mathcal{E}}, \overline{\mathcal{E}}) \otimes \mathbb{C}[x_0] \simeq \bigoplus_{\rho \in \mathrm{Char}(\Gamma)} \mathrm{hom}(\mathcal{E}, \mathcal{E}(\rho)) \otimes \mathbb{C}[x_0]. \tag{4.4}$$

Since $\deg(x_0) = \chi_0$, by taking the Γ -invariant part of (4.4) and using (4.3), one obtains

$$\mathrm{hom}(\mathcal{F}, \mathcal{F}) \simeq \bigoplus_{i=0}^{\infty} \mathrm{hom}(\mathcal{E}, \mathcal{E}(-i\chi_0)) \simeq \bigoplus_{i=0}^{\infty} \mathrm{hom}(\mathcal{E}, \theta^i(\mathcal{E})) \simeq \Pi_n(\mathcal{A}),$$

which shows the quasi-equivalence (1.3).

If n is greater than 1, then the degree of $x_1 \cdots x_{n+1}$ in $\mathrm{Char}(\Gamma_{\mathbf{w}}) \otimes \mathbb{Q} \cong \mathbb{Q}$ is greater than the degree of $\mathbf{w}$, which is turn is greater than the degree of any element of the Jacobi ring

$$\mathrm{Jac}_{\mathbf{w}} := \mathbb{C}[x_1, \dots, x_{n+1}] / (\partial_{x_1} \mathbf{w}, \dots, \partial_{x_{n+1}} \mathbf{w})$$

of $\mathbf{w}$, and the proof of [AGV85, Section 12.6, Theorem] shows that the polynomial $\mathbf{w} + x_0 \cdots x_{n+1}$ considered as an element of $\mathbb{C}[[x_0]][x_1, \dots, x_{n+1}]$ (that is, a formal one-parameter deformation of a formal germ of $\mathbf{w}$) is right equivalent to $\mathbf{w}$ by a formal coordinate change (that is, there exists a $\varphi \in \mathrm{Aut}_{\mathbb{C}[[x_0]]} \mathbb{C}[[x_0]][x_1, \dots, x_{n+1}]$ such that $\varphi^*(\mathbf{w} + x_0 \cdots x_{n+1}) = \mathbf{w}$). The proof moreover shows that one can choose φ to be $\Gamma_{\mathbf{w}}$ -equivariant, which implies that for any $i \in \{1, \dots, n+1\}$, the coefficient $a_{i, m_1, \dots, m_{n+1}}(x_0)$ of the expansion $\varphi^*(x_i) = \sum_{m_1, \dots, m_{n+1}=0}^{\infty} a_{i, m_1, \dots, m_{n+1}}(x_0) x_1^{m_1} \cdots x_{n+1}^{m_{n+1}}$ is a monomial in x_0 since the degree of x_0 in $\mathrm{Char}(\Gamma_{\mathbf{w}}) \otimes \mathbb{Q}$ is negative. In particular, one has $\varphi \in \mathrm{Aut}_{\mathbb{C}[[x_0]]} \mathbb{C}[[x_0]][x_1, \dots, x_{n+1}]$. This means that the formal completion of $(\mathbb{A}^{n+2}, \mathbf{w} + x_0 \cdots x_{n+1})$ along $x_1 = \cdots = x_{n+1} = 0$ is isomorphic to that of $(\mathbb{A}^{n+2}, \mathbf{w})$ as a pair of a formal scheme and a regular function on it, so that

$$\mathrm{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w} + x_0 \cdots x_{n+1}) \simeq \mathrm{mf}(\mathbb{A}^{n+2}, \Gamma_{\mathbf{w}}, \mathbf{w})$$

by [Orl11, Theorem 2.10], and the quasi-equivalence (1.2) is proved.

5. Hochschild cohomology of the derived n -preprojective algebra

5.1 Hochschild cohomology via matrix factorizations

We use the same notation as in Section 4 and set

$$V := \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \cdots \oplus \mathbb{C}x_{n+1}.$$

For $\gamma \in \Gamma$, let V_γ be the subspace of γ -invariant elements in V , let S_γ be the symmetric algebra of V_γ , let $\mathbf{w}_\gamma$ be the restriction of $\mathbf{w}$ to $\text{Spec } S_\gamma$, and let N_γ be the Γ -stable complement of V_γ in V so that $V \cong V_\gamma \oplus N_\gamma$ as a Γ -module. Then [Dyc11, CT13, Seg13, BFK14] (cf. also [LU18, Theorem 3.1]) shows that $\text{HH}^t(\text{mf}(\mathbb{A}^{n+2}, \Gamma, \mathbf{w}))$ is isomorphic to

$$\bigoplus_{\substack{\gamma \in \ker \chi, \, l \geq 0 \\ t - \dim N_\gamma = 2u}} (H^{-2l}(d\mathbf{w}_\gamma) \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{(u+l)\chi} \oplus \bigoplus_{\substack{\gamma \in \ker \chi, \, l \geq 0 \\ t - \dim N_\gamma = 2u+1}} (H^{-2l-1}(d\mathbf{w}_\gamma) \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{(u+l+1)\chi}. \quad (5.1)$$

Here $H^i(d\mathbf{w}_\gamma)$ is the i th cohomology of the Koszul complex

$$C^*(d\mathbf{w}_\gamma) := \{\cdots \rightarrow \Lambda^2 V_\gamma^\vee \otimes S_\gamma(-2\chi) \rightarrow V_\gamma^\vee \otimes S_\gamma(-\chi) \rightarrow S_\gamma\}, \quad (5.2)$$

where the rightmost term S_γ sits in cohomological degree 0 and the differential is the contraction with $d\mathbf{w}_\gamma \in (V_\gamma \otimes S_\gamma)_\chi$.

If $\mathbf{w}_\gamma$ has an isolated critical point at the origin, then the cohomology of (5.2) is concentrated in degree 0, so that only the summand

$$(\text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{u\chi}$$

with $l = 0$ in (5.1) contributes to $\text{HH}^{2u+\dim N_\gamma}$.

If V_γ contains $\mathbb{C}x_0$, then the Koszul complex $C^*(d\mathbf{w}_\gamma)$ is isomorphic to the tensor product of $C^*(d\mathbf{w}'_\gamma)$ and the complex $\{\mathbb{C}x_0^\vee \otimes \chi^\vee \otimes \mathbb{C}[x_0] \rightarrow \mathbb{C}[x_0]\}$ concentrated in cohomological degree $[-1, 0]$ with the zero differential, where $\mathbf{w}'_\gamma$ is the restriction of $\mathbf{w}$ to the complement V'_γ of $\mathbb{C}x_0$ in V_γ . If $\mathbf{w}'_\gamma$ has an isolated critical point at the origin, then $C^*(d\mathbf{w}'_\gamma)$ is quasi-isomorphic to $\text{Jac}_{\mathbf{w}'_\gamma}$ concentrated in cohomological degree 0, so that only the summands

$$(\text{Jac}_{\mathbf{w}'_\gamma} \otimes \mathbb{C}[x_0] \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{u\chi} \quad (5.3)$$

and

$$(\mathbb{C}x_0^\vee \otimes \text{Jac}_{\mathbf{w}'_\gamma} \otimes \mathbb{C}[x_0] \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)_{u\chi} \quad (5.4)$$

with $l = 0$ in (5.1) contribute to $\text{HH}^{2u+\dim N_\gamma}$ and $\text{HH}^{2u+\dim N_\gamma+1}$, respectively.

Remark 5.1. Although (5.1) may not look identical to [BFK14, Theorem 1.2], the proof in [BFK14, Section 5] actually shows (5.1). One way to think about (5.1) is the following: If we set $\mathbf{w} = 0$ and forget Γ , then the Hochschild–Kostant–Rosenberg theorem gives a quasi-isomorphism of the Hochschild cochain complex of $\text{coh } \mathbb{A}^{n+2}$ and

$$S \rightarrow V^\vee \otimes S \rightarrow \Lambda^2 V^\vee \otimes S \rightarrow \cdots \quad (5.5)$$

as complexes of $\mathbb{C}$ -vector spaces (which lifts to a quasi-isomorphism of L_∞ -algebras by the Kontsevich formality). If we introduce the potential $\mathbf{w}$, then the complex (5.5) acquires an additional differential $\Lambda^i V^\vee \otimes S \rightarrow \Lambda^{i-1} V^\vee \otimes S$ defined as the contraction with $d\mathbf{w} \in V \otimes S$, which decreases the cohomological grading by 1 so that the cohomological grading is collapsed to $\mathbb{Z}/2\mathbb{Z}$. The introduction of Γ lifts the grading to $\mathbb{Z}$ again and produces ‘twisted sectors’ from the orbifold HKR

theorem, leading to (5.1); recall the isomorphism $(\chi) \simeq [2]$ of endofunctors of $\text{mf}(\mathbb{A}^{n+2}, \Gamma, \mathbf{w})$ and the orbifold HKR theorem

$$\text{HH}^*([X/G]) = \left(\bigoplus_{g \in G} \bigoplus_{p+q=*} H^{p-\text{codim } X^g} (X^g, \Lambda^q T_{X^g} \otimes \Lambda^{\text{codim } X^g} N_{X^g/X}) \right)_G$$

for global quotients appearing, for example, in [ACH19, Corollary 1.17].

Remark 5.2. The Hochschild cohomology of a graded algebra B (with no differential) has a bi-grading such that

$$\text{HH}^{r+s}(B)^s := \text{Ext}_{B^{\text{op}} \otimes B}^r(B, B[s]) .$$

When B is the trivial extension algebra B^n of a finite-dimensional algebra A , by introducing a $\mathbb{G}_m$ -action on B^n such that A has weight 0 and $A^\vee[-n]$ has weight n , the s -grading on $\text{HH}^*(B^n)$ can be described as the weight of the induced $\mathbb{G}_m$ -action.

For any positive integer m , the underlying ungraded algebra of the trivial extension algebras B^{mn} is isomorphic to B^n , and only the cohomological gradings are different; that of the former is m times that of the latter. It follows that one has an isomorphism

$$\text{HH}^{r+ms}(B^{mn})^{ms} \cong \text{HH}^{r+s}(B^n)^s$$

of vector spaces for any positive integer m such that the parities of n and mn are the same (note that the signs in the Hochschild complex depend on the parity of the cohomological grading).

When Q is a Dynkin quiver, one can transport the $\mathbb{G}_m$ -action on B_Q^n to $\mathcal{G}_Q^n$ through the Koszul duality (2.1), so that g for $g \in Q_1$ has weight 0, while g^* for $g \in Q_1$ has weight $-n$, and h_v for $v \in Q_0$ has weight $-n$. This makes the isomorphism (2.2) $\mathbb{G}_m$ -equivariant, so that the $\mathbb{G}_m$ -weights on both sides agree.

Since $\mathbf{w}$ does not depend on x_0 , the $\mathbb{G}_m$ -action on $\mathbb{A}^{n+2}$ such that the weight of x_i is $-n$ for $i = 0$ and 0 for $i \in \{1, \dots, n+1\}$ keeps $\mathbf{w}$ invariant. This induces a $\mathbb{G}_m$ -action on $\text{mf}(\mathbb{A}^{n+2}, \Gamma, \mathbf{w})$, and hence on B_Q^n , whose weight is 0 on A_Q and n on $A_Q^\vee[-n]$ just as in [LU18]. This allows us to compute the s -grading on $\text{HH}^*(B_Q^n)$ as the $\mathbb{G}_m$ -weight on (5.1). This $\mathbb{G}_m$ -action is mirror to the one introduced in [SS12] and studied further for type A Milnor fibers in [Sei12].

5.2 Type A_ℓ

Consider the case

$$\mathbf{w} = x_1^{\ell+1} + x_2^2 + \dots + x_{n+1}^2 \in \mathbb{C}[x_0, x_1, \dots, x_{n+1}] ,$$

$$\Gamma = \Gamma_{\mathbf{w}} := \{ \gamma = (t_0, t_1, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+2} \mid t_1^{\ell+1} = t_2^2 = \dots = t_{n+1}^2 = t_0 t_1 \dots t_{n+1} \} ,$$

so that $\ker \chi \cong \mu_{\ell+1} \times (\mu_2)^n$ and $\text{Char}(\Gamma)$ is generated by χ and $\chi_i = \deg x_i$ for $i \in \{0, \dots, n+1\}$ with relations

$$\chi = (\ell+1)\chi_1 = 2\chi_2 = \dots = 2\chi_{n+1} = \chi_0 + \dots + \chi_{n+1} .$$

5.2.1 For any $\gamma \in \ker \chi$, one has

$$\text{Jac}_{\mathbf{w}_\gamma} = \begin{cases} \mathbb{C}[x_0] \otimes \mathbb{C}[x_1]/(x_1^\ell) , & \mathbb{C}x_0 \oplus \mathbb{C}x_1 \subset V_\gamma , \\ \mathbb{C}[x_0] , & \mathbb{C}x_0 \subset V_\gamma \text{ and } \mathbb{C}x_1 \not\subset V_\gamma , \\ \mathbb{C}[x_1]/(x_1^\ell) , & \mathbb{C}x_0 \not\subset V_\gamma \text{ and } \mathbb{C}x_1 \subset V_\gamma , \\ \mathbb{C} , & \text{otherwise} . \end{cases}$$

If we write an element of $\text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ as $x_0^{k_0} x_1^{k_1} \otimes x_{j_1}^\vee \wedge x_{j_2}^\vee \wedge \cdots \wedge x_{j_s}^\vee$, where $k_0 = 0$ if $\mathbb{C}x_0 \not\subset V_\gamma$ and $k_1 = 0$ if $\mathbb{C}x_1 \not\subset V_\gamma$, then its degree is given by

$$k_0\chi_0 + k_1\chi_1 - \chi_{j_1} - \cdots - \chi_{j_s},$$

which can be proportional to χ only if V_γ is either V , $\mathbb{C}x_0 \oplus \mathbb{C}x_1$, $\mathbb{C}x_0$, or 0 . We now deal with each of these cases in turn.

5.2.2 One has $V_\gamma = V$ if and only if γ is the identity element. The degree of $x_0^{k_0} x_1^{k_1} \in \text{Jac}_{\mathbf{w}}$ is

$$k_0\chi - (k_0 - k_1)\chi_1 - k_0\chi_2 - \cdots - k_0\chi_{n+1},$$

which is proportional to χ if and only if k_0 is even and $\ell + 1$ divides $k_0 - k_1$. Such an element can be written as $\mathbf{a}_{k,m} := x_0^{k+m(\ell+1)} x_1^k$, where $k \in \{0, \dots, \ell - 1\}$ and $m \in \mathbb{N}$ satisfies

- if ℓ is even, then the parities of k and m agree, and
- if ℓ is odd, then k is even.

Since

$$\begin{aligned} \deg(x_0^{k+m(\ell+1)} x_1^k) &= (k + m(\ell + 1))\chi - m\chi - \frac{1}{2}(k + m(\ell + 1))n\chi \\ &= ((k + m\ell) - \frac{1}{2}(k + m(\ell + 1))n)\chi, \end{aligned}$$

the element $x_0^{k+m(\ell+1)} x_1^k$ for such (k, m) contributes $\mathbb{C}((k + m(\ell + 1))n)$ to HH^t for $t = 2(k + m\ell) - (k + m(\ell + 1))n$ by (5.3). Similarly, for each such (k, m) , the element

$$\mathbf{a}_{k,m} := x_0^\vee \otimes x_0^{k+m(\ell+1)+1} x_1^k \in \mathbb{C}x_0^\vee \otimes \text{Jac}_{\mathbf{w}}$$

contributes $\mathbb{C}((k + m(\ell + 1))n)$ to HH^{t+1} for $t = 2(k + m\ell) - (k + m(\ell + 1))n$ by (5.4).

5.2.3 One has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1$ if and only if n is even and $\gamma = (1, 1, -1, \dots, -1)$. The degree of $x_0^{k_0} x_1^{k_1} \otimes x_2^\vee \wedge \cdots \wedge x_{n+1}^\vee \in \text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ is given by

$$k_0\chi + (k_1 - k_0)\chi_1 - (k_0 + 1)\chi_2 - \cdots - (k_0 + 1)\chi_{n+1},$$

which is proportional to χ if and only if k_0 is odd and $\ell + 1$ divides $k_1 - k_0$. Such an element can be written as

$$\mathbf{a}_{k,m} := x_0^{k+m(\ell+1)} x_1^k \otimes x_2^\vee \wedge \cdots \wedge x_{n+1}^\vee,$$

where $k \in \{0, \dots, \ell - 1\}$ and $m \in \mathbb{N}$ satisfies

- if ℓ is even, then the parities of k and m differ, and
- if ℓ is odd, then k is odd.

Since the degree of this element is $((k + m\ell) - \frac{1}{2}(k + m(\ell + 1) + 1)n)\chi$, each such (k, m) contributes $\mathbb{C}((k + m(\ell + 1))n)$ to HH^t for

$$\begin{aligned} t &= 2((k + m\ell) - \frac{1}{2}(k + m(\ell + 1) + 1)n) + \dim N_\gamma \\ &= 2(k + m\ell) - (k + m(\ell + 1))n. \end{aligned}$$

Similarly, for each such (k, m) , there is an element $\mathbf{a}_{k,m}$ contributing $\mathbb{C}((k + m(\ell + 1))n)$ to HH^{t+1} for $t = 2(k + m\ell) - (k + m(\ell + 1))n$.

5.2.4 One has $V_\gamma = \mathbb{C}x_0$ if and only if both ℓ and n are odd and $\gamma = (1, -1, \dots, -1)$. The degree of $x_0^{k_0} \otimes x_1^\vee \wedge \dots \wedge x_{n+1}^\vee \in \text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ is given by

$$k_0\chi - (k_0 + 1)\chi_1 - (k_0 + 1)\chi_2 - \dots - (k_0 + 1)\chi_{n+1},$$

which is proportional to χ if and only if $\ell + 1$ divides $k_0 + 1$. Such an element can be written as

$$\mathbf{b}_m := x_0^{m(\ell+1)-1} \otimes x_1^\vee \wedge \dots \wedge x_{n+1}^\vee$$

for $m \in \mathbb{N} \setminus \{0\}$. Since the degree of this element is $((m\ell - 1) - \frac{1}{2}m(\ell + 1)n)\chi$, each such element contributes $\mathbb{C}((m(\ell + 1) - 1)n)$ to HH^t for

$$t = 2((m\ell - 1) - \frac{1}{2}m(\ell + 1)n) + \dim N_\gamma = (2m\ell - 1) - (m(\ell + 1) - 1)n.$$

Similarly, for each $m \in \mathbb{N}$, the element

$$\beta_m := x_0^\vee \otimes x_0^{m(\ell+1)} \otimes x_1^\vee \wedge \dots \wedge x_{n+1}^\vee \in \mathbb{C}x_0^\vee \otimes \text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

contributes $\mathbb{C}(m(\ell + 1) - 1)n$ to HH^{t+1} for $t = (2m\ell - 1) - (m(\ell + 1) - 1)n$.

5.2.5 For $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$, one has $V_\gamma = 0$ if and only if $t_i \neq 1$ for all $i \in \{0, \dots, n + 1\}$. This is the case if and only if $t_2 = \dots = t_{n+1} = -1$, $t_1 \in \mu_{\ell+1} \setminus \{1\}$, and

$$t_0 = (-1)^n t_1^{-1} \neq 1. \quad (5.6)$$

If n is odd, then (5.6) holds if and only if $t_1 \neq -1$, so that the number of such γ is ℓ if ℓ is even and $\ell - 1$ if ℓ is odd. If n is even, then (5.6) always holds, and the number of such γ is ℓ . Each such γ contributes $\mathbb{C}(-n)$ to HH^n .

5.2.6 To sum up, the Hochschild cohomology group has a basis consisting of the following elements:

- $\mathbf{a}_{k,m}$ of degree $2(k + m\ell) - (k + m(\ell + 1))n$ and weight $-(k + m(\ell + 1))n$, and $\mathbf{\alpha}_{k,m}$ of degree $2(k + m\ell) - (k + m(\ell + 1))n + 1$ and weight $-(k + m(\ell + 1))n$, where
 - * if n is even, then (k, m) runs over $\{0, \dots, \ell - 1\} \times \mathbb{N}$, and
 - * if n is odd, then (k, m) runs over those pairs in $\{0, \dots, \ell - 1\} \times \mathbb{N}$ for which
 - the parities of k and m agree if ℓ is even,
 - k is even if ℓ is odd;
- if both n and ℓ are odd,
 - * $\mathbf{b}_m$ of degree $2m\ell - 1 - (m(\ell + 1) - 1)n$ and weight $-(m(\ell + 1) - 1)n$ for $m \in \mathbb{N} \setminus \{0\}$, and
 - * β_m of degree $2m\ell - (m(\ell + 1) - 1)n$ and weight $-(m(\ell + 1) - 1)n$ for $m \in \mathbb{N}$; and
- $\mathbf{s}_h$ of degree n and weight n , where h runs over
 - * $\{1, 2, \dots, \ell - 1\}$ if both ℓ and n are odd,
 - * $\{1, 2, \dots, \ell\}$ otherwise.

5.2.7 As an example, consider the case $\ell = 1$. Note that the A_1 -Milnor fiber is symplectomorphic to the cotangent bundle T^*S^n . The Hochschild cohomology group in this case is spanned by

- $\mathbf{a}_{0,m}$ for $m \in \mathbb{N}$ of degree $-2m(n - 1)$ and weight $-2mn$,
- $\mathbf{\alpha}_{0,m}$ for $m \in \mathbb{N}$ of degree $-2m(n - 1) + 1$ and weight $-2mn$,

and, if n is odd, in addition to the above,

- $\mathbf{b}_m$ for $m \in \mathbb{N} \setminus \{0\}$ of degree $-(2m-1)(n-1)$ and weight $-(2m-1)n$,
- β_m for $m \in \mathbb{N}$ of degree $-(2m-1)(n-1)+1$ and weight $-(2m-1)n$,

and, if n is even, in addition to the above,

- $\mathbf{s}_1$ of degree n and weight n .

This is consistent with the isomorphism

$$\mathrm{SH}^*(T^*S^n) \cong H_{n-*}(\mathcal{L}S^n),$$

which is a special case of the isomorphism between the symplectic cohomology of the cotangent bundle and the homology of the free loop space [Vit18, Theorem 3.1] (see, for example, [CJY04, Theorem 2] for the homology of the free loop space of spheres).

Another example is the case when $n = 2$ and ℓ is arbitrary. In this case, $\mathrm{SH}^*(\check{w}^{-1}(1))$ was computed in [EL17b] as a bigraded ring. This is compatible with the computation given here.

5.3 Type D_ℓ

The Berglund–Hübsch transform of the invertible polynomial

$$\check{\mathbf{v}} = y_1^{\ell-1} + y_1 y_2^2 + y_3^2 + \cdots + y_{n+1}^2$$

defining the D_ℓ -singularity is given by

$$\mathbf{v} = y_1^{\ell-1} y_2 + y_2^2 + \cdots + y_{n+1}^2,$$

and one has

$$\Gamma_{\mathbf{v}} = \{\gamma = (t_1, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+1} \mid t_1^{\ell-1} t_2 = t_2^2 = \cdots = t_{n+1}^2\}.$$

By completing the square and rescaling, one has $\mathbf{v}(y) = \mathbf{w}(x(y))t$, where

$$x_1 = (-1/4)^{1/(2n-2)} y_1, \quad x_2 = y_2 + \frac{1}{2} y_1^{\ell-1}, \quad x_3 = y_3, \quad \dots, \quad x_{n+1} = y_{n+1}, \quad (5.7)$$

and $\mathbf{w} = x_1^{2\ell-2} + x_2^2 + \cdots + x_{n+1}^2$. Although the change of variables (5.7) is neither linear nor diagonal, the induced action of $\Gamma_{\mathbf{v}}$ on $\mathrm{Spec} \mathbb{C}[x_1, \dots, x_{n+1}]$ remains linear and diagonal, so that one can identify $\Gamma_{\mathbf{v}}$ with a proper subgroup of $\Gamma_{\mathbf{w}}$.

Therefore, we will work with

$$\mathbf{w} = x_1^{2\ell-2} + x_2^2 + \cdots + x_{n+1}^2 \in \mathbb{C}[x_0, x_1, \dots, x_{n+1}]$$

with the non-maximal group

$$\Gamma = \{\gamma = (t_0, t_1, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+2} \mid t_1^{\ell-1} t_2 = t_2^2 = \cdots = t_{n+1}^2 = t_0 t_1 \cdots t_{n+1}\}.$$

One has $\ker \chi \cong \mu_{2\ell-2} \times (\mu_2)^{n-1}$, and $\mathrm{Char}(\Gamma)$ is generated by χ and $\chi_i = \deg x_i$ for $i \in \{0, \dots, n+1\}$ with relations

$$\chi = (\ell-1)\chi_1 + \chi_2 = 2\chi_2 = \cdots = 2\chi_{n+1} = \chi_0 + \cdots + \chi_{n+1}. \quad (5.8)$$

The relations (5.8) imply

$$\begin{aligned} \chi_2 &= \chi - (\ell-1)\chi_1, & \chi &= (2\ell-2)\chi_1, \\ \chi_0 &= \chi - \chi_1 - \cdots - \chi_n = (\ell-2)\chi_1 - \chi_3 - \cdots - \chi_{n+1}. \end{aligned}$$

5.3.1 For any $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$, one has

$$\text{Jac}_{\mathbf{w}_\gamma} = \begin{cases} \mathbb{C}[x_0] \otimes \mathbb{C}[x_1]/(x_1^{2\ell-3}), & \mathbb{C}x_0 \oplus \mathbb{C}x_1 \subset V_\gamma, \\ \mathbb{C}[x_0], & \mathbb{C}x_0 \subset V_\gamma \text{ and } \mathbb{C}x_1 \not\subset V_\gamma, \\ \mathbb{C}[x_1]/(x_1^{2\ell-3}), & \mathbb{C}x_0 \not\subset V_\gamma \text{ and } \mathbb{C}x_1 \subset V_\gamma, \\ \mathbb{C}, & \text{otherwise.} \end{cases}$$

If we write an element of $\text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ as $x_0^{k_0} x_1^{k_1} \otimes x_{j_1}^\vee \wedge x_{j_2}^\vee \wedge \dots \wedge x_{j_s}^\vee$, where $k_0 = 0$ if $\mathbb{C}x_0 \not\subset V_\gamma$ and $k_1 = 0$ if $\mathbb{C}x_1 \not\subset V_\gamma$, then its degree is given by

$$k_0 \chi_0 + k_1 \chi_1 - \chi_{j_1} - \dots - \chi_{j_s},$$

which can be proportional to χ only if $V_\gamma \cap (\mathbb{C}x_3 \oplus \dots \oplus \mathbb{C}x_{n+1})$ is either $\mathbb{C}x_3 \oplus \dots \oplus \mathbb{C}x_{n+1}$ or 0, that is,

$$t_3 = \dots = t_{n+1} = \pm 1. \quad (5.9)$$

We will assume this condition for the rest of Section 5.3.

5.3.2 One has $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$ if and only if

$$t_1^{\ell-1} t_2 = t_2^2 = \dots = t_{n+1}^2 = t_0 t_1 \dots t_{n+1} = 1.$$

If $t_0 = 1$, then one has $t_2^2 = \dots = t_{n+1}^2 = t_1 \dots t_{n+1} = 1$, so that $t_1 = (t_2 \dots t_{n+1})^{-1} = \pm 1$.

If $t_1 = 1$, then one has $t_2 = t_2^2$, so that $t_2 = 1$ and $t_3^2 = \dots = t_{n+1}^2 = t_0 t_3 \dots t_{n+1} = 1$. Under the assumption (5.9), one has $t_0 = 1$ if and only if $(t_3)^{n-1} = 1$, that is, $t_3 = 1$ or n is even.

If $t_1 = -1$, then $t_2 = (-1)^{\ell-1}$, and one has $t_0 = 1$ if and only if $(-1)^\ell t_3^{n-1} = 1$. It follows that

– V_γ contains $\mathbb{C}x_0$ if and only if

- * $\gamma = (1, \dots, 1)$, where $V_\gamma = V$;
- * $\gamma = (1, 1, 1, -1, \dots, -1)$ with odd n , where $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$;
- * $\gamma = (1, -1, -1, 1, \dots, 1)$ with even ℓ , where $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_3 \oplus \dots \oplus \mathbb{C}x_{n+1}$;
- * $\gamma = (1, -1, -1, -1, \dots, -1)$ with even ℓ and odd n , where $V_\gamma = \mathbb{C}x_0$; or
- * $\gamma = (1, -1, 1, -1, \dots, -1)$ with odd ℓ and even n , where $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_2$.

5.3.3 Note for later use that the smallest positive integer k such that the degree of x_0^k is proportional to χ is $2\ell - 2$. One has

$$\deg x_0^{2\ell-2} = (2\ell - 2)(\chi - \chi_1 - \dots - \chi_{n+1}) = ((2\ell - 3) - (\ell - 1)n)\chi.$$

5.3.4 One has $V_\gamma = V$ if and only if γ is the identity element. The degree of $x_0^{k_0} x_1^{k_1} \in \text{Jac}_{\mathbf{w}}$ is

$$k_0 \chi - (k_0 - k_1) \chi_1 - k_0 \chi_2 - \dots - k_0 \chi_{n+1},$$

which is proportional to χ if and only if k_0 is even and $2\ell - 2$ divides $k_0 - k_1$. Such an element can be written as

$$\mathbf{a}_{k,m} := x_0^{2k+(2\ell-2)m} x_1^{2k}$$

for $(k, m) \in \{0, \dots, \ell - 2\} \times \mathbb{N}$ which contributes $\mathbb{C}((2k + (2\ell - 2)m)n)$ to HH^t for $t = 4k + (4\ell - 6)m - (2k + (2\ell - 2)m)n$ since $\deg x_0^{2k} x_1^{2k} = (2k - kn)\chi$. Similarly, for each $(k, m) \in \{0, \dots, \ell - 2\} \times \mathbb{N}$, there is an element $\mathbf{a}_{k,m}$ contributing $\mathbb{C}((2k + (2\ell - 2)m)n)$ to HH^t for $t = 4k + 1 + (4\ell - 6)m - (2k + (2\ell - 2)m)n$.

5.3.5 One has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$ if and only if $\gamma = (1, 1, 1, -1, \dots, -1)$ and n is odd. The degree of $x_0^{k_0} x_1^{k_1} \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee \in \text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ is

$$k_0\chi - (k_0 - k_1)\chi_1 - k_0\chi_2 - (k_0 + 1)\chi_3 - \dots - (k_0 + 1)\chi_{n+1},$$

which is proportional to χ if and only if k_0 is odd and $2\ell - 2$ divides $k_0 - k_1 - (\ell - 1)$. Such an element can be written as

$$\mathbf{b}_{k,m} := x_0^{k+\ell-1+(2\ell-2)m} x_1^k \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee$$

for $(k, m) \in \{(k, m) \in \{0, \dots, 2\ell - 4\} \times \mathbb{Z} \mid k + \ell \text{ is even and } k + \ell - 1 + m(2\ell - 2) \geq 0\}$.

It contributes $\mathbb{C}((k + \ell - 1 + (2\ell - 2)m)n)$ to HH^t for

$$\begin{aligned} t &= 2 \deg(x_0^{k+\ell-1+(2\ell-2)m} x_1^k \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee) / \chi + \dim N_\gamma \\ &= 2k + 2\ell - 3 + (4\ell - 6)m - (k + \ell - 1 - (2\ell - 2)m)n \end{aligned}$$

since

$$\begin{aligned} \deg(x_0^{k+\ell-1} x_1^k \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee) \\ &= (k + \ell - 1)\chi - (\ell - 1)\chi_1 - (k + \ell - 1)\chi_2 - (k + \ell)\chi_3 - \dots - (k + \ell)\chi_{n+1} \\ &= (k + \ell - 1 - \tfrac{1}{2}(k + \ell)n) \chi. \end{aligned}$$

Similarly, for each

$$(k, m) \in \{(k, m) \in \{0, \dots, 2\ell - 4\} \times \mathbb{Z} \mid k + \ell \text{ is even and } k + \ell + m(2\ell - 2) \geq 0\},$$

the element

$$\beta_{k,m} := x_0^\vee \otimes x_0^{k+\ell+(2\ell-2)m} x_1^k \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee \in ((\mathbb{C}x_0)^\vee \otimes \text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee)^\Gamma$$

contributes $\mathbb{C}(k + \ell - 1 + (2\ell - 2)m)$ to HH^t for

$$t = 2k + 2\ell - 2 + (4\ell - 6)m - (k + \ell - 1 - (2\ell - 2)m)n.$$

5.3.6 One has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_3 \oplus \dots \oplus \mathbb{C}x_{n+1}$ if and only if $\gamma = (1, -1, -1, 1, \dots, 1)$ with even ℓ . An element of $\text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ whose degree is proportional to χ can be written as

$$\mathbf{c}_m := x_0^{\ell-2+(2\ell-2)m} \otimes x_1^\vee \wedge x_2^\vee$$

for $m \in \mathbb{N}$, which contributes $\mathbb{C}((\ell - 2 + (2\ell - 2)m)n)$ to HH^t for

$$\begin{aligned} t &= 2 \deg(x_0^{\ell-2} \otimes x_1^\vee \wedge x_2^\vee) / \chi + \dim N_\gamma \\ &= 2\ell - 4 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n \end{aligned}$$

since $\deg(x_0^{\ell-2} \otimes x_1^\vee \wedge x_2^\vee) = (\ell - 3 - \tfrac{1}{2}(\ell - 2)n) \chi$.

Similarly, for each $m \in \mathbb{N}$, there is an element γ_m contributing $\mathbb{C}((\ell - 2 + (2\ell - 2)m)n)$ to HH^t for $t = 2\ell - 3 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$.

5.3.7 One has $V_\gamma = \mathbb{C}x_0$ if and only if ℓ is even, n is odd, and $\gamma = (1, -1, \dots, -1) \in \ker \chi$. The degree of

$$x_0^{k_0} \otimes x_1^\vee \wedge \dots \wedge x_{n+1}^\vee \in \text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

is given by

$$k_0\chi - (k_0 + 1)\chi_1 - (k_0 + 1)\chi_2 - \dots - (k_0 + 1)\chi_{n+1},$$

which is proportional to χ if and only if $2\ell - 2$ divides $k_0 + 1$. Such an element can be written as

$$\mathbf{d}_m := x_0^{-1+m(2\ell-2)} \otimes x_1^\vee \wedge \cdots \wedge x_{n+1}^\vee$$

for $m \in \mathbb{N} \setminus \{0\}$. Since $\deg(x_0^{-1} \otimes x_1^\vee \wedge \cdots \wedge x_{n+1}^\vee) = -\chi$, each such element contributes $\mathbb{C}((-1 + (2\ell - 2)m)n)$ to HH^t for

$$\begin{aligned} t &= 2 \deg(x_0^{-1+(2\ell-2)m} \otimes x_1^\vee \wedge \cdots \wedge x_{n+1}^\vee) / \chi + \dim N_\gamma \\ &= -1 + (4\ell - 6)m - (-1 + (2\ell - 2)m)n. \end{aligned}$$

Similarly, for each $m \in \mathbb{N}$, the element

$$\delta_m := x_0^\vee \otimes x_0^{m(2\ell-2)} \otimes x_1^\vee \wedge \cdots \wedge x_{n+1}^\vee \in \mathbb{C}x_0^\vee \otimes \mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

contributes $\mathbb{C}((-1 + (2\ell - 2)m)n)$ to HH^t for $t = (4\ell - 6)m - (-1 + (2\ell - 2)m)n$.

5.3.8 One has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_2$ if and only if ℓ is odd, n is even, and $\gamma = (1, -1, 1, -1, \dots, -1)$. The degree of

$$x_0^{k_0} \otimes x_1^\vee \wedge x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee \in \mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

is given by

$$k_0\chi_0 - \chi_1 - \chi_3 - \cdots - \chi_{n+1} = (k_0(\ell - 2) - 1)\chi_1 - (k_0 + 1)\chi_3 - \cdots - (k_0 + 1)\chi_{n+1},$$

which is proportional to χ if and only if k_0 is odd and $2\ell - 2$ divides $k_0(\ell - 2) - 1$. Such an element can be written as

$$\mathbf{e}_m := x_0^{\ell-2+(2\ell-2)m} \otimes x_1^\vee \wedge x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee$$

for $m \in \mathbb{N}$, which contributes $\mathbb{C}((\ell - 2 + (2\ell - 2)m)n)$ to HH^t for

$$t = 2\ell - 4 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$$

since $\deg(x_0^{\ell-2} \otimes x_1^\vee \wedge x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee) = \frac{1}{2}(2\ell - 4 - (\ell - 1)n)\chi$.

Similarly, for each $m \in \mathbb{N}$, there is an element ϵ_m contributing $\mathbb{C}((\ell - 2 + (2\ell - 2)m)n)$ to HH^t for $t = 2\ell - 3 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$.

5.3.9 Now, we move on to the case when $\mathbb{C}x_0 \not\subset V_\gamma$. We divide it into three cases:

$$\mathbb{C}x_1 \subset V_\gamma, \quad \mathbb{C}x_1 \not\subset V_\gamma \text{ and } V_\gamma \neq 0, \quad V_\gamma = 0.$$

5.3.10 Set $\zeta := \exp(2\pi\sqrt{-1}/(2\ell - 2))$. For a given $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$, we write $t_1 = \zeta^p$ for $p \in \{0, \dots, 2\ell - 3\}$. Then one has $t_2 = (-1)^p$, so that

- V_γ contains $\mathbb{C}x_1$ if and only if $p = 0$, and
- V_γ contains $\mathbb{C}x_2$ if and only if p is even.

5.3.11 If $\mathbb{C}x_0 \not\subset V_\gamma$ and $\mathbb{C}x_1 \subset V_\gamma$, then one has that $\gamma = (-1, 1, 1, -1, \dots, -1)$, n is even, and $V_\gamma = \mathbb{C}x_1 \oplus \mathbb{C}x_2$. The element

$$x_1^{\ell-2} \otimes x_0^\vee \wedge x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee$$

has degree $(\ell - 2)\chi_1 - \chi_0 - \chi_3 - \cdots - \chi_{n+1} = 0$, so that it contributes $\mathbb{C}(-n)$ to HH^t for $t = \dim N_\gamma = n$, and this is the only contribution.

5.3.12 If $\mathbb{C}x_0 \not\subset V_\gamma$, $\mathbb{C}x_1 \not\subset V_\gamma$, and $V_\gamma \neq 0$, then V_γ is either $V_\gamma = \mathbb{C}x_2$, $\mathbb{C}x_2 \oplus \cdots \oplus \mathbb{C}x_{n+1}$ or $\mathbb{C}x_3 \oplus \cdots \oplus \mathbb{C}x_{n+1}$. No such γ contributes to HH^* since $\mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ is spanned by a single element whose degree is not proportional to χ .

5.3.13 One has $V_\gamma = 0$ if and only if $t_3 = \cdots = t_{n+1} = -1$, $t_1 = \zeta^{2m+1}$ for $m \in \{0, \dots, \ell-2\}$ and $t_0 = (-1)^n \zeta^{-2m-1} \neq 1$. The number of such γ is $\ell-2$ if ℓ is even and n is odd, and $\ell-1$ otherwise. Each such γ contributes $\mathbb{C}(-n)$ to HH^n .

5.3.14 To sum up, the Hochschild cohomology group has a basis consisting of the following elements:

- $\mathbf{a}_{k,m}$ of degree $4k + (4\ell - 6)m - (2k + (2\ell - 2)m)n$ and weight $-(2k + (2\ell - 2)m)n$ for $(k, m) \in \{0, \dots, \ell-2\} \times \mathbb{N}$,
- $\mathbf{\alpha}_{k,m}$ of degree $4k + 1 + (4\ell - 6)m - (2k + (2\ell - 2)m)n$ and weight $-(2k + (2\ell - 2)m)n$ for $(k, m) \in \{0, \dots, \ell-2\} \times \mathbb{N}$,
- if n is odd, $\mathbf{b}_{k,m}$ of degree $2k + 2\ell - 3 + (4\ell - 6)m - (k + \ell - 1 - (2\ell - 2)m)n$ and weight $-(k + \ell - 1 + (2\ell - 2)m)n$ for $\{(k, m) \in \{0, \dots, 2\ell-4\} \times \mathbb{Z} \mid k + \ell \text{ is even and } k + \ell - 1 + m(2\ell - 2) \geq 0\}$,
- if n is odd, $\mathbf{\beta}_{k,m}$ of degree $2k + 2\ell - 2 + (4\ell - 6)m - (k + \ell - 1 - (2\ell - 2)m)n$ and weight $-(k + \ell - 1 - (2\ell - 2)m)n$ for $\{(k, m) \in \{0, \dots, 2\ell-4\} \times \mathbb{Z} \mid k + \ell \text{ is even and } k + \ell + m(2\ell - 2) \geq 0\}$,
- if ℓ is even, $\mathbf{c}_m$ of degree $2\ell - 4 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$ and weight $-(\ell - 2 + (2\ell - 2)m)n$ for $m \in \mathbb{N}$,
- if ℓ is even, $\mathbf{\gamma}_m$ of degree $2\ell - 3 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$ and weight $-(\ell - 2 + (2\ell - 2)m)n$ for $m \in \mathbb{N}$,
- if ℓ is even and n is odd, $\mathbf{d}_m$ of degree $-1 + (4\ell - 6)m - (-1 + (2\ell - 2)m)n$ and weight $-(-1 + (2\ell - 2)m)n$ for $m \in \mathbb{N} \setminus \{0\}$,
- if ℓ is even and n is odd, $\mathbf{\delta}_m$ of degree $(4\ell - 6)m - (-1 + (2\ell - 2)m)n$ and weight $-(-1 + (2\ell - 2)m)n$ for $m \in \mathbb{N}$,
- if ℓ is odd and n is even, $\mathbf{e}_m$ of degree $2\ell - 4 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$ and weight $-(\ell - 2 + (2\ell - 2)m)n$ for $m \in \mathbb{N}$,
- if ℓ is odd and n is even, $\mathbf{\epsilon}_m$ of degree $2\ell - 3 + (4\ell - 6)m - (\ell - 2 + (2\ell - 2)m)n$ and weight $-(\ell - 2 + (2\ell - 2)m)n$ for $m \in \mathbb{N}$, and
- $\mathbf{s}_h$ of degree n and weight n , where h runs over a set consisting of
 - * $\ell - 2$ elements if ℓ is even and n is odd,
 - * $\ell - 1$ elements if both ℓ and n are odd, and
 - * ℓ elements otherwise.

5.4 Type E_6

Consider the case

$$\mathbf{w} = x_1^4 + x_2^3 + x_3^2 + \cdots + x_{n+1}^2 \in \mathbb{C}[x_0, x_1, \dots, x_{n+1}],$$

$$\Gamma = \Gamma_{\mathbf{w}} := \{\gamma = (t_0, t_1, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+2} \mid t_1^4 = t_2^3 = t_3^2 = \cdots = t_{n+1}^2 = t_0 t_1 \cdots t_{n+1}\},$$

so that $\ker \chi \cong \mu_4 \times \mu_3 \times (\mu_2)^{n-1}$ and $\text{Char}(\Gamma)$ is generated by χ and $\chi_i = \deg x_i$ for $i \in \{0, \dots, n+1\}$ with relations

$$\chi = 4\chi_1 = 3\chi_2 = 2\chi_3 = \dots = 2\chi_{n+1} = \chi_0 + \dots + \chi_{n+1}.$$

5.4.1 For any $\gamma \in \ker \chi$, one has

$$\text{Jac}_{\mathbf{w}_\gamma} \cong \begin{cases} \mathbb{C}[x_0], & \mathbb{C}x_0 \subset V_\gamma \\ \mathbb{C}, & \mathbb{C}x_0 \not\subset V_\gamma \end{cases} \otimes \begin{cases} \mathbb{C}[x_1]/(x_1^3), & \mathbb{C}x_1 \subset V_\gamma \\ \mathbb{C}, & \mathbb{C}x_1 \not\subset V_\gamma \end{cases} \otimes \begin{cases} \mathbb{C}[x_2]/(x_2^2), & \mathbb{C}x_2 \subset V_\gamma \\ \mathbb{C}, & \mathbb{C}x_2 \not\subset V_\gamma \end{cases}.$$

If we write an element of $\text{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ as $x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_{j_1}^\vee \wedge x_{j_2}^\vee \wedge \dots \wedge x_{j_s}^\vee$, where $k_i = 0$ if $\mathbb{C}x_i \not\subset V_\gamma$ for $i = 0, 1, 2$, then its degree is given by

$$k_0\chi_0 + k_1\chi_1 + k_2\chi_2 - \chi_{j_1} - \dots - \chi_{j_s},$$

which can be proportional to χ only if $V_\gamma \cap (\mathbb{C}x_3 \oplus \dots \oplus \mathbb{C}x_{n+1})$ is either $\mathbb{C}x_3 \oplus \dots \oplus \mathbb{C}x_{n+1}$ or 0. We will assume this condition for the rest of Section 5.4 and divide the analysis into the following three cases:

$$\mathbb{C}x_0 \subset V_\gamma, \quad \mathbb{C}x_0 \not\subset V_\gamma \text{ and } V_\gamma \neq 0, \quad V_\gamma = 0.$$

5.4.2 Since $t_0 = 1$ implies $t_2 = 1$ and $t_1 = \pm 1$, one has the following:

- V_γ contains $\mathbb{C}x_0$ if and only if
 - * $\gamma = (1, \dots, 1)$, where $V_\gamma = V$;
 - * $\gamma = (1, 1, 1, -1, \dots, -1)$ with odd $n \geq 3$, where $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$; or
 - * $\gamma = (1, -1, 1, -1, \dots, -1)$ with even n , where $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_2$.

5.4.3 One has $V_\gamma = V$ if and only if γ is the identity element. The degree of $x_0^{k_0} x_1^{k_1} x_2^{k_2} \in \text{Jac}_{\mathbf{w}}$ is

$$k_0\chi - (k_0 - k_1)\chi_1 - (k_0 - k_2)\chi_2 - k_0\chi_3 - \dots - k_0\chi_{n+1},$$

which is proportional to χ if and only if

- 4 divides $k_0 - k_1$ and 3 divides $k_0 - k_2$ if $n = 1$, and
- 4 divides $k_0 - k_1$, 3 divides $k_0 - k_2$, and k_0 be even if $n > 1$.

Thus, for $n = 1$, we must have

$$5k_0 + 3k_1 + 4k_2 = 12m \tag{5.10}$$

for $m \in \mathbb{N}$, in which case one has $\deg x_0^{k_0} x_1^{k_1} x_2^{k_2} = m\chi$.

For each $m \in \mathbb{N}$ such that $5 \nmid m$, the equation (5.10) has a unique solution with $(k_1, k_2) \in \{0, 1, 2\} \times \{0, 1\}$, and if $5 \mid m$, then there are precisely two contributions with $(k_1, k_2) = (0, 0)$ and $(k_1, k_2) = (2, 1)$ such that $(k_1, k_2, m) \in \{0, 1, 2\} \times \{0, 1\} \times \mathbb{N}$ except if $m = 0$, in which case only $(k_1, k_2) = (0, 0)$ contributes. Each such (k_1, k_2, m) contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} for $t = 2m$.

For $n > 1$, the condition that k_0 is even forces $k_1 \neq 1$, and the possible (k_0, k_1, k_2) and

$t = 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2}) / \chi$ are given by

(k_1, k_2)	k_0	t
$(0, 0)$	$12m$	$22m - 12mn$
$(0, 1)$	$4 + 12m$	$8 + 22m - (4 + 12m)n$
$(2, 0)$	$6 + 12m$	$12 + 22m - (6 + 12m)n$
$(2, 1)$	$10 + 12m$	$20 + 22m - (10 + 12m)n$

(5.11)

for $m \in \mathbb{N}$. Each (k_0, k_1, k_2) from (5.11) contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

5.4.4 One has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2 \subsetneq V$ for $\gamma \in \ker \chi$ if and only if n is an odd integer greater than or equal to 3 and $\gamma = (1, 1, 1, -1, \dots, -1)$. The degree of

$$x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee \in \mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

is given by

$$k_0 \chi - (k_0 - k_1) \chi_1 - (k_0 - k_2) \chi_2 - (k_0 + 1) \chi_3 - \cdots - (k_0 + 1) \chi_{n+1},$$

which is proportional to χ if and only if k_0 is odd, 4 divides $k_0 - k_1$, and 3 divides $k_0 - k_2$. This forces $k_1 = 1$, and the possible (k_0, k_1, k_2) and

$$t = 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee) / \chi + \dim N_\gamma$$

are given by

(k_1, k_2)	k_0	t
$(1, 0)$	$9 + 12m$	$17 + 22m - (9 + 12m)n$
$(1, 1)$	$1 + 12m$	$3 + 22m - (1 + 12m)n$

(5.12)

for $m \in \mathbb{N}$. Each (k_0, k_1, k_2) from (5.12) contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

5.4.5 One has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_2$ if and only if n is even and $\gamma = (1, -1, 1, -1, \dots, -1) \in \ker \chi$. The degree of

$$x_0^{k_0} x_2^{k_2} \otimes x_1^\vee \wedge x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee \in \mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

is given by

$$k_0 \chi - (k_0 + 1) \chi_1 - (k_0 - k_2) \chi_2 - (k_0 + 1) \chi_3 - \cdots - (k_0 + 1) \chi_{n+1},$$

which is proportional to χ if and only if 4 divides $k_0 + 1$ and 3 divides $k_0 - k_2$. The possible (k_0, k_2) and

$$t = 2 \deg(x_0^{k_0} x_2^{k_2} \otimes x_1^\vee \wedge x_3^\vee \wedge \cdots \wedge x_n^\vee) / \chi + \dim N_\gamma$$

are given by

k_2	k_0	t
0	$3 + 12m$	$6 + 22m - (3 + 12m)n$
1	$7 + 12m$	$14 + 22m - (7 + 12m)n$

(5.13)

for $m \in \mathbb{N}$. Each (k_0, k_2) from (5.13) contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

5.4.6 If $V_\gamma = \mathbb{C}x_1$, then one has

$$\deg(x_1^{k_1} \otimes x_0^\vee \wedge x_2^\vee \wedge \cdots \wedge x_{n+1}^\vee) = -\chi_0 + k_1\chi_1 - \chi_2 - \cdots - \chi_{n+1} = -\chi + (k_1 + 1)\chi_1,$$

which is not proportional to χ for any $k_1 \in \{0, 1, 2\}$. Similarly, γ with $\mathbb{C}x_0 \not\subset V_\gamma$ and $V_\gamma \neq 0$ does not contribute to HH^* .

5.4.7 One has $V_\gamma = 0$ if and only if $t_1 \in (\mu_4 \setminus \{1\})$, $t_2 \in (\mu_3 \setminus \{1\})$, and $t_3 = \cdots, t_{n+1} = -1$, since $t_2 \neq 1$ implies $t_0 = (-1)^{n-1}t_1^{-1}t_2^{-1} \neq 1$. There are six such γ , and each of them contributes $\mathbb{C}(-n)$ to HH^n .

5.5 Type E_7

Consider the case

$$\mathbf{w} = x_1^3x_2 + x_2^3 + x_3^2 + \cdots + x_{n+1}^2 \in \mathbb{C}[x_0, x_1, \dots, x_{n+1}],$$

$$\Gamma = \Gamma_{\mathbf{w}} := \{\gamma = (t_0, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+2} \mid t_1^3t_2 = t_2^3 = t_3^2 = \cdots = t_{n+1}^2 = t_0 \cdots t_{n+1}\},$$

so that $\ker \chi \cong \mu_9 \times (\mu_2)^{n-1}$ and $\mathrm{Char}(\Gamma)$ is generated by χ and $\chi_i = \deg x_i$ for $i \in \{0, \dots, n+1\}$ with relations

$$\chi = 3\chi_1 + \chi_2 = 3\chi_2 = 2\chi_3 = \cdots = 2\chi_{n+1} = \chi_0 + \cdots + \chi_{n+1}.$$

These relations imply

$$\chi_2 = \chi - 3\chi_1, \quad 9\chi_1 = 2\chi,$$

$$\chi_0 = \chi - \chi_1 - \cdots - \chi_{n+1} = 2\chi_1 - \chi_3 - \cdots - \chi_{n+1}.$$

5.5.1 For any $\gamma \in \ker \chi$, the intersection $V_\gamma \cap (\mathbb{C}x_1 \oplus \mathbb{C}x_2)$ can be $\mathbb{C}x_1 \oplus \mathbb{C}x_2$, $\mathbb{C}x_2$, or 0, where $\mathrm{Jac}_{\mathbf{w}_\gamma}$ is isomorphic to $\mathbb{C}[x_1, x_2]/(3x_1^2x_2, x_1^3 + 3x_2^2)$, $\mathbb{C}[x_2]/(3x_2^2)$, or $\mathbb{C}$, respectively. A basis of $\mathbb{C}[x_1, x_2]/(3x_1^2x_2, x_1^3 + 3x_2^2)$ is given by $\{1, x_1, x_1^2, x_1^3, x_1^4, x_2, x_1x_2\}$.

If we write an element of $\mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ as $x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_{j_1}^\vee \wedge x_{j_2}^\vee \wedge \cdots \wedge x_{j_s}^\vee$, then its degree is given by

$$k_0\chi_0 + k_1\chi_1 + k_2\chi_2 - \chi_{j_1} - \cdots - \chi_{j_s},$$

which can be proportional to χ only if $V \cap (\mathbb{C}x_3 \oplus \cdots \oplus \mathbb{C}x_{n+1})$ is either $\mathbb{C}x_3 \oplus \cdots \oplus \mathbb{C}x_{n+1}$ or 0. We assume this condition for the rest of Section 5.5.

5.5.2 For $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$, one has $t_1^2 = t_0t_3 \cdots t_{n+1} = \pm t_0$ and $t_2^2 = t_0t_1t_3 \cdots t_{n+1} = \pm t_0t_1$, so that the condition $t_0 = 1$ implies $t_1^2 = \pm 1$ and $t_2^2 = \pm t_1$, which together with $t_2^3 = 1$ imply $t_1 = t_2 = 1$. Hence, one has $\mathbb{C}x_0 \subset V_\gamma$ if and only if either $V_\gamma = V$ or $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$.

5.5.3 One has $V_\gamma = V$ if and only if γ is the identity element. The degree of $x_0^{k_0} x_1^{k_1} x_2^{k_2} \in \mathrm{Jac}_{\mathbf{w}}$ is

$$\begin{aligned} & k_0(2\chi_1 - \chi_3 - \cdots - \chi_{n+1}) + k_1\chi_1 + k_2(\chi - 3\chi_1) \\ & = k_2\chi + (2k_0 + k_1 - 3k_2)\chi_1 - k_0\chi_3 - \cdots - k_0\chi_{n+1}, \end{aligned}$$

which is proportional to χ if and only if

- 9 divides $2k_0 + k_1 - 3k_2$ if $n = 1$, and
- 9 divides $2k_0 + k_1 - 3k_2$ and k_0 is even if $n > 1$.

For $n = 1$, one has

$$t := 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2}) / \chi = 2k_2 + \frac{4}{9}(2k_0 + k_1 - 3k_2).$$

The possible (k_0, k_1, k_2) and t are given by

(k_1, k_2)	k_0	t
$(0, 0)$	$9m$	$8m$
$(1, 0)$	$4 + 9m$	$4 + 8m$
$(2, 0)$	$8 + 9m$	$8 + 8m$
$(3, 0)$	$3 + 9m$	$4 + 8m$
$(4, 0)$	$7 + 9m$	$8 + 8m$
$(0, 1)$	$6 + 9m$	$6 + 8m$
$(1, 1)$	$1 + 9m$	$2 + 8m$

(5.14)

for $m \in \mathbb{N}$. Each (k_0, k_1, k_2) from (5.14) contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

In addition, for the case $(k_1, k_2) = (2, 0)$, the element $x_0^\vee \otimes x_1^2$ corresponding to $m = -1$ in (5.14) has degree 0 and contributes $\mathbb{C}(-1)$ to HH^1 .

For $n > 1$, one has

$$t := 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2}) / \chi = 2k_2 + \frac{4}{9}(2k_0 + k_1 - 3k_2) - k_0(n - 1).$$

The possible (k_0, k_1, k_2) and t are given by

(k_1, k_2)	k_0	t
$(0, 0)$	$18m$	$34m - 18mn$
$(1, 0)$	$4 + 18m$	$8 + 34m - (4 + 18m)n$
$(2, 0)$	$8 + 18m$	$16 + 34m - (8 + 18m)n$
$(3, 0)$	$12 + 18m$	$24 + 34m - (12 + 18m)n$
$(4, 0)$	$16 + 18m$	$32 + 34m - (16 + 18m)n$
$(0, 1)$	$6 + 18m$	$12 + 34m - (6 + 18m)n$
$(1, 1)$	$10 + 18m$	$20 + 34m - (10 + 18m)n$

(5.15)

for $m \in \mathbb{N}$. Each (k_0, k_1, k_2) from (5.15) contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

5.5.4 For $n > 1$, in addition, one has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$ if and only if n is odd and $\gamma = (1, 1, 1, -1, \dots, -1)$. The degree of

$$x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee \in \mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$$

is given by

$$\begin{aligned} & k_0(2\chi_1 - \chi_3 - \cdots - \chi_{n+1}) + k_1\chi_1 + k_2(\chi - 3\chi_1) - \chi_3 - \cdots - \chi_{n+1} \\ & = k_2\chi + (2k_0 + k_1 - 3k_2)\chi_1 - (k_0 + 1)\chi_3 - \cdots - (k_0 + 1)\chi_{n+1}, \end{aligned}$$

which is proportional to χ if and only if 9 divides $2k_0 + k_1 - 3k_2$ and k_0 is odd. The possible (k_0, k_1, k_2) and

$$t := 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee) / \chi + \dim N_\gamma$$

are given by

(k_1, k_2)	k_0	t
$(0, 0)$	$9 + 18m$	$17 + 34m - (9 + 18m)n$
$(1, 0)$	$13 + 18m$	$25 + 34m - (13 + 18m)n$
$(2, 0)$	$17 + 18m$	$33 + 34m - (17 + 18m)n$
$(3, 0)$	$3 + 18m$	$7 + 34m - (3 + 18m)n$
$(4, 0)$	$7 + 18m$	$15 + 34m - (7 + 18m)n$
$(0, 1)$	$15 + 18m$	$29 + 34m - (15 + 18m)n$
$(1, 1)$	$1 + 18m$	$3 + 34m - (1 + 18m)n$

(5.16)

for $m \in \mathbb{N}$. Each (k_0, k_1, k_2) from (5.16) contributes $\mathbb{C}(k_0n)$ to HH^t and HH^{t+1} .

In addition, for the case $(k_1, k_2) = (2, 0)$, the element $x_0^\vee \otimes x_1^2 \otimes x_3^\vee \wedge \cdots \wedge x_{n+1}^\vee$ corresponding to $m = -1$ in (5.16) has degree 0 and contributes $\mathbb{C}(-n)$ to HH^n .

5.5.5 One has $V_\gamma = 0$ for $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$ if and only if $t_1 \in \mu_9 \setminus \{1\}$, $t_2 := t_1^{-3} \neq 1$, and $t_3 = \cdots = t_{n+1} = -1$, in which case one has $t_0 = (-1)^{n-1} t_1^2 \neq 1$. The set $\{t_1 \in \mu_9 \mid t_1^3 \neq 1\}$ consists of six elements, each of which contributes $\mathbb{C}(-n)$ to HH^n .

5.6 Type E_8

Consider the case

$$\mathbf{w} = x_1^5 + x_2^3 + x_3^2 + \cdots + x_{n+1}^2 \in \mathbb{C}[x_0, x_1, \dots, x_{n+1}]$$

$$\Gamma = \Gamma_{\mathbf{w}} := \{\gamma = (t_0, \dots, t_{n+1}) \in (\mathbb{G}_m)^{n+2} \mid t_1^5 = t_2^3 = t_3^2 = \cdots = t_{n+1}^2 = t_0 \cdots t_{n+1}\},$$

so that $\ker \chi \cong \mu_5 \times \mu_3 \times (\mu_2)^{n-1}$ and $\mathrm{Char}(\Gamma)$ is generated by χ and $\chi_i = \deg x_i$ for $i \in \{0, \dots, n+1\}$ with relations

$$\chi = 5\chi_1 = 3\chi_2 = 2\chi_3 = \cdots = 2\chi_{n+1} = \chi_0 + \cdots + \chi_{n+1}.$$

5.6.1 If we write an element of $\mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ as $x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_{j_1}^\vee \wedge x_{j_2}^\vee \wedge \cdots \wedge x_{j_s}^\vee$, then its degree is given by

$$k_0\chi_0 + k_1\chi_1 + k_2\chi_2 - \chi_{j_1} - \cdots - \chi_{j_s},$$

which can be proportional to χ only if $V \cap (\mathbb{C}x_3 \oplus \cdots \oplus \mathbb{C}x_{n+1})$ is either $\mathbb{C}x_3 \oplus \cdots \oplus \mathbb{C}x_{n+1}$ or 0. We assume this condition for the rest of Section 5.6.

5.6.2 Since $t_0 = 1$ implies $t_1 = t_2 = 1$, one has $\mathbb{C}x_0 \subset V_\gamma$ if and only if either $V_\gamma = V$ or $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$.

5.6.3 One has $V_\gamma = V$ if and only if γ is the identity element. The degree of $x_0^{k_0} x_1^{k_1} x_2^{k_2} \in \mathrm{Jac}_{\mathbf{w}}$ is

$$k_0\chi - (k_0 - k_1)\chi_1 - (k_0 - k_2)\chi_2 - k_0\chi_3 - \cdots - k_0\chi_{n+1},$$

which is proportional to χ if and only if

- 5 divides $k_0 - k_1$ and 3 divides $k_0 - k_2$ if $n = 1$, and
- 5 divides $k_0 - k_1$, 3 divides $k_0 - k_2$, and k_0 is even if $n > 1$.

For $n = 1$, we must have

$$7k_0 + 3k_1 + 5k_2 = 15m \quad (5.17)$$

for $m \in \mathbb{N}$, in which case one has

$$t := 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2}) / \chi = 2m.$$

For each $m \in \mathbb{N}$ such that $7 \nmid m$, the equation (5.17) has a unique solution with $(k_1, k_2) \in \{0, 1, 2, 3\} \times \{0, 1\}$, and if $7 \mid m$, then there are precisely two contributions with $(k_1, k_2) = (0, 0)$ and $(k_1, k_2) = (3, 1)$ such that $(k_1, k_2, m) \in \{0, 1, 2, 3\} \times \{0, 1\} \times \mathbb{N}$ except if $m = 0$, in which case only $(k_1, k_2) = (0, 0)$ contributes.

For $n > 1$, we must have

$$7k_0 + 3k_1 + 5k_2 = 15m \quad (5.18)$$

for $m \in \mathbb{N}$, and in addition k_0 must be in $2\mathbb{N}$. Thus, we can rewrite (5.18) as

$$k_0 = 6k_1 + 10k_2 + 30m'$$

with $m' = k_0/2 - m$. One has

$$t := 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2}) / \chi = 12k_1 + 20k_2 + 58m' - (6k_1 + 10k_2 + 30m')n.$$

Each $(k_1, k_2, m') \in \{0, 1, 2, 3\} \times \{0, 1\} \times \mathbb{N}$ contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

5.6.4 If $n > 1$, in addition, one has $V_\gamma = \mathbb{C}x_0 \oplus \mathbb{C}x_1 \oplus \mathbb{C}x_2$ if and only if n is odd and $\gamma = (1, 1, 1, -1, \dots, -1)$. The degree of $x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee \in \mathrm{Jac}_{\mathbf{w}_\gamma} \otimes \Lambda^{\dim N_\gamma} N_\gamma^\vee$ is

$$k_0\chi - (k_0 - k_1)\chi_1 - (k_0 - k_2)\chi_2 - (k_0 + 1)\chi_3 - \dots - (k_0 + 1)\chi_{n+1},$$

which is proportional to χ if and only if

$$14k_0 + 6k_1 + 10k_2 = 30m \quad (5.19)$$

for $m \in \mathbb{Z}$, and in addition we must have k_0 odd. Thus, again we can rewrite (5.19) as

$$k_0 = 15 + 6k_1 + 10k_2 + 30m',$$

where $m' = (k_0 - 1)/2 - m$. One has

$$\begin{aligned} t &:= 2 \deg(x_0^{k_0} x_1^{k_1} x_2^{k_2} \otimes x_3^\vee \wedge \dots \wedge x_{n+1}^\vee) / \chi + \dim N_\gamma \\ &= 2 \left(k_0 - \frac{1}{5}(k_0 - k_1) - \frac{1}{3}(k_0 - k_2) - \frac{1}{2}(k_0 + 1)(n - 1) \right) + (n - 1) \\ &= 29 + 12k_1 + 20k_2 + 58m' - (15 + 6k_1 + 10k_2 + 30m')n. \end{aligned}$$

Each $(k_1, k_2, m') \in \{0, 1, 2, 3\} \times \{0, 1\} \times \mathbb{Z}$ such that $15 + 6k_1 + 10k_2 + 30m' \geq 0$ contributes $\mathbb{C}(k_0 n)$ to HH^t and HH^{t+1} .

5.6.5 An element $\gamma = (t_0, \dots, t_{n+1}) \in \ker \chi$ satisfies $V_\gamma = 0$ if and only if $t_1 \in \boldsymbol{\mu}_5 \setminus \{1\}$, $t_2 \in \boldsymbol{\mu}_3 \setminus \{1\}$, $t_3 = \dots = t_{n+1} = -1$, and $t_0 = (-1)^{n-1}(t_1 t_2)^{-1}$. There are eight such elements, each of which contributes $\mathbb{C}(-n)$ to HH^n .

ACKNOWLEDGEMENTS

We thank the anonymous referees for reading the manuscript carefully and suggesting many improvements and corrections.

REFERENCES

- AAK16 M. Abouzaid, D. Auroux, and L. Katzarkov, *Lagrangian fibrations on blowups of toric varieties and mirror symmetry for hypersurfaces*, Publ. Math. Inst. Hautes Études Sci. **123** (2016), 199–282; doi:[10.1007/s10240-016-0081-9](https://doi.org/10.1007/s10240-016-0081-9).
- Abo11 M. Abouzaid, *A topological model for the Fukaya categories of plumbings*, J. Differential Geom. **87** (2011), no. 1, 1–80; doi:[10.4310/jdg/1303219772](https://doi.org/10.4310/jdg/1303219772).
- ACH19 D. Arinkin, A. Căldăraru, and M. Hablicsek, *Formality of derived intersections and the orbifold HKR isomorphism*, J. Algebra **540** (2019), 100–120; doi:[10.1016/j.jalgebra.2019.08.002](https://doi.org/10.1016/j.jalgebra.2019.08.002).
- AGV85 V.I. Arnol’d, S.M. Guseĭn-Zade, and A.N. Varchenko, *Singularities of differentiable maps, Vol. I. The classification of critical points, caustics and wave fronts*, Monogr. Math., vol. 82 (Birkhäuser Boston, Inc., Boston, MA, 1985); doi:[10.1007/978-1-4612-5154-5](https://doi.org/10.1007/978-1-4612-5154-5).
- AK19 M.A. Armenta and B. Keller, *Derived invariance of the Tamarkin–Tsygan calculus of an algebra*, C. R. Math. Acad. Sci. Paris **357** (2019), no. 3, 236–240; doi:[10.1016/j.crma.2019.01.007](https://doi.org/10.1016/j.crma.2019.01.007).
- AR87 M. Auslander and I. Reiten, *Almost split sequences for $\mathbf{Z}$ -graded rings*, Singularities, Representation of Algebras, and Vector Bundles (Lambrecht, 1985), Lecture Notes in Math., vol. 1273 (Springer, Berlin, 1987), 232–243; doi:[10.1007/BFb0078847](https://doi.org/10.1007/BFb0078847).
- Aur07 D. Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. GGT **1** (2007), 51–91.
- Aur09 ———, *Special Lagrangian fibrations, wall-crossing, and mirror symmetry*, in *Surveys in Differential Geometry, Vol. XIII, Geometry, Analysis, and Algebraic Geometry: Forty Years of the Journal of Differential Geometry*, Surv. Differ. Geom., vol. 13 (Int. Press, Somerville, MA, 2009), 1–47; doi:[10.4310/SDG.2008.v13.n1.a1](https://doi.org/10.4310/SDG.2008.v13.n1.a1).
- BEE12 F. Bourgeois, T. Ekholm, and Y. Eliashberg, *Effect of Legendrian surgery (with an appendix by Sheel Ganatra and Maksim Maydanskiy)*, Geom. Topol. **16** (2012), no. 1, 301–389; doi:[10.2140/gt.2012.16.301](https://doi.org/10.2140/gt.2012.16.301).
- BFK14 M. Ballard, D. Favero, and L. Katzarkov, *A category of kernels for equivariant factorizations and its implications for Hodge theory*, Publ. Math. Inst. Hautes Études Sci. **120** (2014), 1–111; doi:[10.1007/s10240-013-0059-9](https://doi.org/10.1007/s10240-013-0059-9).
- BH93 P. Berglund and T. Hübsch, *A generalized construction of mirror manifolds*, Nuclear Phys. B **393** (1993), no. 1–2, 377–391; doi:[10.1016/0550-3213\(93\)90250-S](https://doi.org/10.1016/0550-3213(93)90250-S).
- BH95 P. Berglund and M. Henningson, *Landau–Ginzburg orbifolds, mirror symmetry and the elliptic genus*, Nuclear Phys. B **433** (1995), no. 2, 311–332; doi:[10.1016/0550-3213\(94\)00389-V](https://doi.org/10.1016/0550-3213(94)00389-V).
- CDGG17 B. Chantraine, G. Dimitroglou Rizell, P. Ghiggini, and R. Golovko, *Geometric generation of the wrapped Fukaya category of Weinstein manifolds and sectors*, 2017, [arXiv:1712.09126](https://arxiv.org/abs/1712.09126).
- CJY04 R.L. Cohen, J.D.S. Jones, and J. Yan, *The loop homology algebra of spheres and projective spaces*, Categorical Decomposition Techniques in Algebraic Topology (Isle of Skye, 2001), Progr. Math., vol. 215 (Birkhäuser, Basel, 2004), 77–92; doi:[10.1007/978-3-0348-7863-0_5](https://doi.org/10.1007/978-3-0348-7863-0_5).
- CT13 A. Căldăraru and J. Tu, *Curved A_∞ algebras and Landau–Ginzburg models*, New York J. Math. **19** (2013), 305–342.
- Dyc11 T. Dyckerhoff, *Compact generators in categories of matrix factorizations*, Duke Math. J. **159** (2011), no. 2, 223–274; doi:[10.1215/00127094-1415869](https://doi.org/10.1215/00127094-1415869).
- EE07 P. Etingof and C.-H. Eu, *Hochschild and cyclic homology of preprojective algebras of ADE quivers*, Mosc. Math. J. **7** (2007), no. 4, 601–612; doi:[10.17323/1609-4514-2007-7-4-601-612](https://doi.org/10.17323/1609-4514-2007-7-4-601-612).
- Ekh19 T. Ekholm, *Holomorphic curves for Legendrian surgery*, 2019, [arXiv:1906.07228](https://arxiv.org/abs/1906.07228).

- EL17a T. Ekholm and Y. Lekili, *Duality between Lagrangian and Legendrian invariants*, 2017, [arXiv:1701.01284](#).
- EL17b T. Evgü and Y. Lekili, *Koszul duality patterns in Floer theory*, *Geom. Topol.* **21** (2017), no. 6, 3313–3389; [doi:10.2140/gt.2017.21.3313](#).
- EL19 ———, *Fukaya categories of plumbings and multiplicative preprojective algebras*, *Quantum Topol.* **10** (2019), no. 4, 777–813; [doi:10.4171/qt/131](#).
- EP15 A.I. Efimov and L. Positselski, *Coherent analogues of matrix factorizations and relative singularity categories*, *Algebra Number Theory* **9** (2015), no. 5, 1159–1292; [doi:10.2140/ant.2015.9.1159](#).
- ES98a K. Erdmann and N. Snashall, *On Hochschild cohomology of preprojective algebras. I*, *J. Algebra* **205** (1998), no. 2, 391–412; [doi:10.1006/jabr.1998.7547](#).
- ES98b ———, *On Hochschild cohomology of preprojective algebras. II*, *J. Algebra* **205** (1998), no. 2, 413–434; [doi:10.1006/jabr.1997.7329](#).
- ES98c ———, *Preprojective algebras of Dynkin type, periodicity and the second Hochschild cohomology*, *Algebras and Modules, II* (Geiranger, 1996), CMS Conf. Proc., vol. 24 (Amer. Math. Soc., Providence, RI, 1998), 183–193.
- Eu10 C.-H. Eu, *The calculus structure of the Hochschild homology/cohomology of preprojective algebras of Dynkin quivers*, *J. Pure Appl. Algebra* **214** (2010), no. 1, 28–46; [doi:10.1016/j.jpaa.2009.04.018](#).
- FMT05 Y. Félix, L. Menichi, and J.-C. Thomas, *Gerstenhaber duality in Hochschild cohomology*, *J. Pure Appl. Algebra* **199** (2005), no. 1–3, 43–59; [doi:10.1016/j.jpaa.2004.11.004](#).
- FU11 M. Futaki and K. Ueda, *Homological mirror symmetry for Brieskorn–Pham singularities*, *Selecta Math. (N.S.)* **17** (2011), no. 2, 435–452; [doi:10.1007/s00029-010-0055-6](#).
- FU13 ———, *Homological mirror symmetry for singularities of type D*, *Math. Z.* **273** (2013), no. 3–4, 633–652; [doi:10.1007/s00209-012-1024-x](#).
- Gan12 S. Ganatra, *Symplectic cohomology and duality for the wrapped Fukaya category*, Ph.D. Thesis, Massachusetts Institute of Technology, 2012, available at <https://dspace.mit.edu/handle/1721.1/73362>.
- Gao17 Y. Gao, *Functors of wrapped Fukaya categories from Lagrangian correspondences*, 2017, [arXiv:1712.00225](#).
- GDT90 I. M. Gel’fand, Yu. L. Daletskiĭ, and B. L. Tsygan, *On a variant of noncommutative differential geometry*, *Soviet Math. Dokl.* **40** (1990), no. 2, 422–426.
- Gin06 V. Ginzburg, *Calabi–Yau algebras*, 2006, [arXiv:math.AG/0612139](#).
- Hab20 M. Habermann, *Homological mirror symmetry for invertible polynomials in two variables*, 2020, [arXiv:2003.01106](#).
- Her19 E. Herscovich, *Hochschild (co)homology of Koszul dual pairs*, *J. Noncommut. Geom.* **13** (2019), no. 1, 59–85; [doi:10.4171/JNCG/323](#).
- HS20 M. Habermann and J. Smith, *Homological Berglund–Hübsch mirror symmetry for curve singularities*, *J. Symplectic Geom.* **18** (2020), no. 6, 1515–1574; [doi:10.4310/JSG.2020.v18.n6.a2](#).
- IT13 O. Iyama and R. Takahashi, *Tilting and cluster tilting for quotient singularities*, *Math. Ann.* **356** (2013), no. 3, 1065–1105; [doi:10.1007/s00208-012-0842-9](#).
- Kel11 B. Keller, *Deformed Calabi–Yau completions (with an appendix by Michel Van den Bergh)*, *J. reine angew. Math.* **654** (2011), 125–180; [doi:10.1515/CRELLE.2011.031](#).
- KMVdB11 B. Keller, D. Murfet, and M. Van den Bergh, *On two examples by Iyama and Yoshino*, *Compos. Math.* **147** (2011), no. 2, 591–612; [doi:10.1112/S0010437X10004902](#).
- Knö87 H. Knörrer, *Cohen–Macaulay modules on hypersurface singularities. I*, *Invent. Math.* **88** (1987), no. 1, 153–164; [doi:10.1007/BF01405095](#).

- KST07 H. Kajiura, K. Saito, and A. Takahashi, *Matrix factorization and representations of quivers. II. Type ADE case*, Adv. Math. **211** (2007), no. 1, 327–362; doi:[10.1016/j.aim.2006.08.005](https://doi.org/10.1016/j.aim.2006.08.005).
- KvK16 M. Kwon and O. van Koert, *Brieskorn manifolds in contact topology*, Bull. Lond. Math. Soc. **48** (2016), no. 2, 173–241; doi:[10.1112/blms/bdv088](https://doi.org/10.1112/blms/bdv088).
- Li19 Y. Li, *Koszul duality via suspending Lefschetz fibrations*, J. Topol. **12** (2019), no. 4, 1174–1245; doi:[10.1112/topo.12113](https://doi.org/10.1112/topo.12113).
- LU18 Y. Lekili and K. Ueda, *Homological mirror symmetry for Milnor fibers via moduli of A_∞ -structures*, 2018, [arXiv:1806.04345](https://arxiv.org/abs/1806.04345).
- Orl11 D. Orlov, *Formal completions and idempotent completions of triangulated categories of singularities*, Adv. Math. **226** (2011), no. 1, 206–217; doi:[10.1016/j.aim.2010.06.016](https://doi.org/10.1016/j.aim.2010.06.016).
- Seg13 E. Segal, *The closed state space of affine Landau–Ginzburg B-models*, J. Noncommut. Geom. **7** (2013), no. 3, 857–883; doi:[10.4171/JNCG/137](https://doi.org/10.4171/JNCG/137).
- Sei01 P. Seidel, *More about vanishing cycles and mutation*, Symplectic Geometry and Mirror Symmetry (Seoul, 2000) (World Sci. Publ., River Edge, NJ, 2001), 429–465; doi:[10.1142/9789812799821_0012](https://doi.org/10.1142/9789812799821_0012).
- Sei10 ———, *Suspending Lefschetz fibrations, with an application to local mirror symmetry*, Comm. Math. Phys. **297** (2010), no. 2, 515–528; doi:[10.1007/s00220-009-0944-8](https://doi.org/10.1007/s00220-009-0944-8).
- Sei12 ———, *Lagrangian homology spheres in (A_m) Milnor fibres via $\mathbb{C}^*$ -equivariant A_∞ -modules*, Geom. Topol. **16** (2012), no. 4, 2343–2389; doi:[10.2140/gt.2012.16.2343](https://doi.org/10.2140/gt.2012.16.2343).
- SS12 P. Seidel and J. P. Solomon, *Symplectic cohomology and q -intersection numbers*, Geom. Funct. Anal. **22** (2012), no. 2, 443–477; doi:[10.1007/s00039-012-0159-6](https://doi.org/10.1007/s00039-012-0159-6).
- Tak05 A. Takahashi, *Matrix factorizations and representations of quivers I*, 2005, [arXiv:math.AG/0506347](https://arxiv.org/abs/math/0506347).
- TT00 D. Tamarkin and B. Tsygan, *Noncommutative differential calculus, homotopy BV algebras and formality conjectures*, Methods Funct. Anal. Topology **6** (2000), no. 2, 85–100.
- Ueb16 P. Uebele, *Symplectic homology of some Brieskorn manifolds*, Math. Z. **283** (2016), no. 1–2, 243–274; doi:[10.1007/s00209-015-1596-3](https://doi.org/10.1007/s00209-015-1596-3).
- Vit18 C. Viterbo, *Functors and computations in Floer homology with applications Part II*, [arXiv:1805.01316](https://arxiv.org/abs/1805.01316).

Yankı Lekili y.lekili@imperial.ac.uk

Department of Mathematics, Imperial College London, South Kensington, London SW7 2AZ, UK

Kazushi Ueda kazushi@ms.u-tokyo.ac.jp

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan