NOTE

Synthesis and Crystal Structure of Terephthalaldehyde-*bis*-(thiosemicarbazone)-*bis*-(dimethyl sulfoxide) Solvate

WEI SU^{1,2}, PEIYUAN LI^{1,*}, LIFENG LIU², RUI CHEN^{1,3,*} and LINI HUO¹

¹College of Pharmacy, Guangxi University of Chinese Medicine, Nanning 530001, P.R. China

²College of Chemistry and Life Science, Guangxi Teachers Education University, Nanning 530001, P.R. China

³Faculty of Chinese Medicine Science Guangxi University of Chinese Medicine, Nanning 530001 P.R. China

*Corresponding authors: E-mail: lipearpear@yahoo.cn; chrui1980@163.com

(Received: 3 December 2011;

Accepted: 26 October 2012)

AJC-12340

A thiosemicarbazone single crystal synthesized with a formula as $C_{10}H_{12}N_6S_2 \cdot 2C_2H_6OS$, was synthesized by 1,4-phthalaldehyde and 4-methylthiosemicarbazide. The crystal is triclinic, space group P-1 with unit cell parameters: $a = 5.9519(5) \text{ \AA}$, $b = 7.4721(7) \text{ \AA}$, $c = 12.5450(13) \text{ \AA}$, $\alpha = 77.348(8)^\circ$, $\beta = 78.672(8)^\circ$, $\gamma = 79.193(7)^\circ$, $V = 527.64(8) \text{ \AA}^3$, $Z = 1$, $M_r = 436.63$, $D_c = 1.374 \text{ g/cm}^3$, $\mu = 0.471 \text{ mm}^{-1}$, $F(000) = 230$, $R = 0.0844$, $wR = 0.2607$ for 1902 reflections with $I > 2\sigma(I)$.

Key Words: Thiosemicarbazone, Terephthalaldehyde, Single crystal, Dimethyl sulfoxide.

Thiosemicarbazones, which possess several donor atoms and can bind to a metal atom both in neutral as well as in anionic forms and have shown a number of bonding modes, has attracted intensive attentions due to its broad spectrum of pharmacological properties, including antitumor, antifungal, antibacterial, antiviral and antimalarial activities¹⁻⁴. Herein, we report a novel terephthalaldehyde *bis*-(thiosemicarbazone)-*bis*-(dimethyl sulfoxide) solvate single crystal synthesized with a formula $C_{10}H_{12}N_6S_2 \cdot 2C_2H_6OS$.

All reagents for synthesis were commercially available and employed as received or purified by standard methods prior to use. Analyses for carbon, hydrogen and nitrogen were performed on a Perkin-Elmer 140 °C analyzer.

Synthesis: For the synthesis of present compound, 0.134 g 1,4-phthalaldehyde (1 mmol) and 0.212 g 4-methylthiosemicarbazide were diluted using 10 mL mixture solution of ethane and water (1:1). The solution was refluxed and then cooled to room temperature and filtered. The filtration residue was recrystallized using dimethyl sulfoxide/acetone (1:1). Single crystals suitable for X-ray analysis were grown from the dimethyl sulfoxide/ acetone solution by slow evaporation at room temperature in air.

Crystal structure determination: A single crystal of compound with dimensions of 0.21 mm × 0.14 mm × 0.10 mm was selected for crystallographic data collection at 293 (2) K and structure determination on a Bruker SMART CCD-4 K diffractometer employing graphite-monochromated MoK α

radiation ($\lambda = 0.71073 \text{ \AA}$). A total of 4031 reflections were collected in the range of $5.66^\circ \leq \theta \leq 50.5^\circ$, of which 1902 reflections were unique with $R_{int} = 0.0414$. The data were collected using SMART and reduced by the program SAINT. All the structures were solved by direct methods and refined by full-matrix least squares method on F_{obs}^2 by using SHELXTL-PC software package. Non-hydrogen atoms were placed in geometrically calculated positions. Hydrogen atoms were added according to theoretical model. The final full-matrix least-squares refinement including 126 variable parameters for 1902 reflections with $I > 2\sigma(I)$ and converged with unweighted and weighted agreement factors of

$$R_1 = \Sigma(|F_o| - |F_c|) / \Sigma |F_o| = 0.0844 \quad (1)$$

$$\text{and } wR_2 = \{ \Sigma [w(F_o^2 - F_c^2)^2] / \Sigma w(F_o^2)^2 \}^{1/2} = 0.2607 \quad (2)$$

where, $w = 1/[\sigma^2(F_o^2) + (0.1014P)^2]$ and $P = (F_o^2 + 2F_c^2)/3$. The maximum and minimum peaks on the final difference Fourier map are corresponding to 0.766 and -0.525 e/Å³, respectively.

The selected bond lengths and bond angles are listed in Table-1. Fig. 1 shows the molecular structure of the present compound. Fig. 2 shows the packing diagram of the present compound. The present compound crystallizes in the triclinic system of P-1 space group. The structure of the terephthalaldehyde-*bis*-(thiosemi-carbazone)-*bis*-(dimethyl sulfoxide) solvate consists of the thiosemicarbazone and dimethyl sulfoxide. The average S-C bond distance in thiosemicarbazone

TABLE-1
 SELECTED BOND DISTANCES (Å) AND ANGLES (°)

S(1A)-O(1A)	1.522(6)	S(1A)-C(6A)	1.769(9)	S(1A)-C(7A)	1.782(8)
S(2A)-C(1A)	1.700(7)	N(1A)-C(1A)	1.318(9)	N(2A)-N(3A)	1.365(8)
N(2A)-C(1A)	1.365(8)	N(3A)-C(2A)	1.278(9)	C(2A)-C(3A)	1.459(10)
C(3A)-C(4A)	1.391(10)	C(3A)-C(5N)	1.410(11)	C(4A)-C(5A)	1.379(10)
C(5A)-C(3N)	1.410(11)	O(1A)-S(1A)-C(6A)	106.2(4)	O(1A)-S(1A)-C(7A)	106.5(4)
C(6A)-S(1A)-C(7A)	97.8(4)	C(1A)-N(2A)-N(3A)	119.2(6)	C(1A)-N(3A)-N(2A)	116.6(6)
N(1A)-C(1A)-S(2A)	123.3(6)	N(1A)-C(1A)-N(2A)	117.3(6)	N(2A)-C(1A)-S(2A)	119.4(5)
N(3A)-C(2A)-C(3A)	122.1(7)	N(4A)-C(3A)-C(2A)	119.7(7)	N(4A)-C(3A)-C(5N)	119.0(7)
N(5N)-C(3A)-C(2A)	121.3(7)	N(5A)-C(4A)-C(3A)	121.3(7)	N(4A)-C(5A)-C(3N)	119.7(7)

moiety is 1.700 Å, while that in dimethyl sulfoxide is 1.769 Å. And the average C-N bond distance is 1.320 Å.

Fig. 1. Molecular structure of the terephthalaldehyde *bis*-(thiosemicarbazone)-*bis*-(dimethyl sulfoxide) solvate

Fig. 2. View of a 3D supramolecular framework of the terephthalaldehyde *bis*-(thiosemicarbazone)-*bis*-(dimethyl sulfoxide) solvate

Conclusion

Crystal structure of a novel terephthalaldehyde *bis*-(thiosemicarbazone) *bis*-(dimethyl sulfoxide) solvate has been synthesized and characterized by elemental analysis and X-ray diffraction analysis.

ACKNOWLEDGEMENTS

This research is supported by the Key Project of Chinese Ministry of Education (Grant No. 2010168), Guangxi Natural Science Foundation (Grant No. 2010GXNSFB013014), National Natural Science Foundation of China (Grant No. 21261005, 51263002) and Scientific Research Fund of Guangxi Provincial Education Department (Grant No. 201106LX273).

REFERENCES

1. T.S. Lobana, G. Bawa, R.J. Butcher and C.W. Liu, *Z. Anorg. Allg. Chem.*, **635**, 355 (2009).
2. R. Pingaew, S. Prachayasittikul and S. Ruchirawat, *Molecules*, **15**, 988 (2010).
3. T.S. Lobana, G. Bawa and R.J. Butcher, *Inorg. Chem.*, **47**, 1488 (2008).
4. K. Lee, P. Thanigaimalai, V.K. Sharma, M. Kim, E. Roh, B. Hwang, Y. Kim and S. Jung, *Bioorg. Med. Chem. Lett.*, **20**, 6794 (2010).