NOTE

Microwave Assisted Synthesis of Arylpyrazoles Using Montmorillonite K-10

FARHAD HATAMJAFARI

Department of Chemistry, Faculty of Science, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran

Corresponding author: E-mail: f_hatamjafari@tonekaboniau.ac.ir

(Received: 22 December 2011;

Accepted: 29 October 2012)

AJC-12350

Montmorillonite K-10 clay catalyzed Friedlander condensation of Baylis-Hillman adducts with phenyl hydrazine hydrochloride afforded 1,5-diarylpyrazoles.

Key Words: Arylpyrazole, Montmorillonite K-10, Microwave, Baylis-Hillman.

Heterocyclic compounds are valuable compounds and many applications have been reported. Nitrogen containing five and six membered heterocyclic compounds have been used as a scaffold to synthesize numerous therapeutic molecules¹. We have synthesized a number of heterocyclic compounds²⁻⁶. Arylpyrazole is an important class of organic compounds, which received a considerable attention due to their wide range of biological activities and widely used as pharmaceuticals, agrochemicals, antiinflammatory, antiviral, antibacterial⁷⁻¹³. In our ongoing research prompted by our interest in multiple component reactions and as part of programs in the area of heterocyclic compounds containing nitrogen¹⁴ and due to the resultant pharmacological interest in compounds which belong to the diarylpyrazoles, although this reaction done previously in other conditions^{15,16}, herein we report in a different condition using microwave irradiation, one pot reaction, a short time with high yields and easy separation of product for the construction of some 1,5-diarylpyrazole derivatives, *via* condensation of Baylis-Hillman adduct and phenyl hydrazine under microwave irradiation (**Scheme-I**).

Scheme-I

All the chemicals were obtained from Merck or Fluka. All reactions were carried out in a CEM MARS 5TM microwave oven. Silica gel SILG/UV₂₅₄ plates were used for TLC.

IR spectra were measured on a Shimadzu IR-470 Spectrophotometer. ¹H and ¹³C NMR spectra were determined on Bruker 300 DRX AVANCE instrument at 300 and 75 MHz, respectively.

Synthesis of compounds 4a-e: In a reaction a mixture of montmorillonite K10 (0.3 g) was placed in a mortar followed by a Baylis-Hillman adduct (**4a-e**) (1 mmol) and phenyl hydrazine hydrochloride (1 mmol) in 1,2-dichloroethane (5 mL). These materials were then mixed using a pestle for *ca.* 4 min. The homogenized mixture was transferred to a beaker and irradiated with microwaves for 5 min. The progress of reaction was monitored by TLC. The mixture was diluted with CH₂Cl₂ and washed with water and the organic layer was dried (MgSO₄). Evaporation of the solvent under vacuum provided a residue which was purified by column chromatography (hexan/ethyl acetate of 8/2) to afford the desired pyrazoles (**4a-e**).

Compound 4a: Orange oil; IR (KBr, $\nu_{\max}$, cm⁻¹): 3056, 2974, 1603, 1593, 1495; ¹H NMR (300 MHz, CDCl₃): 1.36 (t, *J* = 7.6 Hz, 3H), 1.94 (s, 3H), 2.75 (q, *J* = 7.6 Hz, 2H), 7.21-7.26 (m, 5H), 7.34 (dd, *J* = 7.6, 1.4 Hz, 1H), 7.55 (dt, *J* = 8.1, 1.4 Hz, 1H), 7.62 (dt, *J* = 7.5, 1.30 Hz, 1H), 7.98 (dd, *J* = 8.1, 1.2 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃): 154.4, 149.6, 140.2, 136.1, 133.5, 133.4, 130.0, 129.3, 127.2, 126.7, 125.0, 124.5, 115.3, 20.6, 13.8, 8.5 ppm.

Compound 4b: Orange oil; IR (KBr, $\nu_{\max}$, cm⁻¹): 3052, 2970, 2965, 2920, 1607, 1552, 1487, 1351, 1455, 900, 750; ¹H NMR (300 MHz, CDCl₃): 1.38 (t, *J* = 7.5 Hz, 3H), 1.95 (s, 3H), 2.7 (q, *J* = 7.5 Hz, 2H), 7.20-7.26 (m, 5H), 7.33 (dd, *J* = 7.5, 1.5 Hz, 1H), 7.58 (dt, *J* = 8.5, 1.5 Hz, 1H), 7.65 (dt, *J* = 7.5, 1.30 Hz, 1H), 8.0 (dd, *J* = 8.5, 1.3 Hz, 1H); ¹³C NMR (75

MHz, CDCl₃): 156.4, 148.2, 141.2, 135.7, 132.2, 135.5, 132.0, 129.7, 128.0, 127.4, 125.5, 124.5, 116.8, 21.8, 14.2, 8.8 ppm.

Compound 4c: Orange oil; IR (KBr, $\nu_{\max}$, cm⁻¹): 3055, 2972, 2920, 2821, 1590, 1456, 1519, 1340, 750; ¹H NMR (300 MHz, CDCl₃): 1.34 (t, J = 7.2 Hz, 3H), 2.12 (s, 3H), 2.79 (q, J = 7.2 Hz, 2H), 7.20 (dd, J = 8.5, 1.3 Hz, 2H), 7.30-7.38 (m, 3H), 7.40 (d, J = 8.5 Hz, 2H), 8.30 (d, J = 8.5 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): 158.0, 148.0, 143.5, 140.8, 135.5, 132.1, 129.2, 128.8, 127.0, 125.9, 120.2, 23.8, 14.5, 8.9 ppm.

Compound 4d: Orange oil; IR (KBr, $\nu_{\max}$, cm⁻¹): 3054, 2962, 2855, 1590, 1568, 1490, 1055, 920, 850, 747, 690; ¹H NMR (300 MHz, CDCl₃): 1.41 (t, J = 7.5 Hz, 3H), 2.4 (s, 3H), 2.9 (q, J = 7.5 Hz, 2H), 7.3 (dt, J = 7.5, 1.2 Hz, 1H), 7.25-7.35 (m, 8H); ¹³C NMR (75 MHz, CDCl₃): 158.0, 144.1, 142.2, 137.7, 138.8, 135.5, 132.1, 129.5, 128.8, 128.1, 127.1, 126.0, 115.6, 21.1, 13.5, 9.1 ppm.

Compound 4e: Orange oil; IR (KBr, $\nu_{\max}$, cm⁻¹): 3055, 2964, 2916, 2873, 1605, 1509, 1455, 1732, 763; ¹H NMR (300 MHz, CDCl₃): 1.38 (t, J = 7.5 Hz, 3H), 2.13 (s, 3H), 2.81 (q, J = 7.5 Hz, 2H), 7.14 (d, J = 7.0 Hz, 2H), 7.20-7.26 (m, 3H), 7.31 (m, 2H), 7.35 (d, J = 7.0 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): 158.5, 152.3, 145.4, 138.3, 135.6, 131.4, 128.3, 126.5, 125.5, 124.8, 118.5, 21.0, 13.1, 9.0.

Baylis-Hillman adducts were prepared by the reaction of ethyl vinyl ketone, arylaldehydes¹⁷. For synthesis of 1,5-diarylpyrazole derivatives under microwave, the reaction of Baylis-Hillman adduct (**1**), phenylhydrazine hydrochloride in 1,2-dichloroethane was used (**Scheme-I**). Therefore preparation of all the 1,5-diarylpyrazoles described in this paper, the reaction was complete within 5-8 min on solid support under microwave irradiation in excellent yields (84-90 %) to afford **4a-e** (Table-1).

TABLE-1
THREE-COMPONENT SYNTHESIS
OF SOME 1,5-DIARYLPYRAZOLES

Entry	R	Ar	Product	Yields (%)	Time (min)
1	-Me	Phenyl	4a	87	5
2	-Et	<i>o</i> -Nitrophenyl	4b	84	8
3	-Et	<i>p</i> -Nitrophenyl	4c	86	7
4	-Et	<i>m</i> -Chlorophenyl	4d	89	7
5	-Et	<i>p</i> -Chlorophenyl	4e	90	6

ACKNOWLEDGEMENTS

The author gratefully acknowledged the financial support from the Research Council of Tonekabon Branch Islamic Azad University.

REFERENCES

1. A.H. Shridhar, J. Keshavayya, H.J. Hoskeri and R.A. Shoukat Ali, *Int. Res. J. Pure Appl. Chem.*, **1**, 119 (2011).
2. J. Azizian, F. Hatamjafari, A.R. Karimi and M. Shaabanzadeh, *Synthesis*, 765 (2006).
3. J. Azizian, M. Shaabanzadeh, F. Hatamjafari and M.R. Mohammadizadeh, *Arkivoc*, 47 (2006).
4. F. Hatamjafari, *Synth. Commun.*, **36**, 3563 (2006).
5. J. Azizian, F. Hatamjafari and A.R. Karimi, *J. Heterocycl. Chem.*, **43**, 1349 (2006).
6. F. Hatamjafari and N. Montazeri, *Turk. J. Chem.*, **33**, 797 (2009).
7. F. Hatamjafari, N. Montazeri and K. Pourshamsian, *Asian J. Chem.*, **23**, 3287 (2011).
8. K.Y. Lee, G. Gowrisankar and J.N. Kim, *Tetrahedron Lett.*, **46**, 5387 (2005).
9. N. Haddad, A. Salvagano and C. Busacca, *Tetrahedron Lett.*, **45**, 5935 (2004).
10. J.W. Lyga, R.M. Patera, M.J. Plummer, B.P. Halling and D.A. Yuhas, *Pestic. Sci.*, **42**, 29 (1994).
11. S. Cacchi, G. Fabrizi and A. Grangio, *Synlett*, 959 (1997).
12. K.Y. Lee, J.M. Kim and J.N. Kim, *Tetrahedron Lett.*, **44**, 6737 (2003).
13. Y.R. Huang and J.A. Katzenellenbogen, *Org. Lett.*, **2**, 2833 (2000).
14. N.P. Peet, E.W. Huber and J.C. Huffman, *J. Heterocycl. Chem.*, **32**, 33 (1995).
15. H.E. Hassaneen, *Molecules*, **16**, 609 (2011).
16. M. Mamaghani and S. Dastmard, *Arkivoc*, 168 (2009).
17. M. Shi, J.K. Jiang and C.Q. Li, *Tetrahedron Lett.*, **43**, 127 (2002).