Synthesis and Antimicrobial Activity of Aluminium(III), Nickel(II) and Zinc(II) Schiff base Complexes Derived from *o*-Phenylenediamine and Salicylaldehyde

MOHIE E.M. ZAYED and REDA M. EL-SHISHTAWY*

Chemistry Department, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

*Corresponding author: Fax: +966 2 695 2292; Tel: +966 564666907; E-mail: elshishtawy@hotmail.com

(Received: 15 March 2012;

Accepted: 19 November 2012)

AJC-12438

N₂O₂ tetradentate Schiff base ligand was synthesized from salicylaldehyde and *o*-phenylenediamine. This ligand was reacted with Al(III), Ni(II) and Zn(II) ions to yield the corresponding complexes. The ligand and its complexes have been characterized by IR, ¹H NMR and elemental analysis. The spectral data of these compounds are discussed in connection with the structural changes due to complexation. The complexes prepared showed good antibacterial activities. The activity data show that the Al(III), Ni(II) and Zn(II) complexes are more potent antibacterial than the parent Schiff base ligand and among all Al(III) complex was the highest.

Key Words: Salicylaldehyde, Phenylenediamine, Tetradentate Schiff base, Characterization, Antimicrobial activity.

INTRODUCTION

Metals complexes using a variety of polydentate ligands are very important in several fields of science and technology¹⁻⁴. They can give a wide variety of model compounds, to mimic, simulate, or modify biological and physical properties^{5,6}. An important number of polydentate ligands that have been reported to date are Schiff bases⁷⁻¹⁰. A large number of Schiff bases and their complexes are of significant interest and attention because of their biological activity including anti-tumor, antibacterial, fungicidal and anti-carcinogenic properties^{11,12}. Tetradentate Schiff bases derived from salicylaldehyde and diamine has been the subject of many studies^{13,14}.

We report in this paper, the synthesis and antibacterial activity of *bis*-hydroxybenzylidene-1,2-phenylene diamine Schiff base and its Al(III), Ni(II) and Zn(II) complexes. The prepared ligand and complexes were characterized by elemental analysis, infrared and ¹H NMR spectra. Also, assessment for the antimicrobial activity of these compounds against *Escherichia coli*, *Staphylococcus aureus* and *Candida albicans* were reported.

EXPERIMENTAL

o-Phenylenediamine, salicylaldehyde, aluminum chloride hexahydrate, zinc nitrate and nickel acetate tetrahydrate were of the highest purity available, purchased from Sigma-Aldrich company and used as received. Solvents were of analytical grade. ¹H NMR spectra were recorded in DMSO-*d*₆ solutions on a Bruker Avance 600 MHz spectrometer. Infrared spectra were performed on a Perkin Elmer spectrum 100 FTIR

spectrometer. Elemental microanalyses were performed at the Cairo University Microanalytical Center. Melting points were determined in open capillary tubes in a Stuart Scientific melting point apparatus SMP₃ and are uncorrected.

Synthesis of N₂O₂ Schiff base (H₂L): A mixture of *o*-phenylenediamine (1.08 g, 10 mmol) and salicylaldehyde (2.44, 20 mmol) in absolute ethanol (30 mL) was refluxed for 2 h. The orange solid Schiff base was filtered off and recrystallized from dioxane. Yield 85 % (2.67 g); m.p. 162-164 °C; Anal. calcd. for C₂₀H₁₆N₂O₂ (316.36) C, 75.93; H, 5.10; N, 8.85. Found C, 76.56; H, 4.78; N, 8.96. IR (cm⁻¹): 2710 (OH), 1611 (C=N), 1585, 1561, 1480 (C=C), 1276 (C-O). ¹H NMR (δ, ppm): 12.97 (s, 2H, =Ar-OH), 8.95 (s, 2H, CH=N), 6.96-7.68 (m, 12H, Ar-H).

Synthesis of metal complexes: To a hot solution of the Schiff base (0.22 g, 0.7 mmol) in 10 mL methanol/chloroform (3:1 v/v) containing triethylamine (0.076 g, 0.75 mmol) a hot solution of metal salt (0.7 mmol) in 10 mL methanol was added. The reaction mixture was refluxed for 2 h. The mixture was concentrated and cooled. The complex that separated out was filtered, washed twice with cold ethanol and dried.

AlCl(L)·2H₂O: m.p. > 340 °C; Anal. calcd. for C₂₀H₁₈N₂O₄AlCl (412.80) C, 58.19; H, 4.39; N, 6.79. Found C, 58.74; H, 4.16; N, 6.88. IR (cm⁻¹): 2892 (OH), 1617 (C=N), 1589, 1546, 1473 (C=C), 1308 (C-O). ¹H NMR (δ, ppm): 9.34 (s, 2H, CH=N), 6.85-8.16 (m, 12H, Ar-H).

Ni(L): m.p. >340 °C; anal. calcd. for C₂₀H₁₄N₂O₂Ni (373.03) C 64.40, N, 7.51. Found C, 64.81; N, 7.66. IR (cm⁻¹):

2920 (OH), 1602 (C=N), 1575, 1512, 1438 (C=C), 1337 (C-O). ^1H NMR (δ , ppm): 8.93 (s, 2H, CH=N), 6.68-8.18 (m, 12H, Ar-H).

$\text{Zn}(\text{L})\cdot\text{H}_2\text{O}$: m.p. > 340 °C; anal. calcd. for $\text{C}_{20}\text{H}_{16}\text{N}_2\text{O}_3\text{Zn}$ (397.74) C 60.39, N, 7.04. Found C, 60.71; N, 7.20. IR (cm^{-1}): 2923 (OH), 1614 (C=N), 1587, 1529, 1459 (C=C), 1323 (C-O). ^1H NMR (δ , ppm): 9.04 (s, 2H, CH=N), 6.52-7.93 (m, 12H, Ar-H).

Antimicrobial activity: The antimicrobial activities were carried out according to the conventional agar diffusion test¹⁵ using plate-technique method. The tested microorganisms were gram positive bacteria (*Staphylococcus aureus* NRRL B-313), NRRL B-94, gram negative bacterium (*Escherichia coli* NRRL B-210) and yeast (*Candida albicans* NRRL Y-477). The used medium was nutrient agar and the incubation period of the tested microorganisms was 24 h.

RESULTS AND DISCUSSION

Synthesis of the H_2L ligand and its metal complexes are shown in **Scheme-I**. The ligand was easily obtained by condensation of 1:2 molar ratio of *o*-phenylenediamine and salicylaldehyde, respectively. This ligand is tetradentate N_2O_2 donor set and reacts by equimolar ratios with Al(III), Ni(II) and Zn(II) ions to yield stable complexes in good yield. The elemental analysis data of the H_2L and its complexes are consistent with the calculated results from the empirical formula of each compound.

Scheme-I: Synthesis of N_2O_2 Schiff base and its metal complexes

Infrared spectra: Tracing the characteristic peaks (C-O, C=N, OH) in H_2L before and after complexation would be useful to confirm the complexation. The vibrational peak of C-O at 1276 cm^{-1} present in H_2L changed upon complexation to appear at 1308 , 1337 and 1323 cm^{-1} for $\text{AlCIL}\cdot 2\text{H}_2\text{O}$, NiL and $\text{ZnL}\cdot\text{H}_2\text{O}$, respectively.

Also, the vibrational peak of C=N at 1611 cm^{-1} present in H_2L changed upon complexation to appear at 1617 , 1602 and 1614 cm^{-1} for $\text{AlCIL}\cdot 2\text{H}_2\text{O}$, NiL and $\text{ZnL}\cdot\text{H}_2\text{O}$, respectively. This

frequency shift indicates the coordination of the azomethine nitrogen and bonding with oxygen in all the complexes.

Furthermore, the IR spectra of the ligands also showed a broad and a weak band at 2710 cm^{-1} , which confirms the intramolecular hydrogen bonded OH^{13,16}. This weak peak become stronger and broader up to 3400 cm^{-1} upon complexation with Al(III) and Zn(II) indicating that water molecules are coordinated to the central metal ion as supported by the results of elemental analysis.

^1H NMR spectra: As reported in the experimental section, the chemical shift observed for the OH protons in the ligand (12.97 ppm) was not observed in any of the complexes. This confirms the bonding of oxygen to the metal ions (C-O-M). The same result was confirmed by the IR spectra. Also, the variation of the chemical shift of CH=N and aromatic protons in H_2L and its metal complexes clearly indicates that the magnetic environment has become different upon complexation and dependent on the metal type.

Antimicrobial activity: The synthesized ligand and its complexes were assessed for their antimicrobial activity. As shown in Table-1, the results revealed that complexes are more antibacterial than the ligand. On the other hand, antiyeast activity was not observed for all compounds. Al(III) and Zn(II) complexes were much more bacterial active than Ni(II) and among all Al(III) complex was the highest.

TABLE-1
ANTIMICROBIAL ACTIVITY DATA FOR THE
LIGAND AND ITS COMPLEXES

Ligand/ complex	Inhibition zone ^a		
	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Candida albicans</i>
H_2L	++	-	-
$\text{AlCIL}\cdot 2\text{H}_2\text{O}$	+++	++	-
NiL	+++	-	-
$\text{ZnL}\cdot\text{H}_2\text{O}$	++	++	-

^aInhibition zone diameter: -, no activity; +, 1–5 mm; ++, 6–9 mm; +++, more than 10 mm inhibition zone

The activities of these complexes may be explained on the basis of chelation theory; chelation reduces the polarity of the metal atom mainly because of partial sharing of its positive charge with the donor groups and possible π -electron delocalization within the whole chelate ring. Also, chelation increases the lipophilic nature of the central atom which subsequently favours its permeation through the lipid layer of the cell membrane¹⁷.

REFERENCES

- X.-G. Zhou, J.-S. Huang, Z.-Y. Zhou, K.-K. Cheung and Ch.-M. Che, *Inorg. Chim. Acta*, **331**, 194 (2002).
- D.M. Boghael and S. Mohebi, *J. Mol. Catal. A*, **179**, 41 (2002).
- A. Pui, I. Berdan, I. Morgenstern-Badarau, A. Gref and M. Peree-Fauvet, *Inorg. Chim. Acta*, **320**, 167 (2001).
- L.A. Kovbasyuk, I.O. Fritzky, V.N. Kokozyay and T.S. Iskenderov, *Polyhedron*, **16**, 1723 (1997).
- M.R. Bermejo, A. Sousa, A. García-Deibe, M. Maneiro, J. Sanmartín and M. Fondo, *Polyhedron*, **18**, 511 (1999).
- F. Tuna, L. Patron and M. Andruh, *Inorg. Chem. Commun.*, **6**, 30 (2003).
- G.B. Gill, G. Pattenden and S.J. Reynolds, *J. Chem. Soc., Perkin Trans. I*, 369 (1994); D.J. Coveney, V.F. Patel, G. Pattenden and D.M. Thompson, *J. Chem. Soc., Perkin Trans. I*, 2721 (1990).

8. R. Irie, K. Noda, Y. Ito, N. Matsumoto and T. Katsuki, *Tetrahedron: Asymm.*, **2**, 481 (1991).
9. L. Deng and E.N. Jacobsen, *J. Org. Chem.*, **57**, 4320 (1992); E.N. Jacobsen, L. Deng, Y. Furukawa and L.E. Martinez, *Tetrahedron*, **50**, 4323 (1994).
10. K. Imagawa, T. Negata, T. Yamada and T. Mukaiyama, *Chem. Lett.*, **527**, (1994); C. Bolm, *Angew. Chem.*, **103**, 414 (1991); *Angew. Chem. Ed. Engl.*, **30**, 403 (1991).
11. S. Ren, R. Wang, K. Komatsu, P. Bonaz-Krause, Y. Zyrianov, C.F. Mckenna, C. Csipke, Z.A. Tokes and E.J. Lien, *J. Med. Chem.*, **45**, 410 (2002).
12. N. Raman, A. Thangaraja and C. Kulandaisamy, *Transition Met. Chem.*, **28**, 29 (2003).
13. J.-N. Liu, B.-W. Wu, B. Zhang and Y.C. Liu, *Turk. J. Chem.*, **30**, 41 (2006).
14. A.A.A. Aziz, *J. Mol. Struct.*, **979**, 77 (2010) and references cited therein.
15. D. Greenwood, *Antimicrobial Chemotherapy, Part II. Laboratory Aspects of Antimicrobial Therapy*, Bailliere, Tindall, London, p. 71 (1983).
16. M.R. Mahmoud, S.A. Ibrahim and N.M. Ismail, *Monatsh. Chem.*, **116**, 167 (1985).
17. M.M. Abd-Elzaher, *Appl. Organometal. Chem.*, **18**, 149 (2004) and references cited therein.