Structural Identification of Chemical Constituents from *Scutellaria baicalensis* by HPLC-ESI-MS/MS and NMR Spectroscopy

WENHUI LIN¹, SHUANG LIU^{2,3} and BIN WU^{3,*}

¹School of Chemical and Environmental Engineering, Shanghai Institute of Technology, Shanghai 201418, P.R. China

²Haerbin University of Commerce, Haerbin 150076, P.R. China

³Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 501 Haik Rd, Zhangjiang, Pudong 201203, Shanghai, P.R. China

*Corresponding author: Email: wubin1031@hotmail.com; 893898379@qq.com; linwu1217@hotmail.com

(Received: 26 March 2012;

Accepted: 14 January 2013)

AJC-12697

Scutellariae Radix, the dried roots of *Scutellaria baicalensis* Georgi, which is one of the famous herb widely used in the treatment of pyrexia, jaundice and hypertension. To directly identify the compounds of the extract of the *S. baicalensis* was performed using HPLC-ESI-MS/MS. Chromatographic (analytical and preparative HPLC), chemical (hydrolysis) and spectroscopic (UV, ¹H and ¹³C NMR, MS) techniques were applied for separation, isolation, purification and identification of new compounds. A systematic phytochemical investigation of *S. baicalensis* led to the isolation of 7 new and 40 known chemical constituents. By HPLC-ESI-MS/MS and NMR spectral analyses, the new compounds were identified as 7-hydroxy-3,5,8-trimethoxy-3',4'-methylenedioxy flavone-7-O-β-glucopyranoside (1), 7-hydroxy-5,6,8-trimethoxy-3',4'-methylenedioxy flavone-7-O-β-glucopyranoside (2), patuletin-7-β-glucuronide (3), pinobanksin-6-C-glucopyranosyl-8-C-arabinopyranoside (4), chrysin-3-C-α-arabinopyranosyl-8-C-β-glucopyranoside (11), 6-sulfoxy-chrysin-7-glucoside (12) and 5-hydroxy-6,7,4'-trimethoxy-flavanone 5-sulfate (24).

Key Words: *Scutellaria baicalensis*; Structural elucidation, Chemical constituents, HPLC-MS/MS, NMR.

INTRODUCTION

Huangqin (*Radix scutellariae*) is an important traditional Chinese medicine prepared from the roots of *Scutellaria baicalensis* Georgi (Labiateae family). It is widely used in traditional Chinese medicine to treat inflammation, pyrexia, jaundice, hepatitis and hypertension¹. In the clinic, it is widely applied to cure pneumonia, hypertension, jaundice, dysentery and intestinal catarrh, pyogenic infection, etc.². It acts as a key ingredient in a number of formulae such as Shuanghuanglian oral liquid, Huangqin compound recipes granule, Yinhuang table, etc. Flavonoids and their derivatives are the main components of the *S. baicalensis*. Including baicalin, baicalein, oroxylin A, wogonoside, wogonin, apigenin and scutellarein, etc.³.

This paper describes 7 new and 38 known chemical constituents, including 2 flavone sulfates representing the first report of natural flavone sulfates from this genus and 5 novel flavone glycosides, from the aerial parts of this plant. The molecular structures of these new compounds were elucidated based on ¹D and ²D NMR spectra and comparisons with literature data.

EXPERIMENTAL

Optical rotations were measured using a Perkin-Elmer 241 MC polarimeter. IR spectra were recorded using a Perkin-Elmer 577 spectrometer. UV spectra were obtained on a Shimadzu UV-2401PC spectrophotometer. HR-TOF-MS was performed using a Waters Q-TOF Ultima. NMR experiments were run on a Varian Mercury-400 spectrometer (USA). Chemical shifts are given on a δ (ppm) scale using TMS as an internal standard. HPLC was performed using Davisil® ODS (35-60 μm, 60Å, Grace) on a BUCHI C-615 pump (UV-detector C-635 and C-660 fraction collector, BUCHI, Switzerland). Semi-preparative HPLC was carried out using EasySep TM-1010 instrument equipped with a Daisogel C₁₈ column (12 μm, 20 mm × 250 mm).

HPLC conditions: An Agilent series 1100 HPLC instrument (Waldbronn, Germany) equipped with a quaternary pump, a UV detector and a column compartment was used for analyses. The samples were separated on a Apollo C₁₈ column (5 μm, 4.6 × 250 mm, Grace), including an Easy Guard Kit C₁₈ (4 × 2 mm, Dikma) guard column. A gradient elution of 0.5 % aqueous formic acid (A) and acetonitrile (B) was used

Fig. 1. A representative HPLC-ESI-MSn analysis of the extract of *S. baicalensis*. (a) HPLC-UV chromatogram at 280 nm (b) LC-negative ion ESI-MS total ion current (TIC) profile

as follows: 20 % B in the first 10 min, 20 %-25 % B at 10-11 min, then B held at 25 % for 14 min, linearly gradient to 35 % B at 26 min and hold for 15 min, 35 % - 55 % B at 41-45 min, linearly gradient to 100 % B at 50 min and hold for 5 min. The column was maintained at 25 °C. Detection wavelengths were set at 280 nm. The flow rate was 0.8 mL/min (Fig. 1).

Mass spectrometric conditions: HPLC-MS experiments were performed with an Agilent 1100 series LC/MSD Trap mass spectrometer was connected to an Agilent 1100 HPLC instrument *via* an ESI source. The LC/MSD Trap software (5.3 version) was applied for system operation and data collection. The mass spectra were recorded in negative mode, drying gas flow rate 10 L/min, drying gas temperature 350 °C, nebulizer 35 psi., capillary voltage 3500 V, mass range 50-1500 *m/z*.

The roots of *Scutellaria baicalensis* Georgi were collected from Xi'an, ShanXi province, China and authenticated by Professor Shen Jingui of Shanghai Institute of Materia Medica, Chinese Academy of Sciences. A voucher specimen (SIMM-HCG-HQ0905) was deposited in the Herbarium of Shanghai Institute of Materia Medica, Chinese Academy of Sciences.

Extraction and isolation: The roots of *S. baicalensis* (5 Kg) were extracted twice with 95 % ethanol at room temperature. The EtOH extracts were combined and evaporated under reduced pressure to give a residue (677 g). A portion of

the extract (384 g) was suspended in distilled water and chromatographed using medium-pressure ODS column chromatography (column dimensions 100 mm × 460 mm) eluting followed by increasing concentrations of MeOH in H₂O (10, 20, 30, 50, 80 and 100 % MeOH; fraction volume 1200 mL). Give 48 fractions (355 g), which were combined into seven main groups (A-G). The fraction A (33.8 g) eluted with 10 % MeOH was rechromatographed using semi-preparative HPLC to give compound **1** (5.4 mg) and compound **2** (6.0 mg). Fraction B (26.4 g) eluted with 13 % MeOH was repeatedly chromatographed using semi-preparative HPLC to yield compound **3** (4.5 mg) and compound **4** (5.4 mg). The fraction C (11.4 g) eluted with 15 % MeOH was rechromatographed using semi-preparative HPLC to yield compound **11** (15.7 mg). The fraction D (93.9 g) eluted with 17 % MeOH was rechromatographed using semi-preparative HPLC to give compound **12** (21.1mg). The fraction F (73.9 g) eluted with 22 % MeOH was rechromatographed using semi-preparative HPLC to give compound **24** (6.8 mg).

Experimental data for the new flavonoid glycosides

7-Hydroxy-3,5,8-trimethoxy-3',4'-methylenedioxy flavone-7-O- β -glucopyranoside (1**):** Pale yellow needles crystals; m.p.: 284-285 °C; UV (MeOH) λ_{max} (log ϵ): 256 (4.21), 273 (4.23), 350 (4.23); IR (KBr, ν_{max} , cm^{-1}): 3100 (OH), 1620 (C=O), 1605, 1590, 1495; ESI-MS [*M*-H]⁻*m/z*, 533; HR-

ESI-MS [M-H]⁻ *m/z*, 533.1274 (calcd. for C₂₅H₂₆O₁₃, 533.1295); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.69 (1H, dd, *J* = 8.5, 2.0 Hz, H-6'), 7.54 (1H, d, *J* = 2.0 Hz, H-2'), 7.13 (1H, d, *J* = 8.5 Hz, H-5'), 6.41 (1H, s, H-6), 6.13 (2H, s, -O-CH₂-O-), 5.11 (1H, d, *J* = 7.1 Hz, Glc-1), 3.90 (1H, m, Glc-6), 3.81 (3H, s, 5-OCH₃), 3.75 (3H, s, 8-OCH₃), 3.72 (3H, s, 3-OCH₃), 3.71 (1H, m, Glc-6), 3.46 (1H, m, Glc-5), 3.38 (1H, m, Glc-3), 3.35 (1H, m, Glc-4), 3.21 (1H, m, Glc-2); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 174.8 (C-4), 157.1 (C-5), 155.4 (C-7), 152.0 (C-2), 151.3 (C-3'), 149.2 (C-9), 147.5 (C-4'), 140.9 (C-3), 131.0 (C-8), 124.8 (C-1'), 119.0 (C-6'), 109.3 (C-5'), 109.0 (C-10), 108.1 (C-2'), 103.1 (-O-CH₂-O-), 99.5 (Glc-1), 94.2 (C-6), 77.0 (Glc-5), 76.3 (Glc-3), 73.1 (Glc-2), 69.6 (Glc-4), 61.2 (8-OCH₃), 60.7 (Glc-6), 59.7 (3-OCH₃), 56.4 (5-OCH₃); ESI-MS/MS data (Table-1).

7-Hydroxy-5,6,8-trimethoxy-3',4'-methylenedioxy flavone-7-O-β-glucopyranoside (2): Pale yellow needles; UV (MeOH) λ_{max} (log ε): 257, 350 nm. IR (KBr, ν_{max}, cm⁻¹): 3100 (OH), 1620 (C=O), 1605, 1590, 1495; ESI-MS [M-H]⁻ *m/z*, 533; HR-ESI-MS [M-H]⁻ *m/z*, 533.1275 (calcd. for C₂₅H₂₆O₁₃, 533.1295); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.69 (1H, dd, *J* = 8.8, 2.3 Hz, H-6'), 7.55 (1H, d, *J* = 2.3 Hz, H-2'), 7.14 (1H, d, *J* = 8.8 Hz, H-5'), 6.63 (1H, s, H-3), 6.11 (2H, s, -O-CH₂-O-), 5.08 (1H, d, *J* = 7.3 Hz, Glc-1), 3.92 (3H, s, 5-OCH₃), 3.90 (1H, m, Glc-6), 3.89 (3H, s, 8-OCH₃), 3.85 (3H, s, 6-OCH₃), 3.71 (1H, m, Glc-6), 3.46 (1H, m, Glc-5), 3.39 (1H, m, Glc-3), 3.35 (1H, m, Glc-4), 3.21 (1H, m, Glc-2); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 175.1 (C-4), 160.6 (C-2), 151.5 (C-3'), 150.3 (C-6), 149.1 (C-8), 147.8 (C-4'), 147.6 (C-9), 144.0 (C-5), 137.1 (C-7), 124.3 (C-1'), 119.3 (C-6'), 110.1 (C-5'), 109.5 (C-10), 109.0 (C-2'), 106.3 (C-3), 103.2 (-O-CH₂-O-), 99.9 (Glc-1), 77.1 (Glc-5), 76.4 (Glc-3), 73.1 (Glc-2), 69.5 (Glc-4), 61.5 (5-OCH₃), 61.2 (6-OCH₃), 61.0 (8-OCH₃, Glc-6); ESI-MS/MS data (Table-1).

Patuletin-7-β-glucuronide (3): Pale yellow crystals; UV (MeOH) λ_{max} (log ε): 255, 267sh, 336 nm; IR (KBr, ν_{max}, cm⁻¹): 3327, 1654, 1610, 1504, 1457, 881, 805; ESI-MS [M-H]⁻ *m/z*, 507; HR-ESI-MS [M-H]⁻ *m/z*, 507.0760 (calcd. for C₂₂H₂₀O₁₄, 507.0775); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 12.31 (1H, s, 5-OH), 9.92 (1H, s, 3'-OH), 9.80 (1H, s, 3-OH), 9.40 (4'-OH), 7.67 (1H, d, *J* = 2.1 Hz, H-2'), 7.56 (1H, dd, *J* = 7.8, 2.1 Hz, H-6'), 6.89 (1H, d, *J* = 7.8 Hz, H-5'), 6.51 (1H, s, H-8), 5.17 (1H, d, *J* = 7.4 Hz, GlcA-1), 4.15 (1H, d, *J* = 9.5 Hz, GlcA-5), 3.81 (3H, s, 6-OCH₃), 3.67 (1H, t, *J* = 9.5 Hz, GlcA-4), 3.59 (2H, m, GlcA-2, GlcA-3); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 176.3 (C-4), 169.9 (GlcA-6), 157.3 (C-7), 151.5 (C-9), 151.4 (C-5), 147.5 (C-4'), 146.5 (C-2), 145.3 (C-3'), 135.1 (C-3), 130.9 (C-6), 122.1 (C-1'), 119.8 (C-6'), 115.5 (C-5'), 114.7 (C-2'), 103.8 (C-10), 101.2 (GlcA-1), 94.5 (C-8), 75.8 (GlcA-5), 75.3 (GlcA-3), 72.8 (GlcA-2), 71.2 (GlcA-4), 59.9 (6-OCH₃); ESI-MS/MS data (Table-1).

Pinobankasin-6-C-glucopyranosyl-8-C-arabinopyranoside (4): Colourless needles; m.p.: 164-166 °C; [α]_D²⁵ = 10.1 (c 0.5, MeOH). UV (MeOH) λ_{max} (log ε): 380, 315 nm; IR (KBr, ν_{max}, cm⁻¹): 3460, 3120, 1650, 1620, 1480, 1285, 1178; ESI-MS [M-H]⁻ *m/z*, 565; HR-ESI-MS [M-H]⁻ *m/z*, 565.1563 (calcd. for C₂₆H₃₀O₁₄, 565.1558); ¹H NMR (CD₃OD, 400 MHz) δ 7.49 (2H, m, H-2', H-6'), 7.27 (1H, m, H-4'), 7.25

(2H, m, H-3', H-5'), 5.07 (1H, d, *J* = 12.8 Hz, H-3), 4.51 (1H, d, *J* = 12.8 Hz, H-2), 4.80 (1H, d, *J* = 9.6 Hz, Glc-1), 4.64 (1H, d, *J* = 9.0 Hz, Ara-1), 4.01 (1H, m, Ara-2), 3.93 (1H, m, Ara-5), 3.89 (1H, m, Glc-2), 3.88 (1H, m, Ara-4), 3.70 (1H, m, Glc-6), 3.69 (1H, m, Ara-5), 3.54 (1H, m, Glc-6), 3.53 (1H, m, Ara-3), 3.28 (3H, m, Glc-3, Glc-4, Glc-5); ¹³C NMR (CD₃OD, 100 MHz) δ 195.6 (C-4), 165.1 (C-7), 161.5 (C-5), 160.5 (C-9), 136.3 (C-1'), 129.2 (C-3', C-5'), 128.7 (C-2', C-6'), 127.5 (C-4'), 106.9 (C-6), 105.4 (C-8), 96.8 (C-10), 83.6 (C-2), 77.7 (Glc-3, Glc-5), 75.2 (Ara-3), 74.2 (Ara-1), 73.2 (Glc-1), 72.7 (C-3), 71.1 (Glc-4), 70.8 (Glc-2), 69.6 (Ara-5), 69.1 (Ara-2), 68.6 (Ara-4), 62.2 (Glc-6); ESI-MS/MS data (Table-1).

Chrysin 3-C-a-arabinopyranosyl-8-C-β-glucopyranoside (11): Pale yellow needles crystals; m.p.: 274-276 °C; UV (MeOH) λ_{max} (log ε): 269, 316 nm; IR (KBr, ν_{max}, cm⁻¹): 3264 (-OH), 1650 (-C=O), 1595, 1342, 1216, 845; ESI-MS [M-H]⁻ *m/z*, 547; HR-ESI-MS [M-H]⁻ *m/z*, 547.1435 (calcd. for C₂₆H₂₈O₁₃, 547.1452); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 12.90 (1H, s, 5-OH), 10.51 (1H, s, 7-OH), 8.13 (2H, m, H-2', H-6'), 7.70 (1H, m, H-4'), 7.66 (2H, m, H-3', H-5'), 6.27 (1H, s, H-6), 4.80 (1H, d, *J* = 9.4 Hz, Ara-1), 4.76 (1H, d, *J* = 9.8 Hz, Glc-1), 4.00 (2H, m, Glc-2, Ara-2), 3.79 (1H, m, Ara-5), 3.77 (1H, m, Ara-4), 3.74 (1H, m, Glc-6), 3.57 (1H, m, Ara-5), 3.52 (1H, m, Glc-6), 3.42 (1H, m, Ara-3), 3.36 (1H, m, Glc-4), 3.30 (1H, m, Glc-3), 3.27 (1H, m, Glc-5); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 178.3 (C-4), 161.0 (C-5, C-7), 160.1 (C-2), 154.3 (C-9), 131.9 (C-4'), 129.3 (C-1'), 128.7 (C-3', C-5'), 126.1 (C-2', C-6'), 101.3 (C-10), 115.3 (C-3), 105.1 (C-8), 95.8 (C-6), 81.5 (Glc-5), 79.1 (Glc-3), 74.6 (Ara-1), 74.5 (Glc-1), 73.9 (Ara-3), 71.5 (Glc-2), 70.8 (Glc-4), 70.1 (Ara-5), 68.9 (Ara-2), 68.4 (Ara-4), 60.9 (Glc-6); ESI-MS/MS data (Table-1).

6-Sulfoxy-chrysin-7-glucoside (12): Pale yellow needles crystals; UV λ_{max} (log ε): 267, 313nm; IR (KBr, ν_{max}, cm⁻¹): 3348, 1655, 1617, 1510, 1068, 854, 768, 686; ESI-MS [M-H]⁻ *m/z*, 511; HR-ESI-MS [M-H]⁻ *m/z*, 511.0562 (calcd. for C₂₁H₂₀O₁₃S, 511.0547); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 12.63 (1H, s, 5-OH), 8.08 (2H, m, H-2', H-6'), 7.58 (3H, m, H-3', H-4', H-5'), 7.36 (1H, s, H-8), 7.11 (1H, s, H-3), 4.53 (1H, d, *J* = 6.8 Hz, Glc-1), 3.92 (1H, m, Glc-6), 3.71 (1H, m, Glc-6), 3.45 (1H, m, Glc-5), 3.37 (1H, m, Glc-3), 3.35 (1H, m, Glc-4), 3.20 (1H, m, Glc-2); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ 182.6 (C-4), 166.5 (C-7), 164.5 (C-2), 164.2 (C-5), 153.2 (C-9), 132.5 (C-4'), 130.4 (C-1'), 129.3 (C-3', C-5'), 126.7 (C-2', C-6'), 105.7 (C-3, C-10), 101.3 (Glc-1), 94.7 (C-6), 94.4 (C-8), 77.0 (Glc-5), 76.2 (Glc-3), 73.5 (Glc-2), 70.0 (Glc-4), 61.0 (Glc-6); ESI-MS/MS data (Table-1).

5-Hydroxy-6,7,4'-trimethoxy flavanone 5-sulfate (24): Pale yellow crystals; m.p.: 164-166 °C; UV (MeOH) λ_{max} (log ε): 225, 287, 340nm; IR (KBr, ν_{max}, cm⁻¹): 3400, 2905, 1630, 1565, 1510, 1445, 1270, 1250, 1190, 1155, 1105, 1005, 965, 890 and 830; ESI-MS [M-H]⁻ *m/z*, 409; HR-ESI-MS [M-H]⁻ *m/z*, 409.0590 (calcd. for C₁₈H₁₈O₉S, 409.0594); ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.48 (2H, d, *J* = 8.8 Hz, H-2', H-6'), 6.95 (2H, d, *J* = 8.8 Hz, H-3', H-5'), 6.71 (1H, s, H-8), 5.57 (1H, dd, *J* = 8.8, 3.8 Hz, H-2), 3.85 (3H, s, 7-OCH₃), 3.80 (3H, s, 6-OCH₃), 3.78 (3H, s, 4'-OCH₃), 3.35 (1H, dd, *J* =

TABLE-1
 CHARACTERIZATION OF COMPOUNDS BY HPLC-ESI-MS/MS FROM *S. baicalensis*

No.	R _t (min)	UV λ _{max} (nm)	ESI-MS/MS m/z (abundance)	Identification results
1	3.3	256,273,350	MS ² [533]: 516(1.2), 317(8.4), 353(27.3), 341(7.8), 329(0.6), 313(2.3)	7-Hydroxy-3,5,8-trimethoxy-3',4'-(methylenedioxy) flavone-7-O-β-glucopyranoside
2	3.4	258,350	MS ² [533]: 516(1.2), 317(8.4), 353(27.3), 341(7.8), 313(2.3)	7-Hydroxy-5,6,8-trimethoxy-3',4'-(methylenedioxy) flavone-7-O-β-glucopyranoside
3	4.8	255,267,336	MS ² [507]: 491(26.5), 399(99.3), 331(31.1)	Patuletin-7-β-glucuronide
4	4.9	280,315	MS ² [565]: 547(100.0), 487(50.7), 457(75.9), 445 (43.9), 367(86.0), 337(51.5)	Pinobankasin-6-C-glucopyranosyl-8-C-arabinopyranoside
5	5.2	255,370	MS ² [657]: 495(100.0), 333(0.2)	6,8-Dihydroxy quercetin diglucopyranoside
6	5.5	282,330	MS ² [671]: 509(100.0), 389(4.4), 347(50.4), 228(1.4)	3,5,7,8,3',4'-Hexahydroxy-6-methoxy flavone diglucopyranoside
7	5.7	271,338	MS ² [563]: 443(100.0), 545(15.0), 503(27.0), 486 (14.6), 473(67.1), 383(33.9), 353(73.1)	Apigenin-6-C-β-D-glucoside-8-C-?-L-arabinoside
8	5.9	271,338	MS ² [563]: 443(100.0), 545(28.5), 503(44.4), 485 (15.6), 473(67.7), 425(26.3), 383 (38.7), 353(34.8)	Apigenin-6-C-α-L-arabinoside-8-C-β-D-glucoside
9	6.4	222,388	MS ² [577]: 457(100.0), 559(8.2), 517(1.3), 487(45.4), 367(21.1), 337(39.8)	Chrysin-6,8-C-diglucoside
10	6.7	224,272,316	MS ² [547]: 487(100.0), 529(14.5), 457(79.3), 427 (79.3), 367(41.0), 337(85.9)	Chrysin-6-C-α-L-arabinoside-8-C-β-D-glucoside
11	7.1	269,316	MS ² [547]: 457(100.0), 529(22.4), 487(55.3), 427 (58.9), 367(48.2), 337(50.3)	Chrysin-3-C-α-arabinopyranosyl-8-C-β-glucopyranoside
12	8.6	267,313	MS ² [511]: 431(100.0), 349(10.6), 269(3.2), 241(1.2)	6-Sulfooxy-chrysin 7-glucoside
13	10.1	220,274,316	MS ² [547]: 457(100.0), 529(10.4), 487(38.1), 427 (22.6), 367(34.6)	Chrysin-6-C-β-D-glucoside-8-C-αL-arabinoside
14	10.6	279,317	MS ² [511]: 269(100.0), 431(20.5), 351(4.2), 349(3.1), 241(8.6)	Bicaicin-7-O-β-D-glucopyranosyl sulfate conjugate
15	11.6	252,347	MS ² [475]: 299(100.0), 300(26.2), 284(5.4), 175(3.7)	Chrysoeriol-7-O-β-D-glucuronide
16	13.6	276,315	MS ² [431]: 269(100.0), 271(3.3), 270(16.5), 201(1.2), 196(1.0)	Baicaicin-7-O-β-D-glucopyranoside
17	14.0	272,316	MS ² [445]: 430(100.0), 283(87.6), 268(82.4), 161 (1.2)	Wogonin-5-O-β-D-glucopyranoside
18	14.7	270,313	MS ² [415]: 295(100.0), 325(17.2), 296(14.2), 267 (3.2)	Chrysin-8-C-β-D-glucopyranoside
19	16.4	276,315	MS ² [445]: 269(100.0), 366(2.5), 341(2.2), 270(19.5), 223(3.3), 175(20.7)	Baicaicin-7-O-glucuronide (baicalin)
20	16.6	263, 323	MS ² [475]: 299(100.0), 301(3.2), 300(19.4), 285(2.2), 284(37.1), 175(5.2)	5,7,6'-trihydroxy-2'-methoxy flavone-7-O-β-D-glucuronide
21	17.4	282,333	MS ² [445]: 269(100.0), 427(2.0), 385(1.0), 175(14.9), 137(2.9)	Apigenin-7-O-β-D-glucuronopyranoside (scutellarin A)
22	17.7	268,336	MS ² [445]: 269(100.0), 244(35.0), 216(1.4), 175(2.6)	2'-Hydroxychrysin-7-glucuronide
23	18.1	254,346	MS ² [463]: 161(100.0), 301(68.4), 283(3.4), 191(8.1), 162(12.7), 139(5.5)	Quercetin-7-O-β-D-glucopyranoside (quercimeritrin)
24	18.4	287,340	MS ² [409]: 329(100.0), 315(0.2), 314(0.9), 299(0.2)	5-Hydroxy-6,7,4'-trimethoxy flavanone-5-sulfate
25	18.9	272,312	MS ² [459]: 283(100.0), 268(8.6), 175(20.7)	Oroxylin A-7-O-β-D-glucuronide
26	19.3	274,349(sh)	MS ² [459]: 283(100.0), 269(5.0), 268(4.2), 176(1.3), 175(24.5)	Wogonin-7-O-β-D-glucuronopyranoside
27	21.5	276,320	MS ² [349]: 269(100.0), 332(0.1), 270(23.0), 225(0.3)	Baicaicin-7-O-sulfate
28	23.1	288	MS ² [303]: 125(100.0), 285(3.1), 275(9.0), 259(8.7), 217(25.3), 177(30.9)	3,5,7,3',4'-pentahydroxy flavanone (taxifolin)
29	24.4	324,287	MS ² [303]: 125(100.0), 285(6.1), 259(15.6), 177 (12.9)	(2R,3R)-3,5,7,2',6'-pentahydroxy flavanone
30	26.4	290,330	MS ² [303]: 125(100.0), 285(4.4), 275(10.2), 259(6.1), 217(13.8), 177(27.6)	3,5,7,2',6'-Pentahydroxy flavanone
31	27.6	254,351	MS ² [301]: 161(100.0), 283(4.8), 273(3.8), 270(1.2), 139(44.0)	3,5,7,3',4'-Pentahydroxy flavone (quercetin)
32	28.1	253,302	MS ² [301]: 161(100.0), 283(7.5), 273(17.9), 269 (18.2), 139(86.4), 133(15.3)	5,7,2',6'-Tetrahydroxy flavonol (viscidulin I)
33	31.1	267,342	MS ² [345]: 330(100.0), 331(17.5), 329(4.9), 328(1.4), 315(5.1)	5,7,2',5'-Tetrahydroxy-8,6'-dimethoxy flavone (viscidulin III)
34	31.5	261,310	MS ² [285]: 125(100.0), 269(5.6), 241(1.0), 177(1.0), 161(28.7), 151(2.6),	5,7,2',6'-Tetrahydroxy flavone
35	34.2	275,340	MS ² [393]: 313(100.0), 331(0.9), 314(8.1), 298(0.7)	5,2'-Dihydroxy-7,8-dimethoxy flavone (panicolin)
36	35.2	272,313	MS ² [283]: 269(100.0), 175(4.1)	5,7-Dihydroxy-6-methoxy flavone (oroxylin A)
37	36.3	262,346	MS ² [285]: 241(11.0), 237(7.6), 227(9.9), 151(10.0)	3,5,7,4'-Tetrahydroxy flavone (kaempferol)
38	36.8	241,253,349	MS ² [285]: 241(6.8), 233(5.8), 231(2.4), 201(6.7), 199(1.2), 175(4.9), 151(13.7)	5,7,3',4'-Tetrahydroxy flavone (luteolin)
39	37.9	222,278,304	MS ² [285]: 267(10.0), 243(2.5), 223(3.5), 218(4.1), 201(9.1), 153(5.3), 133(9.1)	5,7,8,4'-tetrahydroxy flavone (isoscuteallarein)
40	48.5	276,320	MS ² [285]: 268(18.5), 241(7.2), 213(18.7), 205(3.4), 149(1.3)	5,6,7,4'-Tetrahydroxy flavone (scuteallarein)
41	49.6	260(sh),274,362	MS ² [331]: 331(100.0), 300(1.2), 273(10.7), 157(1.8)	5,7,3',4',5'-Pentahydroxy-3,6,8-trimethoxy flavone
42	50.4	274	MS ² [283]: 268(100.0), 283(4.2), 269(16.2), 137(5.4)	5,7-Dihydroxy-8-methoxy flavone (wogonin)
43	51.0	230,253,311(sh)	MS ² [253]: 181(100.0), 208(9.4), 164(20.9), 125 (47.7)	7,4'-Dihydroxy flavone
44	51.1	268,315	MS ² [253]: 153(100.0), 186(4.5), 171(3.9), 145(13.7)	5,7-Dihydroxy flavone (chrysin)
45	51.2	269,324	MS ² [283]: 267(100.0), 136(1.8)	5,7-Dihydroxy-4'-methoxy flavone (acacetin)

Fig. 2. Structures of the new components in the extract of *S. baicalensis*

12.0, 8.8 Hz, H-3), 2.81 (1H, dd, $J = 12.0, 3.8$ Hz, H-3); ^{13}C NMR (DMSO- d_6 , 100 MHz) δ 197.5 (C-4), 159.7 (C-4'), 158.6 (C-7), 155.7 (C-5), 153.5 (C-9), 133.6 (C-6), 130.3 (C-1'), 128.4 (C-2', C-6'), 113.8 (C-3', C-5'), 106.2 (C-10), 90.5 (C-8), 78.9 (C-2), 59.9 (7-OCH₃), 58.2 (6-OCH₃), 55.1 (4'-OCH₃), 42.7 (C-3); ESI-MS/MS data (Table-1).

Supporting information: NMR spectra (^1H NMR, ^{13}C NMR, HMQC and HMBC), HR-ESI-MS of all new compounds and detailed extraction and isolation procedures are given as supporting information.

RESULTS AND DISCUSSION

Systematic phytochemical investigations were carried out to probe the chemical profile of the *S. baicalensis*, the successive purification of the EtOH extract of *S. baicalensis* by medium-pressure ODS column chromatography. Seven new and 38 known compounds were isolated and purified through medium-pressure ODS column, as well as repeated preparative using semi-preparative HPLC.

Compound **1**, obtained as pale yellow needles crystals. The molecular formula of **1** was determined to be C₂₅H₂₆O₁₃ on the basis of negative ion ESI-MS, ^1H and ^{13}C NMR spectral data. In the UV spectrum of **1**, the maximum bands are at 256, 273 and 350 nm. Its IR spectrum showed absorption bands due to hydroxy (3100 cm⁻¹), carbonyl (1620 cm⁻¹), aromatic (1605, 1590, 1495 cm⁻¹) groups. In the ^1H NMR spectrum (Table-1), three singlets (3H, each) were observed at $\delta_{\text{H}} = 3.72$, 3.75 and 3.81, indicating three methoxyl groups. The ^{13}C NMR signals of **1** were assigned with the help of an HMQC experiment, establishing direct C-H bonding. The location of the three methoxyl groups were assigned to C-3, C-5 and C-8 based on the observed HMBC correlations from 3-OCH₃ (δ 3.72), 5-OCH₃ (δ 3.81) and 8-OCH₃ (δ 3.75) to C-3 (δ 140.9), C-5 (δ 157.1) and C-8 (δ 131.0), respectively. This experiment also clarified the site of glycosidation showing a long-range correlation between the anomeric proton of glucose at $\delta_{\text{H}} 5.11$ (d, $J = 7.7$ Hz, H-1") and the oxygenated carbon atom at $\delta_{\text{C}} 155.4$ (C-7). The ^1H and ^{13}C data of the aglycone were similar to those reported for 7-hydroxy-3,5,8-trimethoxy-3',4'-methylenedioxyflavone⁴. Finally, the structure of **1** was established as 7-hydroxy-3,5,8-trimethoxy-3',4'-methylenedioxy flavone-7-O- β -glucopyranoside.

Compound **2**, obtained as pale yellow needles with the molecular formula C₂₅H₂₆O₁₃ (negative ion ESI-MS, ^1H and ^{13}C NMR spectral data). The ^1H and ^{13}C NMR spectra of **2** were similar to those of **1**, which are different with site of methoxyl linkage. Comparison with the spectrum of compound **1** allowed an upfield shift for C-3 signal and a downfield shift for the C-6 signal, furthermore, signals of methoxyl groups could be assigned to C-6. The structure of aglycone was identified by comparison of its MS and ^1H and ^{13}C NMR data with those reported in the literature⁵. Based on these findings, the chemical structure of **2** has been elucidated to be 7-hydroxy-5,6,8-trimethoxy-3',4'-methylenedioxy flavone-7-O- β -glucopyranoside.

Compound **3**, the ^{13}C NMR spectrum of **3** displayed the presence of one glucuronide unit in addition to 16 carbon signals for the aglycone. The chemical shifts of the aglycone carbons were similar to those reported for patuletin⁶. In the HMBC spectrum, long-range correlations were observed from δ 5.17 (H-1 of GlcA) to δ 157.3 (C-7 of the aglycone). Accordingly, the structure of **3** was formulated as patuletin 7- β -glucuronide.

Compound **4** is optically active, $[\alpha]_{\text{D}}^{25} = 10.1$ (c 0.5, MeOH) and its molecular formula was established as C₂₆H₃₀O₁₄ by high resolution ESIMS. Additionally, two anomeric proton resonances were observed at δ 4.64 (d, $J = 9.0$ Hz) and 4.80 (d, $J = 9.6$ Hz), indicating its diglycosidic structure. The sugar functionality was identified as a β -glucopyranose and arabinopyranoside by ^1H and ^{13}C NMR spectral data together with ESI-MS. The ^{13}C NMR spectrum of **4** exhibited 26 carbon resonances: 8 quaternary carbons (C), 16 methine (CH) and 2 methylene (CH₂). In the HMBC spectrum, the proton signal at δ 4.51 (d, $J = 12.8$ Hz, H-2) showed long-range correlations with the carbon signals at δ 160.5 (C-9), whereas the anomeric proton (δ 4.64) showed long-range correlation with C-8 (δ 105.4) and C-9 (δ 160.5), indicating that the arabinose moiety was located at C-8. In the HMBC spectrum, the long-range correlation between the anomeric proton and C-3" (δ 77.7) of the glucose moiety, correlation to C-5 (δ 161.5)/C-6 (δ 106.9)/C-7 (δ 165.1), confirmed the attachment of glucose moiety to the carbon C-6. The structures of aglycone were identified by comparison of its MS and ^1H and ^{13}C NMR data with those reported in the literature⁷. Therefore, the structure

of **4** can be identified as pinobankasin-6-C-glucopyranosyl-8-C-arabinopyranoside.

Compound **11**, the ^1H and ^{13}C NMR spectral data together with ESI-MS supported the presence of diglycosidic. The structure of the sugar moieties could be entirely established on the basis of ^1H and ^{13}C NMR data. In HMBC, the anomeric proton (δ 4.64) showed long-range correlation with C-8 (δ 105.1), C-9 (δ 154.3) and C-3" (δ 79.1) indicating that the glucose moiety was located at C-8. Moreover, this experiment also clarified the site of glycosidation showing a long-range correlation between the anomeric proton of arabinose at δ_{H} 4.80 (d, J = 9.4 Hz) and the carbon atom at δ_{C} 178.3 (C-4). Thus, compound **11** was identified as chrysin 3-C- α -arabinopyranosyl-8-C- β -glucopyranoside by comparing its ^1H and ^{13}C NMR data with previously published data⁸.

Compound **12**, obtained as a yellow amorphous powder. The molecular formula was determined to be $\text{C}_{21}\text{H}_{20}\text{O}_{13}\text{S}$ on the basis of negative ion ESI-MS, ^1H and ^{13}C NMR spectral data. The UV spectrum exhibited maxima at 267 and 313 nm and IR spectrum suggested the presence of hydroxy groups (3348 cm^{-1}). The ^1H and ^{13}C data of the aglycone were similar to those reported for chrysin. The significant shifts of the carbon signals of the *ipso*, *ortho* and *para* positions caused by the sulphation at C-6 are consistent with literature data for flavonoid sulphates⁹. The β -glucopyranosyl unit is linked to C-7 of the aglycone from the chemical shift of this carbon at δ 166.5. Therefore, the structures of compound **12**, 6-sulfoxy-chrysin-7-glucoside, were identified on the basis of comparison of its spectroscopic (NMR, MS) data in comparison with literature values¹⁰.

Compound **24** was isolated as a pale yellow crystal. The molecular formula was established as $\text{C}_{18}\text{H}_{18}\text{O}_9\text{S}$ by ESI-MS, ^1H and ^{13}C NMR spectral data. The IR spectrum showed absorption bands at 1630 (C=O), 1445 (aromatic), 1270 (C-O-) and the UV spectrum exhibited maxima at 225, 287 and 340 nm. The signal at δ 6.71 related to two methylene protons showed long-range correlations with the signals for C-7, C-9, C-6 and C-10, whereas the methoxy protons at δ 3.80 and δ 3.85 showed long-range correlation with C-5 and C-6, respectively. Additional, the signal at δ_{H} 3.78 showed long-range correlations with the signals for C-3', C-5' and 4'-OCH₃ in the B ring. The significant shifts of the carbon signals of the *ipso*, *ortho* and *para* positions caused by the sulphation at C-5. Therefore, indicating the sulphation at C-5. Based on these findings, compound **25** were identified by comparing their physical and spectral data with the literature values: 5-hydroxy-6,7,4'-trimethoxy flavanone 5-sulfate¹¹.

By comparing their physical and spectroscopic data with published data, the structures of the other 38 known chemical constituents were elucidated as 6,8-dihydroxy quercetin diglucopyranoside (**5**)¹², 3,5,7,8,3',4'-hexahydroxy-6-methoxy flavone diglucopyranoside (**6**)¹³, apigenin-6-C- β -D-glucoside-8-C- α -L-arabinoside (**7**)¹⁴, apigenin-6-C- α -L-arabinoside-8-C- β -D-glucoside (**8**)¹⁴, chrysin-6,8-C-diglucoside (**9**)¹⁵, chrysin-6-C- α -L-arabinoside-8-C- β -D-glucoside (**10**)¹⁶, chrysin-6-C- β -D-glucoside-8-C- α -L-arabinoside (**13**)¹⁶, baicalein-7-O- β -D-glucopyranosyl sulfate conjugate (**14**)¹⁷, chrysoeriol-7-O- β -D-glucuronide (**15**)¹⁸, baicalein-7-O- β -D-

glucopyranoside (**16**)¹⁷, wogonin-5-O- β -D-glucopyranoside (**17**)¹⁹, chrysin-8-C- β -D-glucopyranoside (**18**)²⁰, baicalein-7-O-glucuronide (baicalin) (**19**)²¹, 5,7,6'-trihydroxy-2'-methoxy flavone-7-O- β -D-glucuronide (**20**)²², apigenin-7-O- β -D-glucuronopyranoside (scutellarin A) (**21**)²³, 2'-hydroxychrysin-7-glucuronide (**22**)²⁴, quercetin-7-O- β -D-glucopyranoside (quercimeritrin) (**23**)²⁵, oroxylin A-7-O- β -D-glucuronide (**25**)²⁶, wogonin-7-O- β -D-glucuronopyranoside (**26**), baicalein-7-O-sulfate (**27**)²⁷, 3,5,7,3',4'-pentahydroxy flavanone (taxifolin) (**28**)²⁸, (2R, 3R)-3,5,7,2',6'-pentahydroxy flavanone (**29**)²⁹, 3,5,7,2',6'-pentahydroxy flavanone (**30**)³⁰, 3,5,7,3',4'-pentahydroxy flavone (quercetin) (**31**), 5,7,2',6'-tetrahydroxy flavonol (visciclulin I) (**32**)³¹, 5,7,2',5'-tetrahydroxy-8,6'-dimethoxy flavone (viscidulin III) (**33**)³², 5,7,2',6'-tetrahydroxy flavone (**34**)³³, 5,2'-dihydroxy-7,8-dimethoxy flavone (panicolin) (**35**)³⁴, 5,7-dihydroxy-6-methoxy flavone (oroxylin A) (**36**)³⁵, 3,5,7,4'-tetrahydroxy flavone (kaempferol) (**37**)³⁶, 5,7,3',4'-tetrahydroxy flavone (luteolin) (**38**)³⁷, 5,7,8,4'-tetrahydroxy flavone (isoscuteallarein) (**39**)³⁸, 5,6,7,4'-tetrahydroxy flavone (scuteallarein) (**40**)³⁶, 5,7,3',4',5'-pentahydroxy-3,6,8-trimethoxy flavone (**41**)³⁹, 5,7-dihydroxy-8-methoxy flavone (wogonin) (**42**)⁴⁰, 7,4'-dihydroxy flavone (**43**)⁴¹, 5,7-dihydroxy flavone (chrysin) (**44**)⁴², 5,7-dihydroxy-4'-methoxy flavone (acacetin) (**45**)⁴³.

ACKNOWLEDGEMENTS

The authors thank the National Science and Technology Major Project Key New Drug Creation and Manufacturing Program, China (No. 2009ZX09301-001, 2009ZX09308-005) and the National Natural Science Foundation of China (No. 81030065) for financial support of this work.

REFERENCES

1. National Commission of Chinese Pharmacopoeia, Pharmacopoeia of the People's Republic of China. China Medical Technology Press, Beijing (2010).
2. Jiangsu New Medical College, Dictionary of Chinese Materia Medical, Science and Technology Press of Shanghai (1977).
3. H.Z. Wen, S.Y. Xiao, Y.M. Wang and G.A. Luo, *Nat. Prod. Res. Dev.*, **6**, 575 (2004).
4. G. Corinne, M. Frederic, B. Françoise, T. François, V. Jacqueline, S. Thierry and L. Marc, *J. Nat. Prod.*, **62**, 1188 (1999).
5. W.L. Yang, H.F. Chen, B.J. Yu, P.Z. Zhang, Z.H. Yan, L. Jie and H. Xu, *Zhong Yao Cai*, **31**, 1812 (2008).
6. H. Tokunaru, S. Kenichi, Y. Kazuyo, K. Yasuhiko and T. Masao, *Chem. Pharm. Bull.*, **45**, 446 (1997).
7. K. Masanori, Y. Yooko, F. Seigo, U. Akira, N. Tadataka and M. Toshio, *Chem. Pharm. Bull.*, **30**, 1602 (1982).
8. D.W. Aksnes, A. Standnes and O.M. Anderson, *Magn. Reson. Chem.*, **34**, 820 (1996).
9. H.J. Xia, F. Qiu, S. Zhu, T.Y. Zhang, G.X. Qu and X.S. Yao, *Biol. Pharm. Bull.*, **30**, 1308 (2007).
10. E.A. Ali, M. Ibtissam, T. Rachida Chendid, B. Mohamed, E.A. Rachid, E.B. Brahim and L. Mohammed, *Molecules*, **9**, 568 (2004).
11. E. Wollenweber, V.H. Dietz, D. Schillo and G. Schilling, *Zeitsch. Naturforschung, C: J. Biosci.*, **35**, 685 (1980).
12. D.K. Bhardwaj and R.M. Jain, *Proc. Indian Nat. Sci. Acad.*, **49**, 408 (1983).
13. L.C. Dos Santos, C.M. Rodrigues, M.A. Silva, R.G. Coelho, M. Sannomiya and W. Vilegas, *Biochem. System. Ecol.*, **33**, 1159 (2005).
14. C. Xie, N.C. Veitch, P.J. Houghton and S.J. Simmonds, *Chem. Pharm. Bull.*, **51**, 1204 (2003).
15. L. Gobbo-Neto, M.D. Santos, A. Kanashiro, M.C. Almeida, Y.M. Lucisano-Valim, L.C. Lopes, E.P. Souza and N.P. Lopes, *Planta Medica*, **71**, 3 (2005).

16. S. Takagi, M. Yamaki and K. Inoue, *Yakugaku Zasshi*, **101**, 899 (1981).
17. Y. Yuan, W.L. Hou, M.H. Tang, H.D. Luo, L.J. Chen, Y.H. Guan and L.A. Sutherland, *Chromatographia*, **68**, 885 (2008).
18. G. Romussi, N. Fontana and N.D. Tommasi, *Phytother. Res.*, **10**, S84 (1996).
19. Y.Y. Zhang, Y.Z. Guo, H. Ageta, Y. Harigaya, M. Onda, K. Hashimoto, Y. Ikeya, M. Okada and M. Maruno, *J. Chin. Pharm. Sci.*, **6**, 182 (1997).
20. Z.H. Zhou and C.R. Yang, *Yunnan Zhiwu Yanjiu*, **22**, 475 (2000).
21. T. Tomimori, Y. Miyaichi, Y. Imoto, H. Kizu and Y. Tanabe, *Yakugaku Zasshi*, **103**, 607 (1983).
22. J. Han, M. Ye, M. Xu, J.H. Sun, B.R. Wang and D. Guo, *J. Chromatogr. B*, **848**, 355 (2007).
23. T. Hase, K. Ohtani, R. Kasai, K. Yamasaki and C. Picheansoonthon, *Phytochemistry*, **40**, 287 (1995).
24. Y.Q. Wang, K. Matsuzaki, K. Takahashi, T. Okuyama and S. Shibata, *Chem. Pharmaceut. Bull.*, **36**, 3206 (1988).
25. R. Kunde and O. Isaac, *Planta Medica*, **37**, 124 (1979).
26. K. Abe, O. Inoue and E. Yumioka, *Chem. Pharm. Bull.*, **38**, 208 (1990).
27. W.H. Huang, A.R. Lee and C.H. Yang, *Biosci. Biotechnol. Biochem.*, **70**, 2371 (2006).
28. W.L. Mei, W. Ni, Y. Hua and C.X. Chen, *Nat. Prod. Res. Dev.*, **14**, 26 (2002).
29. Y.Y. Zhang and Y.Z. Guo, *Shenyang Yaoxueyuan Xuebao*, **8**, 137 (1991).
30. S. Takagi, M. Yamaki and K. Inoue, *Phytochemistry*, **20**, 2443 (1981).
31. H.Y. Wang, L.H. Xiao, L. Liu and S.X. Xu, *Shenyang Yaoke Daxue Xuebao*, **20**, 339, 348 (2003).
32. M.L. Liu, M.L. Li, X.T. Liang, X.Y. Lin, Q.T. Zheng, S.D. Zhang and F.L. Shen, *Zhongcaoyao*, **17**, 438 (1986).
33. T. Tomimori, Y. Miyaichi and H. Kizu, *Yakugaku Zasshi*, **102**, 388 (1982).
34. M. Inuma, S. Matsuura and K. Kusuda, *Chem. Pharm. Bull.*, **28**, 708 (1980).
35. R.R. Biekofsky, C.A. Buschi and A.B. Pomilio, *Magn Reson. Chem.*, **29**, 569 (1991).
36. L.H. Qin, X.Y. Guo, M. Fan, A.S. Din, N.L. Wang and X.S. Yao, *Shenyang Yaoke Daxue Xuebao*, **23**, 633 (2006).
37. J.Y. Peng, G.R. Fan and Y.T. Wu, *China J. Chin. Mater. Medica*, **31**, 128 (2006).
38. W.H. Huang, Y.B. Li and J.Q. Jiang, *Zhongguo Zhongyao Zazhi*, **34**, 712 (2009).
39. S.G. Yu, N.B. Fang and T.J. Mabry, *Phytochemistry*, **27**, 171 (1988).
40. X. Li, J.H. Wang, X. Li and D.L. Liang, *Shenyang Yaoke Daxue Xuebao*, **19**, 178 (2002).
41. C.N. He, W.W. Gao, W.Y. Xu and J.M. Tong, *Zhong Cao Yao*, **39**, 1783 (2008).
42. Z.T. Ma and X.W. Yang, *Zhongguo Zhongyao Zazhi*, **33**, 2080 (2008).
43. N.Y. Yang, J.H. Duang, P. Li, S.H. Qian and W.C. Hu, *Zhongguo Tianran Yaowu*, **4**, 98 (2006).