Synthesis and Supramolecular Structure of 3,3'-Dibromo-1,1'-[butane-1,4-diylldioxybis(nitrilomethylidyne)]dibenzene

YIN-XIA SUN*, WEI-SHENG MENG, QING-YING LAN, XIAO-YAN ZHANG, FEI-XIA MA and DA-SENG WANG

School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou 730070, P.R. China

*Corresponding author: E-mail: sun_yinxia@163.com

(Received: 3 April 2012;

Accepted: 21 December 2012)

AJC-12594

The compound, 3,3'-dibromo-1,1'-[butane-1,4-diylldioxybis(nitrilo-methylidyne)]dibenzene with the molecular formula $C_{18}H_{18}N_2O_2Br_2$, was synthesized by the reaction of 3-bromobenzaldehyde with 1,4-bis(aminooxy) butane in ethanol. The molecule lies across a crystallographic inversion centre (symmetry code: $-x, -y, -z$) and adopts an extended form. Within the molecule, the two aromatic rings are aligned with the angle of $5.57(3)^\circ$, but extend in opposite directions from the tetramethylene bridge. In the crystal structure, weak intermolecular $C-H\cdots O$ hydrogen bonds and $C-H\cdots\pi$ stacking interactions link the title molecules into an infinite 3D supramolecular network structure.

Key Words: Bisoxime compound, Synthesis, Supramolecular structure.

INTRODUCTION

Salen-type compounds are one of most prevalent mixed-donor ligands in the field of coordination chemistry¹⁻³. They play an important role in the development of coordination chemistry related to catalysis and enzymatic reactions, magnetism and supramolecular architectures. In the past few years, salen-type compounds have been used widely as ligands in the formation of transition metal complexes^{4,5}. Many salen-type complexes have been structurally characterized, but only a relatively small number of free salen-type bisoxime compounds have been characterized. As an extension of our work⁶⁻⁸, we report here the synthesis and crystal structure of a new salen-type bisoxime compound 3,3'-dibromo-1,1'-[butane-1,4-diylldioxybis(nitrilomethylidyne)]dibenzene.

EXPERIMENTAL

3-Bromobenzaldehyde was purchased from Alfa Aesar and used without further purification. 1,4-Bis(aminooxy)butane was synthesized according to an analogous method reported earlier⁶. The other reagents and solvents were analytical grade reagents from Tianjin Chemical Reagent Factory. C, H and N analyses were carried out with a GmbH VariuoEL V3.00 automatic elemental analyzer. IR spectra in the range 4000-400 cm^{-1} were recorded on a VERTEX70 FT-IR spectrophotometer using KBr pellets. The 1H NMR spectra were recorded on a Mercury-400BB spectrometer at room temperature using $CDCl_3$ as solvent. X-Ray single crystal structure was determined on a Bruker Smart APEX CCD area detector. Melting

points were measured by the use of a microscopic melting point apparatus made in Beijing Taike Instrument Limited Company and the thermometer was uncorrected.

General procedure: 3,3'-Dibromo-1,1'-[butane-1,4-diylldioxybis(nitrilomethylidyne)]dibenzene was synthesized according to an analogous method reported earlier^{7,8}. To an ethanolic solution (2 mL) of 3-bromobenzaldehyde (407.7 mg, 2.13 mmol) was added an ethanolic solution (3 mL) of 1,4-bis(aminooxy)butane (128.1 mg, 1.07 mmol). The mixture solution was stirred at 328 K for 4 h. When cooled to room temperature, the precipitate was filtered and washed successively with ethanol and ethanol-hexane (1:4), respectively. The product was dried under vacuum to yield 94.8 mg of the title compound. Yield, 19.6 %. m.p. 323.5-325.5 K. Anal. calcd. (%) for $C_{18}H_{18}N_2O_2Br_2$: C, 47.60; H, 3.99; N, 6.17. Found (%): C, 47.42; H, 3.83; N, 6.00. IR (KBr, ν_{max} , cm^{-1}): C=N, 1625 and C-O, 1176. 1H NMR (400 MHz, $CDCl_3$): 2.05 (t, $J = 4.0$ Hz, 4H, CH_2), 4.22 (t, $J = 4.0$ Hz, 4H, CH_2-O), 6.86 (d, $J = 4.0$ Hz, 2H, PhH), 6.95 (s, 2H, PhH), 7.22 (s, 2H, PhH), 7.36 (d, $J = 4.0$ Hz, 2H, PhH), 8.03 (s, 2H, N=CH). Colourless needle-shaped single crystals suitable for X-ray diffraction studies were obtained after several weeks by slow evaporation from a methanol-acetonitrile mixed solution of 3,3'-dibromo-1,1'-[butane-1,4-diylldioxybis(nitrilomethylidyne)]dibenzene.

X-Ray structure determination: The single crystal of the title compound, with approximate dimensions of 0.48 mm $\times$ 0.38 mm $\times$ 0.30 mm was placed on a Bruker Smart 1000 diffractometer equipped with Apex CCD area detector. The diffraction data were collected using a graphite monochromated

MoK α radiation ($\gamma = 0.71073$ Å) at 298(2) K. The structure was solved by using the program SHELXS-97⁹ and Fourier difference techniques and refined by full-matrix least-squares method on F^2 using SHELXL-97¹⁰. Details of the data collection and refinements of title compound are given in Table-1. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were added theoretically. CCDC: 696136.

TABLE-1
CRYSTAL DATA AND REFINEMENT PARAMETERS
FOR THE TITLE COMPOUND

Empirical formula	C ₁₈ H ₁₈ N ₂ O ₂ Br ₂
Formula weight	454.16
Temperature	298(2) K
Wavelength	0.71073 Å
Crystal system	Monoclinic
Space group	P2(1)/c
Cell dimensions	a = 11.9330(16) Å, b = 4.8703(5) Å, c = 32.423(3) Å β = 99.706(2)°
Volume	1857.4(4) Å ³
Z	4
Density (calculated)	1.624 mg/m ³
Absorption coefficient	4.377 mm ⁻¹
$F(000)$	904
Index ranges	-14 $\leq h \leq$ 11, -5 $\leq k \leq$ 5, -38 $\leq l \leq$ 34
Reflections collected	8610/3256 [R_{int} = 0.0505]
Independent reflections	3256
Data/restraints/parameters	3256/0/217
Goodness of fit indicator	1.005
R [$I > 2\sigma(I)$]	$R_1 = 0.0766$, $wR_2 = 0.1515$
Largest diff. peak and hole	0.524 and -0.538 e. Å

RESULTS AND DISCUSSION

X-Ray crystallographic analysis revealed the crystal structure of title compound. And the structure is shown in Fig. 1. Selected bond distances and angles are listed in Table-2. The crystal structure of the title compound is built up by only the C₁₈H₁₈N₂O₂Br₂ molecules and all bond lengths and angles are in normal ranges^{6,7}. The molecule assumes an extended

Fig. 1. Molecule structure of the title compound

conformation in which two benzaldehyde moieties are well separated from each other with a crystallographic inversion center (symmetry code: -x, -y, -z). Within the molecule, the two aromatic rings are aligned with the angle of 5.57(3)°, but extend in opposite directions from the tetramethylene bridge.

In the crystal structure, C-H $\cdots\pi$ (Ph) stacking interactions and weak intermolecular C-H $\cdots$ O hydrogen bonds link the neighboring molecules into an infinite three-dimensional supramolecular network structure. The title molecules are linked by a pair of intermolecular C9-H9 $\cdots\pi_{\text{centroid}}$ (C13-C18) hydrogen bond interactions into a 1D infinite zigzag chain along the c-axis. Furthermore, this linkage is further stabilized by a pair of intermolecular C16-H16 $\cdots\pi_{\text{centroid}}$ (C6-C11) hydrogen bond interactions to form an infinite 2D crapy layer parallel to the bc-plane (Fig. 2). The intermolecular C-H $\cdots\pi_{\text{centroid}}$ separations are *ca.* 3.214 Å for C16-H16 $\cdots\pi_{\text{centroid}}$ (C6-C11) and *ca.* 3.231 Å for C9-H9 $\cdots\pi_{\text{centroid}}$ (C13-C18), respectively (Table-3) and lie in the accepted distance range for this type of contacts¹¹. In addition, the adjacent crapy layers are further linked by the intermolecular C18-H18 $\cdots$ O2 hydrogen bonds between the oxime oxygen atom and the methylene groups of the benzene ring. It is noteworthy that an infinite helices chain along the b-axis is formed by the C18-H18 $\cdots$ O2 hydrogen bond interactions (Figs. 3 and 4). To sum up, with the help of intermolecular C-H $\cdots$ O and C-H $\cdots\pi$ hydrogen-bonding interactions, the crystal packing shows a self-assembling 3D supramolecular network structure.

TABLE-2
SELECTED BOND LENGTHS (Å) AND ANGLES (°) FOR THE TITLE COMPOUND

Bond	Lengths	Bond	Lengths	Bond	Lengths
Br(1)-C(8)	1.901(8)	C(9)-C(10)	1.368(12)	O(2)-C(4)	1.419(8)
N(1)-C(5)	1.253(9)	C(12)-C(13)	1.480(10)	C(2)-C(3)	1.524(10)
N(2)-C(12)	1.265(9)	C(13)-C(18)	1.413(9)	C(5)-C(6)	93(11)
O(1)-C(1)	1.438(8)	C(15)-C(16)	1.387(11)	C(6)-C(11)	1.411(11)
C(1)-C(2)	1.520(10)	C(17)-C(18)	1.383(10)	C(8)-C(9)	1.380(11)
C(3)-C(4)	1.510(9)	Br(2)-C(15)	1.861(7)	C(10)-C(11)	1.342(11)
C(6)-C(7)	1.376(10)	N(1)-O(1)	1.419(7)	C(13)-C(14)	1.372(9)
C(7)-C(8)	1.387(10)	N(2)-O(2)	1.422(8)	C(14)-C(15)	1.386(10)
C(16)-C(17)	1.376(11)	—	—	—	—
C(5)-N(1)-O(1)	108.6(7)	C(14)-C(13)-C(18)	119.8(7)	C(11)-C(6)-C(5)	118.2(8)
N(1)-O(1)-C(1)	109.7(6)	C(18)-C(13)-C(12)	118.0(7)	C(9)-C(8)-C(7)	121.0(8)
O(1)-C(1)-C(2)	106.6(6)	C(14)-C(15)-C(16)	118.9(8)	C(7)-C(8)-Br(1)	119.4(6)
C(4)-C(3)-C(2)	110.7(6)	C(16)-C(15)-Br(2)	121.8(6)	C(11)-C(10)-C(9)	123.5(9)
N(1)-C(5)-C(6)	122.2(8)	C(16)-C(17)-C(18)	119.0(8)	N(2)-C(12)-C(13)	120.1(7)
C(7)-C(6)-C(5)	121.0(7)	C(12)-N(2)-O(2)	111.1(6)	C(14)-C(13)-C(12)	122.2(7)
C(6)-C(7)-C(8)	118.7(8)	C(4)-O(2)-N(2)	108.3(5)	C(13)-C(14)-C(15)	120.5(7)
C(9)-C(8)-Br(1)	119.6(6)	C(1)-C(2)-C(3)	112.3(7)	C(14)-C(15)-Br(2)	119.3(6)
C(10)-C(9)-C(8)	118.0(8)	O(2)-C(4)-C(3)	108.4(6)	C(17)-C(16)-C(15)	121.9(8)
C(10)-(11)-C(6)	117.7(9)	C(7)-C(6)-C(11)	120.8(7)	C(17)-C(18)-C(13)	119.9(7)

Fig. 2. View of the 1D supramolecular zigzag chain and 2D crapy layers within the title compound (hydrogen atoms, except those forming hydrogen bonds, are omitted for clarity)

Fig. 3. Intermolecular C18-H18...O2 hydrogen bonds. For clarity, the H atoms not involved in the interactions have been omitted

Fig. 4. Helix chain linked by C18-H18...O2 hydrogen bonds. For clarity, the non-contact atoms not involved in the interactions have been omitted

TABLE-3 HYDROGEN-BONDING DATA [Å, °]				
D-H...A	d(D-H)	d(H...A)	d(D...A)	∠D-H...A
C18-H18...O2	0.93	2.51	3.417(3)	161
C16-H16...Cg1 ^a	0.93	3.21	4.012(4)	146
C9-H9...Cg2 ^b	0.93	3.23	4.008(4)	143

^aCg1 is the C6–C11 ring centroids. ^bCg2 is the C13–C18 ring centroid.

ACKNOWLEDGEMENTS

The authors thank Young Scholars Science Foundation of Lanzhou Jiaotong University (2011007) for financial support of this work.

REFERENCES

1. B. Sieklucka, R. Podgajny, P. Przychodzen and T. Korzeniak, *Coord. Chem. Rev.*, **249**, 2203 (2005).
2. J.D. Dyers, S.Y. Que, D. VanDerveer and X.R. Bu, *Inorg. Chim. Acta*, **359**, 197 (2006).
3. R. Klement, F. Stock, H. Elias, H. Paulus, P. Pelikan, P. Valko and M. Mazur, *Polyhedron*, **18**, 3617 (1999).
4. S. Akine, W.K. Dong and T. Nabeshima, *Inorg. Chem.*, **45**, 4677 (2006).
5. W.K. Dong, J.H. Feng and X.Q. Yang, *Synth. React. Inorg. Met.-Org. Nano-Met. Chem.*, **37**, 189 (2007).
6. W.K. Dong, J.H. Feng, L. Wang, L. Xu, L. Zhao and X.Q. Yang, *Transition Met. Chem.*, **32**, 1101 (2007).
7. W.K. Dong, J.G. Duan and G.L. Liu, *Transition Met. Chem.*, **32**, 702 (2007).
8. W.K. Dong, J.Y. Shi, J.K. Zhong, Y.Q. Tian and J.G. Duan, *Chin. J. Inorg. Chem.*, **24**, 10 (2008).
9. G.M. Sheldrick, SHELXS97, Program for Crystal Structure Determination, University of Göttingen, Germany (1996).
10. G.M. Sheldrick, *Acta Cryst.*, **A64**, 112 (2008).
11. J. Lewin Ski, J. Zachara, I. Justyniak and M. Dranka, *Coord. Chem. Rev.*, **249**, 1185 (2005).