Synthesis and Antimicrobial Activity of 4*H*, 4-Thio-2-hepta-*O*-benzoyl- β -D-lactosylimino-3-phenyl-2,3-dihydro-(1,3,5)-triazino-(2,1*b*) 6, 7 or 8-Aryl benzothiozoles (hydrochlorides)

K.M. HEDA* and S.P. DESHMUKH

P.G. Department of Chemistry, Shri Shivaji College, Akola-444 001, India

*Corresponding author: E-mail: kavita.heda25@gmail.com

(Received: 25 June 2012;

Accepted: 8 April 2013)

AJC-13207

A series of novel 4*H*, 4-thio-2-hepta-*O*-benzoyl- β -D-lactosylimino-3-phenyl-2,3-dihydro-(1,3,5)-triazino-(2,1*b*) 6,7 or 8 aryl benzothiozoles (hydrochlorides) have been synthesized by the interaction of several 1-hepta-*O*-benzoyl- β -D-lactosyl-3-aryl benzothiozoyl thiocarbamides with *N*-phenyl isocyanodichloride. These compounds were screened for their antibacterial and antifungal activities against *Escherichia coli*, *Proteus vulgaris*, *Staphylococcus aureus*, *Salmonella typhimurium*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Aspergillus niger* and *Candida albicans*. The newly synthesized compounds have been characterized by analytical and IR, ¹H NMR and mass spectral studies.

Key Words: Benzothiazoles, *N*-phenyl isocyanodichloride, 1,3,5-Triazino compounds.

INTRODUCTION

Benzoylated lactosyl nucleosides having aryl benzothiazoles are an important class of heterocyclic compounds in organic chemistry. Benzothiazoles are bicyclic ring system with multiple applications. They have diverse chemical reactivity and broad spectrum of biological activity including antibacterial and antifungal properties¹⁻³. 2-aminobenzothiazoles shows antitumor⁴ and antimalarial activity⁵. *Bis*-substituted amidino benzothiazole act as potential anti HIV agents⁶. The Schiff-base of benzothiazoles also possesses antitubercular, anticancer, antitumor, antipyretic and sterase inhibitory activity^{7,8}. Nitrogen containing heterocycles with sulphur are an important class of existing drugs. Azoles, triazoles, substituted thiazoles, thiadiazoles have been known to possess wide spectrum of activity. In our laboratory, 1-hepta-*O*-benzoyl- β -D-lactosyl-3-aryl-benzothiazoyl thiocarbamides have been prepared for the first time by the interaction of 1-hepta-*O*-benzoyl- β -D-lactosyl-isothiocyanate-with 2-amino-aryl benzothiazoles⁹⁻¹¹. In view of our interest in *N*-lactosylated heterocyclic compounds, now we report the synthesis of 4*H*-4-thio-2-hepta-*O*-benzoyl- β -D-lactosylimino-3-phenyl-2,3-dihydro-1,3,5-triazino-6,7 or 8-aryl benzothiazoles (hydrochlorides).

EXPERIMENTAL

Melting points determined are uncorrected. IR spectra were recorded in KBr on a FT-IR Perkin-Elmer RXI (4000-

450 cm⁻¹) spectrophotometer. ¹H NMR measurements were performed on a Bruker DRX-300 (300 MHz FT NMR) NMR spectrometer in CDCl₃ solution with TMS as internal reference. The Mass spectra were recorded on a THERMO Finnigan LCQ Advantage max ion trap Mass spectrometer. Optical rotation¹² [α]_D³¹ measured on a equip-tronics digital polarimeter EQ-800 at 31 °C in CHCl₃. Thin layer chromatography¹³ was performed on silica gel G and spots were visualized by iodine vapour.

Synthesis of 4*H*,4-thio-2-hepta-*O*-benzoyl- β -D-lactosylimino-3-phenyl-2,3-dihydro-(1,3,5)-triazino-(2,1*b*)6,7 or 8 phenyl benzothiazoles (3*a*) (hydrochlorides): A mixture of 1-hepta-*O*-benzoyl- β -D-lactosyl-3-phenyl benzothiazoyl-thiocarbamide (**1a-f**) (1.261 g, 0.001 M, in 15 mL CHCl₃) with *N*-phenyl isocyanodichloride (**2**) (0.173 g, 0.001 M, in 5 mL CHCl₃) was gently refluxed for 3 h. The progress of reaction was monitored by TLC. After completion of the reaction, the reaction mixture was brought to room temperature and the solvent removed under reduced pressure to obtain residue. This residue was triturated with petroleum ether (60-80 °C) to afford a pale yellow solid (**3a**). The crude product was purified by chloroform-petroleum ether (0.927 g, 83.43 %), m.p. 160 °C. The physical characterization results are summarized in Table-1 (**3a-f**). The compounds **3b-f** were also prepared by similar methods (**Scheme-I**).

3a) IR (KBr, $\nu_{\max}$ cm⁻¹): 1727.5 (C=O), 1631.1 (C=N), 1175 (C=S), 1269 (C-O), 1098 (lactosyl ring deformation), 770 (1,2- disubstituted ring), 710 (monosubstituted ring) cm⁻¹;

TABLE-1
PHYSICAL DATA FOR CHARACTERIZATION OF COMPOUNDS (3a-f) REACTANT:
N-PHENYL ISOCYANODICHLORIDE (2) (0.173 g, 0.001 M)

1-Hepta-O-benzoyl-β-D-lactosyl-3-aryl benzothiozoyl-thiocarbamides (1a-f)	g	Product	Yield (%)	m.p. (°C)	R _f value (Petroleum ether : EtOAc 6:4)	[α] _D ³¹ CHCl ₃ (c, CHCl ₃)	Analysis (%): found (required)	
							N	S
3-phenyl	1.26	3a	66.18	160	0.80	+142.8 (c, 0.7)	3.93 (3.90)	4.42 (4.45)
3- <i>o</i> -Cl-phenyl	1.29	3b	62.58	150	0.64	+128.5 (c, 0.7)	3.76 (3.80)	4.32 (4.35)
3- <i>m</i> -Cl-phenyl	1.29	3c	69.23	164	0.54	+57.14 (c, 0.7)	3.77 (3.80)	4.36 (4.35)
3- <i>p</i> -Cl-phenyl	1.29	3d	61.18	180	0.72	-155 (c, 0.7)	3.78 (3.80)	4.33 (4.35)
3- <i>o</i> -tolyl	1.27	3e	77.14	154	0.69	+114.2 (c, 0.7)	3.83 (3.86)	4.43 (4.41)
3- <i>p</i> -tolyl	1.27	3f	70	167	0.60	+28.57 (c, 0.7)	3.84 (3.86)	4.38 (4.41)

C and H analysis was found satisfactory in all cases

Scheme-I: i) CHCl₃, refluxed, 3h; where, Bz = COC₆H₅ (Benzoyl); R = (a) Phenyl, (b) *o*-Cl-phenyl, (c) *m*-Cl-phenyl, (d) *p*-Cl-phenyl, (e) *o*-tolyl, (f) *p*-tolyl

¹H NMR (δ in ppm, CDCl₃): δ 8.01-7.16 (44H, m, 7COC₆H₅, C₆H₅, C₆H₄), 6.29-3.71 (14 H, m, lactose ring protons); Mass (*m/z*): 1435 (M⁺), 1330 (M⁺-C₇H₅O), 1300 (M⁺-C₈H₇O₂), 1053 (HBL⁺), 976 (HBL⁺-C₆H₅), 948 (HBL⁺-C₇H₅O), 932 (HBL⁺-C₇H₅O₂), 918 (HBL⁺-C₈H₇O₂), 579 (TBG⁺), (found : C, 63.52; H, 4.06; N, 3.93; S, 4.42, calculated for C₇₆H₅₈N₄O₁₇S₂, 2HCl required C, 63.55; H, 4.04; N, 3.90; S, 4.45; %).

3b) IR (KBr, ν_{max}, cm⁻¹): 2963 (ali C-H), 1727.4 (C=O), 1597.0 (C=N), 1268.4 (C-O), 1101 (lactosyl ring deformation), 771.2 (monosubstituted ring) 710.3 (C-S), 558.6 (C-Cl); ¹H NMR (δ in ppm, CDCl₃): δ 8.13-7.16 (43H, m, 7COC₆H₅, C₆H₅, C₆H₃), 6.75-3.71 (14 H, m, lactose ring protons) mass (*m/z*): 1470 (M⁺), 1435 (M⁺-Cl), 1365 (M⁺-C₇H₅O), 1349 (M⁺-C₇H₅O₂), 1337 (M⁺-C₈H₇O₂), 1053 (HBL⁺), 976 (HBL⁺-C₆H₅), 948 (HBL⁺-C₇H₅O), 932 (HBL⁺-C₇H₅O₂), 579 (TBG⁺), (found : C, 62.03; H, 3.85; N, 3.76; S, 4.32, calculated for C₇₆H₅₇N₄O₁₇S₂, 2HCl required C, 62.02; H, 3.87; N, 3.80; S, 4.35; %).

3e) IR (KBr, ν_{max}, cm⁻¹): 1727.4 (C=O), 1602.1 (C=N), 1267.1 (C-O), 1099.8 (C=S) 1028.3 (lactosyl ring deformation), 863.3 (1,3 disubstituted ring), 770.6 (monosubstituted ring) 604.5 (C-S), ¹H NMR (δ in ppm, CDCl₃): δ 8.07-7.04 (43H, m, 7COC₆H₅, C₆H₅, C₆H₃), 6.58-3.71 (14 H, m, lactose ring protons), 2.53 (3H, s, Ar-CH₃); Mass (*m/z*): 1449 (M⁺), 1345 (M⁺-C₇H₅O), 1329 (M⁺-C₇H₅O₂), 1314 (M⁺-C₈H₇O₂), 1053 (HBL⁺), 976 (HBL⁺-C₆H₅), 948 (HBL⁺-C₇H₅O), 932 (HBL⁺-C₇H₅O₂), 579 (TBG⁺), (Found : C, 63.73; H, 4.12; N, 3.83; S, 4.43, calculated for C₇₆H₅₇O₁₇N₄S₂, 2HCl required C, 63.76; H, 4.14; N, 3.86; S, 4.41; %).

HBL⁺ = Hepta-O-benzoyl-β-D-lactosyl

TBG⁺ = Tetra-O-benzoyl-β-D-galactosyl

RESULTS AND DISCUSSION

Several 4*H*, 4-thio-2-hepta-O-benzoyl-β-D-lactosylimino-3-phenyl-2,3-dihydro-(1,3,5)-triazino-(2,1*b*)6,7 or 8-aryl benzothiozoles (hydrochlorides) (**3a-f**) have been prepared by the interaction of several 1-hepta-O-benzoyl-β-D-lactosyl-3-aryl benzothiozoyl thiocarbamides (**1a-f**) with *N*-phenyl isocyanodichloride (**2**) in CHCl₃. After condensation, the solvent was distilled off to obtain a sticky residue. This residue was triturated several times with petroleum ether (60-80 °C) to afford a pale yellow solid (**3a-f**). The product was found to be non-desulphurizable when boiled with alkaline lead acetate solution. The specific rotations were measured in chloroform. The result is summarized in Table-1. In spectral analysis of products shows bands due to Ar-H, ali. C-H, C=O, C=N, C-N, C-O, C-S, C=S stretching and ¹H NMR spectra of products distinctly displayed signals due to aromatic protons and lactose ring protons. The mass spectrum of products was also observed. The identities of these new *N*-lactosides have been established on the basis of usual chemical transformations and also IR, ¹H NMR and mass spectral studies¹⁴⁻¹⁶.

Antimicrobial activity: All the compounds have been screened for both antimicrobial and antifungal activity by using disc diffusion assay. For this sterile filter paper disc (6 mm) impregnated with fixed doses of compounds was placed on pre-inoculated surface. The disc bearing plates were incubated at 37 °C for 24 h. Inhibition zones read after incubation at 37 °C for 24 h. for bacterial strains and for fungal strains inhibition zones read after incubation at 35 °C for 48 h. The compounds were taken at a concentration or 1 mg/mL using dimethyl sulphoxide as a solvent. Amikacin (100 µg/mL) was used as standard for antibacterial and Fluconazole (100 µg/mL) as a standard for antifungal activity. The compound were screened for antibacterial activity against *Escherichia coli*, *Proteus vulgaris*, *Staphylococcus aureus*, *Salmonella typhi*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* in Mullar-Hilton medium. *Aspergillus niger* and *Candida albicans* in potato dextrose agar medium. It has been observed that all the compounds showed nearly same activity against bacteria and fungi. **3a**, **3b**, **3d** and **3f** exhibits most significant activity against *Salmonella typhi*. All other compounds exhibited low to moderate activity (Table-2).

ACKNOWLEDGEMENTS

The authors acknowledged to SAIF, CDRI, Lucknow for providing the spectral data. Thanks are also due to the Principal

TABLE-2
ANTIMICROBIAL ACTIVITY OF 4H, 4-THIO-2-HEPTA-O-BENZOYL- β -D-LACTOSYLIMINO-3-PHENYL-2, 3-DIHYDRO-(1, 3, 5)-TRIAZINO-(2, 1B) 6, 7 OR 8 ARYL BENZOTHIOZOLES (HYDROCHLORIDES)

Compound	<i>E. coli</i>	<i>S. aureus</i>	<i>P. vulgaris</i>	<i>P. aeruginosa</i>	<i>S. typhi</i>	<i>K. pneumoniae</i>	<i>A. niger</i>	<i>C. albicans</i>
3a	14	10	14	-	17	11	19	20
3b	10	16	-	12	18	13	20	21
3c	13	14	12	13	15	14	17	19
3d	14	15	13	11	16	12	20	19
3e	16	13	10	10	15	10	21	22
3f	13	14	-	17	19	13	20	20
Amikacin	18	21	23	19	20	21	-	-
Fluconazole	-	-	-	-	-	-	24	24
DMSO	-	-	-	-	-	-	-	-

Zone of inhibition in mm (15 or less) resistance, (16-20 mm) moderate and more than (20 mm) sensitive

Dr. S.G. Bhadange for encouragement and providing the necessary research facilities. The authors also thanks Dr. Rupali Mantri (M.D. Microbiology), Government Medical College, Akola, India for her help in screening antimicrobial activity.

REFERENCES

1. M. Lacova, J. Chavaneova, O. Hyblova and S. Varkonda, *Chem. Pap.*, **45**, 411 (1991).
2. L. Ballell, R.A. Field, K. Duncan and R. Young, *J. Antimicrob. Agents Chemother.*, **49**, 2153 (2005).
3. V.P. Trivedi, N.K. Undavia and P.B. Trivedi, *J. Indian Chem. Soc.*, **81**, 506 (2004).
4. T.D. Bradshaw, M.S. Chua, H.L. Browne, V. Tarpan, E.A. Sausville and M.F.G. Stevens, *Br. J. Cancer*, **86**, 1348 (2002).
5. S. Hout, N. Azas, A. Darque, M. Robin, C. Di Giorgio, M. Gasquet, J. Galy and Timan-David, *Parasitology*, **129**, 525 (2004).
6. L. Racane, V. Tralic-Kulenovic, L. Fiseo-Jakie, D.W. Boykin and G. Karmiski-Zamola, *Heterocycles*, **55**, 2085 (2001).
7. M.S. Shingare and D.B. Ingle, *J. Indian Chem. Soc.*, **53**, 1036 (1976).
8. B. Dash and M. Patra, *Indian J. Chem.*, **19B**, 894 (1980).
9. M.G. Dhonde and S.P. Deshmukh, *Int. J. Chem. Sci.*, **3**, 135 (2005).
10. S.P. Mote and S.P. Deshmukh, *Rasayan J. Chem.*, **4**, 29 (2011).
11. G.V. Korpe, S.P. Deshmukh and A.K. Fokmare, *Indian, J. Heterocyc. Chem.*, **10**, 287 (2001).
12. Arnold Weissberger, *Physical Methods of Organic Chemistry*, Part II, Interscience Publisher, INC, New York, edn. 2 (1949).
13. B. Fried and S.H. Sherama, *Thin-Layer Chromatography*, Chromatographic Science Series, New York, Dekkert, edn. 2 (1986).
14. R.M. Silverstein and F.X. Webster, *Spectrometric Identification of Organic Compounds*, John Wiley & Sons, Inc, New York, edn. 6 (2004).
15. N.B. Colthup, L.H. Daly and S.E. Wibereley, *Introduction to Infrared and Raman Spectroscopy* Academic Press, New York (1964).
16. D.H. Williams and L. Fleming, *Spectroscopic Methods in Organic Chemistry*, Tata McGraw-Hill, edn. 4 (1988).