NOTE

Theoretical Studies of the Glycoluril Molecular Clips Using Quantum Calculations

F. GHARAKHANI¹, E. VESSALLY^{1,*}, A. MOHAMMADI² and Z. ALIMADADI³

¹Department of Chemistry, Miyaneh Branch, Islamic Azad University, Miyaneh, Iran

²Payame Noor University, Zanjan, Iran

³Department of Chemistry, Tabriz Branch, Islamic Azad University, Tabriz, Iran

*Corresponding author: E-mail: vessally@yahoo.com

(Received: 24 August 2012;

Accepted: 5 July 2013)

AJC-13771

Theoretical calculations were performed to determine the structure of a simple glycoluril (**1**) and its derivative (**2**). Full geometry optimizations were carried out by MP2 method using 6-311++G** basis set of the GAUSSIAN 98 program. The geometrical parameters including bond lengths (R) and bond angles (A) were estimated at the same level.

Key Words: Glycoluril, Quantum calculations, Molecular clip.

The glycolurils have been received a great deal of attention due to their practical applications, such as fertilizers¹ polymer crosslinking², explosives³, stabilizers of organic compounds⁴, combinatorial chemistry⁵, bleaching activators⁶ and in supramolecular chemistry⁷.

Follow up of our work⁸, the MP2 calculations were carried out on the simple glycoluril (**1**) and its derivative which have two benzene walls, **2**.

Full geometry optimizations were carried out for a simple glycoluril (**1**) and its derivative with two benzene walls, **2**, by the MP2 method using 6-311++G** basis set of the GAUSSIAN 98 program⁹⁻¹¹ (**Scheme-I**). To find a global minimum on a specific surface, all possible conformations of the given species were examined through scanning the specific dihedral angles at B3LYP/6-311++G** level. All calculations were carried out for gas phase at 298 K temperature and 1 atm pressure.

Scheme-I: Molecular structure of glycoluril (**1**) and its derivative (**2**)

Thermal energies (E), enthalpies (H) and Gibbs free energies (G) were calculated for a simple glycoluril, **1** and its

derivative (**2**) with two benzene walls at MP2/6-311++G**//B3LYP/6-311++G** level of theory (**Scheme-I**). Geometrical parameters including bond lengths (R) and bond angles (A) were estimated at the same level.

The C=O bond lengths in **1** and **2** are 1.209 and 1.215 Å, respectively which indicate the C=O bond is stronger for **1** with respect to **2**. Moreover, the N-C (amide) bond lengths in **1** and **2** are 1.391 and 1.386 Å, respectively. Therefore, the C=O bond length in **1** and **2** changed in contrast to the N-C (amide) bond length which indicate that the conjugation of the lone pair electrons of nitrogen atoms with the C=O bond in **2** is more than **1**.

The N-CH bond lengths in **1** and **2** are 1.454 and 1.449 Å, respectively. The CH-CH bond length in **1** and **2** are 1.567 and 1.558 Å, respectively. Two carbon atoms of two benzene rings selected to compute the distance between the two benzene rings in **2**. The distance of two carbon atoms of two benzene rings in **2** is 6.587.

The N-C-N bond angles in **1** and **2** are 116.26° and 84.97°, respectively. It seems that two benzene rings lead to decrease the bond lengths of the N-C-N **2**.

The highest occupied orbital molecular HOMO and lowest unoccupied molecular orbital LUMO of **1** and **2** were calculated. It appears that the HOMO and LUMO of **1** are -11.53 and 1.23 eV, respectively. The HOMO and LUMO of **2** are -8.97 and 1.29 eV, respectively. The energy of the HOMO in **2** is higher with respect to **1** which indicates the nucleophilic character of **2**. The LUMO of **1** is similar to **2** that indicate identical electrophilic character of **1** and **2**.

Conclusion

Full geometry optimizations were carried out for a simple glycoluril (**1**) and its derivative (**2**) with two benzene walls, by DFT method using 6-311++G** basis set of the GAUSSIAN 98 program. The C=O bond length in **1** and **2** changed in contrast to the N-C (amide) bond length which indicate that the conjugation of the lone pair electrons of nitrogen atoms with the C=O bond in **2** is more than **1**. The HOMO and LUMO studies indicated that the nucleophilic character of **2** is more than **1**.

ACKNOWLEDGEMENTS

Islamic Azad University, Miyaneh Branch is gratefully acknowledged for financial support of this research.

REFERENCES

1. T.M. Addiscott and V.H. Thomas, *Chem. Ind. (London)*, 29 (1979).
2. A. Wang and D. Bassett, US Patent 4310450 (1982).
3. J. Boileau, M. Carail, E. Wimmer, R. Gallo and M. Pierrot, *Propell. Explos. Pyrotech.*, **10**, 118 (1985).
4. A. Krause, A. Aumuller, E. Korona and H. Trauth, US Patent 5670613 (1997).
5. K.E. Pryor and J. Rebek, *Org. Lett.*, **1**, 39 (1999).
6. S. Karcher, A. Kornmuller and M. Jekel, *Water Sci. Technol.*, **40**, 425 (1999).
7. A.X. Wu, J.C. Fettingner and L. Isaacs, *Tetrahedron*, **58**, 9769 (2002).
8. E. Vessally and F. Hokmabadi, *Tetrahedron Lett.*, **53**, 5645 (2012).
9. C. Lee, W. Yang and R.G. Parr, *Phys. Rev. B*, **37**, 785 (1988).
10. A.D. Becke, *J. Chem. Phys.*, **98**, 5648 (1993).
11. M.J. Frisch, *et al.*, Gaussian 98, Revision A. 6, Gaussian Inc., Pittsburgh, PA (1998).