Facile Synthesis of 8-Hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]chromene-3-carbaldehyde

V.V.A. KUMAR, D. ADISESHU and K. MOHANRAJU*

Department of Polymer Science and Technology, Sri Krishnadevaraya University, Anantapur-515 003, India

*Corresponding author: Fax:+91 8554 255710; Tel.: 91 8554 255655; E-mail: kumarchesku@gmail.com; kumarchesku@gmail.com

(Received: 5 September 2012;

Accepted: 29 May 2013)

AJC-13561

7-Hydroxy-8-allyl-2H-3-chromene carbaldehyde and *m*-chloro benzoic acid in chloroform on heating to gave 8-hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]chromene-3-carbaldehydes (**3a-d**) and 2H-3-chromene carbaldehyde reaction with malononitrile to gave 9-amino-6H-benzochromene-8,10-dicarbonitriles (**6a-d**) in good yields.

Key Words: *m*-Chloro benzoic acid, 7-Hydroxy-8-allyl chromene carbaldehyde, Piperidine, Knoevenagel condensation.

INTRODUCTION

Chromones and isoflavones constitute an important class of oxygen heterocyclics. Substituted as well as heterocyclic ring fused chromones and isoflavones have a wide range of pharmacological activity¹⁻⁵. Chromones and isoflavones with medicinal use are Khellin a coronary vasodilator. Chromones-2-carboxylate spasmolytic agent and disodium chromo glycate and anti elergitic drug. Genstein having estrogen hormonal activity⁶⁻⁸ and 7-isopropoxy isoflavones for treatment of post-menopausal and senile osteoporosis^{9,10}.

With a view to synthesize a new heterocyclic ring fused chromones and isoflavones, we studied the Knoevenagel condensation of 8-formyl-7-hydroxychromones and isoflavones^{11,12}. Literature shows that Knoevenagel condensation reaction of 2-hydroxy benzaldehyde proceeds *via* acrylo intermediate to gives rise to either three substituted 2-H chromones¹³⁻¹⁵. Selective formation of 2-hydroxy benzoldehyde depends on solvent and structural features of substrate.

EXPERIMENTAL

Melting points were determined on a Polmon instrument (model no. MP 96). IR spectra were recorded on FT-IR Perkin-Elmer 1605 spectrometer and ¹H NMR (200 MHz) and ¹³C NMR (50.3 MHz) were recorded on a Varian Gemini 200 spectrometer using TMS as internal standard (chemical shifts and ppm). UV spectra were obtained on a Shimadzu UV-visible spectrophotometer (model UV-1601). Mass spectra were recorded on a VG micromass70-70H instrument.

General procedure for the synthesis of 8-hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]chromene-3-carbaldehydes (3a-d**)**

7-Hydroxy-8-allyl-2H-3-chromenecarbaldehyde (**2a**) (2.16 g, 10 mmol), *m*-chloro benzoic acid (1.62 g) (10 mmol) were dissolved in dry CHCl₃ (25 mL) and refluxed for 6 h. After cooling to room temperature, the separated *m*-chloro benzoic acid was filtered. The filtrate was washed with aqueous NaHCO₃ (2 %, 3 × 100 mL) to remove traces of *m*-chloro benzoic acid and then with water (2 × 100 mL) dried and concentrated. The residue on chromatography over silica gel by eluting with petroleum ether:ethylacetate (8:2) gave 8-hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]chromene-3-carbaldehyde (**3a**) (1.2 g) (51 % yield), which was recrystallized from chloroform as colourless needles, m.p. 150 °C (**Scheme-I**).

Spectral data

8-Hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]chromene-3-carbaldehydes (3a**):** IR (KBr, $\nu_{\max}$, cm⁻¹): 1014 (C-O), 1624 (C=C) and 1664 (C=O). UV (MeOH): 241 nm (log ϵ 3.7), 263 nm (log ϵ 3.6) and 348 nm (log ϵ 3.7). ¹H NMR (200 MHz) (CDCl₃): δ 2.95 (dd, *J* = 16.0, 8.0 Hz, CH₂-9, H_A), 3.20 (dd, *J* = 16.0, 10.0 Hz, CH₃, H_B), 3.75 (dd, *J* = 16.0, 6.0 Hz, 8-

CH₂OH, H_D), 3.90 (dd, $J = 16.0, 4.0$ Hz, 8-CH₂OH, H_E), 5.05 (s, OCH₂-2), 5.36 (m, H-8), 6.42 (d, $J = 10.0$ Hz, H-6), 7.00 (d, $J = 10.0$ Hz, H-5), 7.18 (s, H-4), 9.50 (s, CHO). FABMS: m/z 233 (M+1), m/z 231 (M-1).

6-Chloro-8-hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]-chromene-3-carbaldehydes (3b): Recrystallized from chloroform as colourless needles, m.p. 155 °C. IR (KBr, $\nu_{\max}$, cm⁻¹): 1014 (C-O), 1624 (C=C) and 1664 (C=O). UV (MeOH): 241 nm (log ϵ 3.7), 263 nm (log ϵ 3.6) and 348 nm (log ϵ 3.7). ¹H NMR (200 MHz) (CDCl₃): δ 2.95 (dd, $J = 16.0, 8.0$ Hz, CH₂-9, H_A), 3.20 (dd, $J = 16.0, 10.0$ Hz, CH₃, H_B), 3.75 (dd, $J = 16.0, 6.0$ Hz, 8-CH₂OH, H_D), 3.90 (dd, $J = 16.0, 4.0$ Hz, 8-CH₂OH, H_E), 5.05 (s, OCH₂-2), 5.36 (m, H-8), 7.00 (d, $J = 10.0$ Hz, H-5), 7.18 (s, H-4), 9.50 (s, CHO). FABMS: m/z 268 (M+1), m/z 266 (M-1).

6-Bromo-8-hydroxymethyl-8,9-dihydro-2H-furo[2,3-h]-chromene-3-carbaldehydes (3c): Recrystallized from chloroform as colourless needles, m.p. 160 °C. IR (KBr, $\nu_{\max}$, cm⁻¹): 1014 (C-O), 1624 (C=C) and 1664 (C=O). UV (MeOH): 241 nm (log ϵ 3.7), 263 nm (log ϵ 3.6) and 348 nm (log ϵ 3.7). ¹H NMR (200 MHz) (CDCl₃): δ 2.95 (dd, $J = 16.0, 8.0$ Hz, CH₂-9, H_A), 3.20 (dd, $J = 16.0, 10.0$ Hz, CH₃, H_B), 3.75 (dd, $J = 16.0, 6.0$ Hz, 8-CH₂OH, H_D), 3.90 (dd, $J = 16.0, 4.0$ Hz, 8-CH₂OH, H_E), 5.05 (s, OCH₂-2), 5.36 (m, H-8), 7.00 (d, $J = 10.0$ Hz, H-5), 7.18 (s, H-4), 9.50 (s, CHO). FABMS: m/z 311 (M+1), m/z 309 (M-1).

8-Hydroxymethyl-8,9-dihydro-6-methyl-2H-furo[2,3-h]-chromene-3-carbaldehydes (3d): Recrystallized from chloroform as colourless needles, m.p. 158 °C. IR (KBr, $\nu_{\max}$, cm⁻¹): 1014 (C-O), 1624 (C=C) and 1664 (C=O). UV (MeOH): 241 nm (log ϵ 3.7), 263 nm (log ϵ 3.6) and 348 nm (log ϵ 3.7). ¹H NMR (200 MHz) (CDCl₃): δ 2.95 (dd, $J = 16.0, 8.0$ Hz, CH₂-9, H_A), 3.20 (dd, $J = 16.0, 10.0$ Hz, CH₃, H_B), 3.75 (dd, $J = 16.0, 6.0$ Hz, 8-CH₂OH, H_D), 3.90 (dd, $J = 16.0, 4.0$ Hz, 8-CH₂OH, H_E), 5.05 (s, OCH₂-2), 5.36 (m, H-8), 7.00 (d, $J = 10.0$ Hz, H-5), 7.18 (s, H-4), 9.50 (s, CHO). FABMS: m/z 247 (M+1), m/z 245 (M-1).

General procedure for the synthesis of 9-amino-6H-benzochromene-8,10-dicarbonitriles (6a-d)

2H-3-Chromene carbaldehyde (3.2 g, 20 mmol) (**4a**) is dissolved in ethanol (20 mL) and piperidine (3 mL) is added and the reaction mixture was stirred for 5 h and then 2.64 g (40 mmol) of malononitrile is added and stirring is continued for 24 h at room temperature. After completion of the reaction, solvent was removed under reduced pressure affording a gum. This on column chromatography over silica gel and elution with petroleum ether:ethylacetate (7:3) gave 9-amino-6H-benzochromene-8,10-dicarbonitrile (**6a**) (2.6 g yield) which was recrystallized from chloroform as pale yellow needles m.p. 172 °C (Scheme-II).

Spectral data

9-Amino-6H-benzochromene-8,10-dicarbonitrile (6a): IR (KBr, $\nu_{\max}$, cm⁻¹): 1027 (C-O), 1203 (C=O), 2360 (CN), 3358 (NH₂). UV (MeOH): 255 nm (log ϵ 4.2), 249 nm (log ϵ 4.0), 349 nm (log ϵ 3.7) and 387 nm (log ϵ 3.9). ¹H NMR (200 MHz) (CDCl₃ + DMSO): δ 4.86 (s, OCH₂), 6.02 (bs, NH₂), 7.02 (bd, $J = 10$ Hz, H-4), 7.15 (dd, $J = 10, 10$ Hz, H-3), 7.40

Scheme-II

(dd, $J = 10.0, 10.0$ Hz, H-2), 7.50 (m, H-2,3,4), 7.45 (s, H-7) 8.38 (dd, $J = 10.0, 2.0$ Hz, H-1). ¹³C NMR (50.3 MHz) (CDCl₃ + DMSO): δ 66.7 (OCH₂-6), 90.5 (C-10), 94.3 (C-8), 116.2 (C=N, C-10), 116.6 (C=N, C-8), 117.8 (C-4), 120.0 (c-10b), 121.6 (C-6a), 122.2 (C-2), 126.0 (C-3), 132.6 (C-1), 134.3 (C-&), 137.6 (C-10a), 153.8 (C-9) 156.5 (C-4a). MS: m/z 247 (M+), 246 (M-1).

9-Amino-4-chloro-6H-benzochromene-8,10-dicarbonitrile (6b): Recrystallized from chloroform as pale yellow needles m.p. 167 °C. IR (KBr, $\nu_{\max}$, cm⁻¹): 2215 (C-N), 2335 (C=O), 3354 (NH₂). UV (MeOH): 241 nm (log ϵ 4.2), 255 nm (log ϵ 4.0), 293 nm (log ϵ 3.7) and 381 nm (log ϵ 3.9). ¹H NMR (200 MHz) (CDCl₃ + DMSO): δ 4.90 (s, OCH₂), 6.30 (bs, NH₂), 7.05 (bd, $J = 10$ Hz, H-4), 6.37 (dd, $J = 10, 10$ Hz, H-3), 7.40 (dd, $J = 10.0, 10.0$ Hz, H-1), 7.50 (m, H-2,3,4), 7.55 (s, H-7) 8.38 (dd, $J = 10.0, 2.0$ Hz, H-1). ¹³C NMR (50.3 MHz) (CDCl₃ + DMSO): δ 64.7 (OCH₂-6), 94.5 (C-10), 95.3 (C-8), 116.2 (C=N, C-10), 116.6 (C=N, C-8), 119.0 (c-10b), 126.6 (C-6a), 122.2 (C-2), 129.0 (C-3), 132.6 (C-1), 134.3 (C-2), 137.6 (C-10a), 153.4 (C-9) 156.5 (C-4a). MS: m/z 281 (M⁺), 217, 189, 164 and 138.

9-Amino-4-bromo-6H-benzochromene-8,10-dicarbonitrile (6c): Recrystallized from chloroform as pale yellow needles m.p. 164 °C. IR (KBr, $\nu_{\max}$, cm⁻¹): 1018 (C-O), 1210 (C=O), 1322 (NH₂). UV (MeOH): 249 nm (log ϵ 4.6), 296 nm (log ϵ 4.4), 337 nm (log ϵ 4.2) and 383 nm (log ϵ 4.5). ¹H NMR (200 MHz) (CDCl₃ + DMSO): δ 4.90 (s, OCH₂), 5.32 (bs, NH₂), 7.00 (bd, $J = 10$ Hz, H-4), 7.55 (dd, $J = 10, 10$ Hz, H-3), 7.50 (m, H-2,3,4), 7.45 (s, H-7) 8.55 (dd, $J = 10.0, 2.0$ Hz, H-1). ¹³C NMR (50.3 MHz) (CDCl₃ + DMSO): δ 67.2 (OCH₂-6), 95.6 (C-10), 95.8 (C-8), 116.0 (C=N, C-10), 116.2 (C=N, C-8), 119.8 (c-10b), 126.6 (C-6a), 120.2 (C-2), 135.0 (C-3), 132.6 (C-1), 134.3 (C-2), 135.6 (C-10a), 153.8 (C-9) 156.5 (C-4a). MS: m/z 326 (M⁺), 218, 189, 164.

9-Amino-4-methyl-6H-benzochromene-8,10-dicarbonitrile (6d): Recrystallized from chloroform as pale yellow needles m.p. 152 °C. IR (KBr, $\nu_{\max}$, cm⁻¹): 2211 (C=N), 2344 (CN), 3367 (NH₂). UV (MeOH): 244 nm (log ϵ 4.1), 302 nm (log ϵ 3.7), 336 nm (log ϵ 3.5) and 383 nm (log ϵ 3.8). ¹H NMR (200 MHz) (CDCl₃ + DMSO): δ 2.25 (s, 4-CH₃), 4.89 (s, OCH₂), 6.30 (bs, NH₂), 7.05 (bd, $J = 10$ Hz, H-2), 7.25 (dd, $J = 10, 10$ Hz, H-3), 7.58 (s, H-7) 8.18 (dd, $J = 10.0, 2.0$ Hz, H-1). ¹³C NMR (50.3 MHz) (CDCl₃ + DMSO): δ 18.2 (4-CH₃), 67.8 (OCH₂-6), 92.4 (C-10), 95.2 (C-8), 116.0 (C=N, C-8), 120.8 (c-10b), 123.6 (C-6a), 124.2 (C-2), 127.0 (C-3), 132.4 (C-1), 136.6 (C-10a), 153.4 (C-9) 155.8 (C-4a). MS: m/z 261 (M⁺), 232, 199, 130.

REFERENCES

1. H. Staudinger, *Justus Liebigs Ann. Chem.*, **356**, 51 (1907).
2. F.H. van der Steen and G. van Koten, *Tetrahedron*, **47**, 7503 (1991).
3. W.R. Vaughan and R.S. Klonowski, *J. Org. Chem.*, **26**, 145 (1961).
4. G.C. Look, M.M. Murphy, D.A. Campbell and M.A. Gallop, *Tetrahedron Lett.*, **36**, 2937 (1995).
5. B.K. Banik, F.F. Becker, *Tetrahedron Lett.*, **41**, 6551 (2000).
6. I. Banik, L. Hackfeld and B.K. Banik, *Heterocycles*, **59**, 505 (2003).
7. D.A. Nelson, *J. Org. Chem.*, **37**, 1447 (1972).
8. S.D. Sharma, G. Singh and P.K. Gupta, *Indian J. Chem.*, **16B**, 74 (1978).
9. V.V. Govande, M. Arun, A.R.A.S. Deshmukh and B.M. Bhawal, *Synth. Commun.*, **30**, 4177 (2000).
10. M. Arun, V.V. Govande, A.R.A.S. Deshmukh and B.M. Bhawal, *Indian J. Chem.*, **41B**, 856 (2002).
11. J.M. Aizpurua, I. Ganboa, F.P. Cossio, A. Gonzalez, A. Arrieta and C. Palomo, *Tetrahedron Lett.*, **25**, 3905 (1984).
12. A. Arrieta, F.P. Cossio and C. Palomo, *Tetrahedron*, **41**, 1703 (1985).
13. M. Miyake, M. Kinsawa and N. Tokutake, *Synthesis*, 1053 (1982).
14. D. Krislmaswamy, V.V. Govande, V.K. Gumaste, B.M. Bhawal and A.R.A.S. Deshmukh, *Tetrahedron*, **58**, 2215 (2002).
15. A. Arrieta, J.M. Aizpurua and C. Palomo, *Tetrahedron Lett.*, **25**, 3365 (1984).