Synthesis and Dipolymer Structure of Salen-type N₂O₂ Ligand 2,2'-[(Pentane-1,5-diylldioxy-bis)-bis-(nitrilomethylidyne)]dinaphthalene

LI ZHAO*, YU-JIE ZHANG, SHOU-TING ZHANG, HONG-XIA GUO and QIAN CHENG

School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou 730070, P.R. China

*Corresponding author: E-mail: zhaoli_72@163.com

(Received: 21 September 2012;

Accepted: 18 February 2013)

AJC-13042

The compound, 2,2'-[(pentane-1,5-diylldioxy-bis)-bis-(nitrilomethylidyne)]dinaphthalene has been synthesized by 1,5-bis-(aminooxy)pentane and 2-naphthaldehyde in ethanol solution and characterized structurally by X-ray crystallography. The molecule of the title compound, C₂₇H₂₆N₂O₂, assumes a dipolymer structure through a weak C-H...C hydrogen-bonding interactions. And the single crystal structure of the compound revealed all-conformation of the (-CH=N-O-(CH₂)₅-O-N=CH-) linkage, which resulted in the structure with two nitrilomethylidyne units apart from each other. The compound is sufficiently stable to resist scrambling of the C=N bonds. This may be ascribed to lower reactivity of the C=N-O bonds toward nucleophiles.

Key Words: Salen-type bisoxime compound, Synthesis, Dipolymer, Structure.

INTRODUCTION

Salen-type bisoxime compounds, consisting of two nitrogen and two oxygen donors, are capable of forming stable metal complexes¹ which contain one, two or more metal centers and form homo- and heteronuclear complexes serving as catalysts², models of reaction centers of metallo-enzymes³, models of reaction centers of metalloenzymes⁴, nonlinear optical materials and molecular recognition and biological agents⁵. And bisoxime complexes offer both high reactivity and selectivity include epoxidation of olefins, asymmetric ring-opening of epoxides, olefin aziridination, olefin cyclopropan-ation and formation of cyclic, linear polycarbonates⁶ and building blocks for cyclic supramolecular structures⁷. Thus, new materials can be produced by using these compounds, which seem to be suitable candidates for further chemical modifications⁸.

Recently, a preferable class of Salen-type bisoxime ligands have been reported by using an O-alkyloxime unit (-CH=N-O-(CH₂)₂-O-N=CH-) instead of the (-CH=N-(CH₂)₂-N=CH-) group and the large electronegativity of oxygen atoms is expected to affect strongly the electronic properties of N₂O₂ coordination sphere, which can lead to different and novel properties and structures of the resulting complexes⁹⁻¹³. Here, we report on the synthesis and dipolymer structure of Salen-type N₂O₂ compound, 2,2'-[(pentane-1,5-diylldioxybis)bis-(nitrilomethylidyne)]dinaphthalene.

EXPERIMENTAL

2-Naphthaldehyde, *N*-hydroxyphthalimide, 1,5-dibromopentane, triethylamine, hydrazine hydrate and anhydrous magnesium sulfate were purchased and used without further purification. The other reagents and solvents were analytical grade reagents from Tianjin Chemical Reagent Factory. C, H and N analyses were carried out with a GmbH VariuoEL V3.00 automatic elemental analyzer. X-ray single crystal structure was determined on a Bruker Smart 1000 CCD area detector. Melting points were measured by the use of a microscopic melting point apparatus made in Beijing Taike Instrument Limited Company and the thermometer was uncorrected.

1,5-Bis-(aminooxy)pentane was synthesized according to an analogous method reported earlier¹⁴. Yield: 53.4 %.

2,2'-[(Pentane-1,5-diylldioxy-bis)-bis-(nitrilomethylidyne)]-dinaphthalene was synthesized according to an analogous method reported earlier¹⁵. To an ethanolic solution (8 mL) of 2-naphthaldehyde (203.6 mg, 1.3 mmol) was added an ethanol solution (5 mL) of 1,5-bis-(aminooxy)pentane (121.8 mg, 0.6 mmol). The reaction mixture was stirred at 328-333 K for 4 h. The formed precipitate was separated by filtration and washed successively with ethanol and *n*-hexane. The product was dried under vacuum to yield 124.6 mg of the title compound. Yield, 65.2 %. m.p. 388-389 K. Anal. calcd. for C₂₇H₂₆N₂O₂ (%): C, 79.00; H, 6.38; N, 6.82. Found (%): C, 79.28; H, 6.25; N, 6.70.

After a solution of the title compound (5.0 mg, 0.012 mmol) in ethanol (4 mL) was allowed to stand at room temperature for about 2 weeks, several colourless block-like single crystals suitable for X-ray crystallographic analysis were obtained.

X-Ray structure determination: The single crystal of the title compound, with approximate dimensions of 0.40 × 0.37 × 0.11 mm was placed on a Bruker Smart 1000 diffractometer equipped with Apex CCD area detector. The diffraction data were collected using a graphite monochromated MoK α radiation ($\lambda = 0.71073$ Å) at 298(2) K. The structure was solved by using the program SHELXS-97 and Fourier difference techniques and refined by full-matrix least-squares method on F² using SHELXL-9716. Details of the data collection and refinements of the title compound are given in Table-1. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were added theoretically. CCDC: 901641.

RESULTS AND DISCUSSION

X-ray crystallographic analysis revealed the crystal structure of the title compound. The structure is shown in Fig. 1 and packing arrangement of the unit cell of the title compound is given in Fig. 2 and the dipolymer structure of the compound is shown in Fig. 3. Crystal data and structure refinement for the title compound are listed in Table-1. Selected bond distances and angles are listed in Table-2. The single crystal structure of the compound is built up by only the C₂₇H₂₆N₂O₂ molecule, in which all bond lengths are in normal ranges. The title compound is a typical Salen-type bisoxime derivative with normal geometric parameters. The X-ray diffraction analysis revealed all-conformation of the (-CH=N-O-(CH₂)₅-O-N=CH-) linkage, which resulted in the structure with two nitrilomethyl

TABLE-1
CRYSTAL DATA AND STRUCTURE REFINEMENT
FOR THE TITLE COMPOUND

Empirical formula	C ₂₇ H ₂₆ N ₂ O ₂
Formula weight	410.58
Temperature, (K)	298(2)
Wavelength (Å)	0.71073
Crystal system	Monoclinic
Space group	C2/c
Cell dimensions, (Å°)	a = 10.2673(9), b = 10.3060(11), c = 10.3060(11), β = 96.4800
Volume, Å ³	4477.8(7)
Z	8
Density (calculated) (mg/m ³)	1.218
Absorption coefficient (mm ⁻¹)	0.077
F(000)	1744
Index ranges	-12 ≤ h ≤ 12, -12 ≤ k ≤ 8, -50 ≤ l ≤ 50
Reflections collected	10790/7230 [R(int) = 0.0645]
Max. and min. transmission	0.9916 and 0.9699
Data/restraints/parameters	7230/2/559
Goodness of fit indicator	1.051
R [I > 2σ(I)]	R ₁ = 0.1365, wR ₂ = 0.2918
Largest diff. peak and hole, eÅ ⁻³	0.446 and -0.223

-idyne units apart from each other. The compound is sufficiently stable to resist scrambling of the C=N bonds. This may be ascribed to lower reactivity of the C=N-O bonds toward nucleophiles.

In the crystal structure, packing arrangement of the unit cell of the title compound revealed that each molecule links another neighboring molecule through a weak C-H...C hydrogen-bonding interactions into a dipolymer structure (Fig. 3).

TABLE-2
SELECTED BOND DISTANCES (Å) AND ANGLES (°) FOR THE TITLE COMPOUND

Bond	Lengths	Bond	Lengths	Bond	Lengths
N(1)-C(6)	1.27(2)	C(6)-C(8)	1.45(2)	C(17)-C(19)	1.45(2)
N(1)-O(1)	1.357(16)	C(7)-C(8)	1.36(2)	C(18)-C(19)	1.42(2)
N(2)-C(17)	1.306(17)	C(7)-C(12)	1.44(2)	C(18)-C(23)	1.46(2)
N(2)-O(2)	1.358(15)	C(8)-C(9)	1.39(2)	C(19)-C(20)	1.35(2)
N(3)-O(3)	1.331(16)	C(9)-C(10)	1.35(2)	C(20)-C(21)	1.29(2)
N(4)-O(4)	1.378(16)	C(10)-C(11)	1.344(2)	C(21)-C(22)	1.38(2)
O(1)-C(1)	1.40(2)	C(11)-C(16)	1.374(2)	C(22)-C(23)	1.42(2)
O(2)-C(5)	1.444(19)	C(11)-C(12)	1.399(2)	C(22)-C(27)	1.45(2)
C(1)-C(2)	1.50(2)	C(12)-C(13)	1.36(2)	C(23)-C(24)	1.37(2)
C(2)-C(3)	1.47(2)	C(13)-C(14)	1.33(2)	C(24)-C(25)	1.323(2)
C(3)-C(4)	1.48(2)	C(14)-C(15)	1.41(2)	C(25)-C(26)	1.36(2)
C(4)-C(5)	1.48(2)	C(15)-C(16)	1.349(2)	C(26)-C(27)	1.46(2)
Bond	Angles	Bond	Angles	Bond	Angles
C(6)-N(1)-O(1)	112.5(17)	C(8)-C(7)-C(12)	120.3(17)	C(13)-C(14)-C(15)	115.0(18)
C(17)-N(2)-O(2)	110.5(14)	C(12)-C(7)-H(7)	119.9	C(16)-C(15)-C(14)	120.7(15)
C(33)-N(3)-O(3)	113.7(17)	C(7)-C(8)-C(9)	123.4(15)	C(15)-C(16)-C(11)	120.2(16)
C(44)-N(4)-O(4)	108.9(16)	C(7)-C(8)-C(6)	120(2)	N(2)-C(17)-C(19)	120.1(19)
N(1)-O(1)-C(1)	117.4(16)	C(9)-C(8)-C(6)	116(2)	C(19)-C(18)-C(23)	116.4(17)
N(2)-O(2)-C(5)	107.6(14)	C(10)-C(9)-C(8)	115.5(18)	C(20)-C(19)-C(18)	122.9(18)
N(3)-O(3)-C(28)	118.2(16)	C(11)-C(10)-C(9)	124.2(19)	C(20)-C(19)-C(17)	120(2)
N(4)-O(4)-C(32)	104.2(15)	C(10)-C(11)-C(16)	116(2)	C(18)-C(19)-C(17)	117(2)
O(1)-C(1)-C(2)	113.1(18)	C(10)-C(11)-C(12)	122.1(17)	C(21)-C(20)-C(19)	123(2)
C(3)-C(2)-C(1)	114.2(19)	C(16)-C(11)-C(12)	121.7(16)	C(20)-C(21)-C(22)	117(2)
C(2)-C(3)-C(4)	115.9(19)	C(13)-C(12)-C(11)	113.5(16)	C(21)-C(22)-C(23)	127.0(19)
C(5)-C(4)-C(3)	118.3(19)	C(13)-C(12)-C(7)	132(2)	C(21)-C(22)-C(27)	114.4(19)
O(2)-C(5)-C(4)	110.6(17)	C(11)-C(12)-C(7)	114.5(16)	C(23)-C(22)-C(27)	118.6(16)
N(1)-C(6)-C(8)	124(2)	C(14)-C(13)-C(12)	129(2)	C(24)-C(23)-C(22)	122.4(19)

Fig. 1. Molecule structure of the title compound with atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 30 % probability level

Fig. 2. Packing arrangement of the unit cell of the title compound. H atoms are omitted for clarity

Fig. 3. Dipolymer structure of the compound, hydrogen atoms, except those forming hydrogen bonds, are omitted for clarity

REFERENCES

1. S. Akine, A. Akimoto, T. Shiga, H. Oshio and T. Nabeshima, *Inorg. Chem.*, **47**, 875 (2008).
2. W.K. Dong, Y.X. Sun, Y.P. Zhang, L. Li, X.N. He and X.L. Tang, *Inorg. Chim. Acta*, **362**, 117 (2009).
3. W.K. Dong, Y.X. Sun, Y.X. Zhao, X.Y. Dong and L. Xu, *Polyhedron*, **29**, 2087 (2010).
4. J.P. Costes, F. Dahan and A. Dupuis, *Inorg. Chem.*, **39**, 165 (2000).
5. S.S. Sun, C.L. Stern, S.T. Nguyen and J.T. Hupp, *J. Am. Chem. Soc.*, **126**, 6314 (2004).
6. S. Yamada, *Coord. Chem. Rev.*, **190**, 537 (1999).
7. A.K. Sharma, F. Lloret and R. Mukherjee, *Inorg. Chem.*, **46**, 5128 (2007).
8. C. Polcar, F. Lambert and M. Cesario, *Eur. J. Inorg. Chem.*, **12**, 2201 (1999).
9. S. Akine, T. Taniguchi and T. Nabeshima, *Chem. Lett.*, 682 (2001).
10. S. Akine, T. Taniguchi and T. Nabeshima, *Inorg. Chem.*, **43**, 6142 (2004).
11. S. Akine, W.K. Dong and T. Nabeshima, *Inorg. Chem.*, **45**, 4677 (2006).
12. W.K. Dong, J.G. Duan and G.L. Liu, *Transition Met. Chem.*, **32**, 702 (2007).
13. W.K. Dong, C.E. Zhu, H.L. Wu, Y.J. Ding and T.Z. Yu, *Synth. React. Inorg. Met.-Org. Nano-Met. Chem.*, **37**, 61 (2007).
14. W.K. Dong, J.H. Feng, L. Wang and L. Xu, *Transition Met. Chem.*, **32**, 1101 (2007).
15. W.K. Dong, J. Yao, Y.X. Sun, L. Li and J.C. Wu, *Synth. React. Inorg. Met.-Org. Nano-Met. Chem.*, **40**, 521 (2010).