Antiviral Fluorenone Derivatives from *Arundina grammifolia*

DE-YUN NIU^{1,2}, JING-MEI HAN², WEI-SONG KONG², ZHU-WEN CUI², QIU-FEN HU¹ and XUE-MEI GAO^{1,*}

¹Key Laboratory of Chemistry in Ethnic Medicinal Resources, State Ethnic Affairs Commission & Ministry of Education, Yunnan University of Nationalities, Kunming 650031, P.R. China

²Key Laboratory of Tobacco Chemistry of Yunnan Province, Yunnan Academy of Tobacco Science, Kunming 650106, P.R. China

*Corresponding authors: E-mail: gao_xuemei@hotmail.com

(Received: 31 December 2012;

Accepted: 1 October 2013)

AJC-14217

A new fluorenone derivative, gramniphinol H (**1**), together with four known fluorenone derivatives (**2-5**) were isolated from the whole plant of *Arundina grammifolia*. Their structures were determined by means of HRESIMS, extensive 1D and 2D NMR spectroscopic studies and chemical evidence. Compound **1** was tested for its cytotoxicity against five human tumor cell lines and it shows modest cytotoxicity against PC3 and SHSY5Y cell with IC₅₀ values of 6.18 and 4.25 μ M, respectively.

Key Words: *Arundina grammifolia*, Fluorenone, Structural elucidation, Cytotoxicity.

INTRODUCTION

Arundina grammifolia (D. Don) Hochr. (bamboo orchid) belongs to the orchid family (Orchidaceae). The plant is used in Chinese folkloric medicine as a detoxifying and diuretic agent, as well as for the treatment of arthritis and inflammation¹. Previous phytochemical studies on *A. grammifolia* has revealed the presence of stilbenoids²⁻⁴, sterols^{5,6}, triterpenes^{7,8} and phenols⁵⁻¹⁰.

In our previous studies, some new phenols possessing antitobacco mosaic virus (anti TMV) property were isolated from *A. grammifolia*^{9,10}. Continuing the efforts to discover bioactive metabolites from local plants, we now investigated the chemical constituents of the whole plant of *A. grammifolia* growing in the Xishuangbanna Prefecture, leading to the isolation of a new fluorenone derivative (**1**) and four known fluorenone derivatives (**2-5**). The structures of the isolated compounds were determined by means of spectroscopic methods including 1D and 2D NMR techniques. Compound **1** was tested for its cytotoxicity against five human tumor cell lines and it shows modest cytotoxicity against PC3 and SHSY5Y cell.

EXPERIMENTAL

General procedures: Optical rotations were measured in a Horiba SEPA-300 polarimeter. UV spectra were obtained on a Shimadzu UV-2401A spectrophotometer and CD spectra were measured on a JASCO J-810 spectropolarimeter. Tenor 27 spectrophotometer was used for scanning IR spectra (KBr

pellets). 1D and 2D NMR spectra were recorded on a DRX-500 spectrometer with TMS as internal standard. Chemical shifts (δ) were expressed in ppm with reference to TMS. HRESIMS was performed on an API QSTAR spectrometer or a VG Autospec-3000 spectrometer. Preparative HPLC was performed on a Shimadzu LC-8A liquid chromatography equipped with ZORBAX PrepHT GF (21.2 mm $\times$ 25 cm, 7 μ m) column or Venusil MP C₁₈ (20 mm $\times$ 25 cm, 5 μ m) column. Column chromatography was performed using Si gel (200-300 mesh, Qing-dao Marine Chemical, Inc., Qingdao, China), Lichroprep RP-18 gel (40-63 μ m, Merck, Darmstadt, Germany), Sephadex LH-20 (Sigma-Aldrich, Inc, USA) or MCI gel (75-150 μ m, Mitsubishi Chemical Corporation, Tokyo, Japan). Column fractions were monitored by TLC and the spots were visualized by heating the plates after spraying with 5 % H₂SO₄ in EtOH.

The whole plant of *Arundina grammifolia* (D. Don)Hochr. was collected in the Xishuangbanna prefecture of Yunnan Province in September 2010. The identification of plant material was verified by Prof. Ning Yuan. A voucher specimen (Ynni-10-09-22) has been deposited in the Key Laboratory of Chemistry of Ethnic Medicinal Resources, Yunnan University of Nationalities.

Extraction and isolation: The air-dried and powdered of whole plant of *A. grammifolia* (2.0 kg) were extracted three times with 70 % aqueous acetone (3.0 L $\times$ 3.5 L) at room temperature and filtered to yield a filtrate, which was successively evaporated under reduced pressure to obtain a crude extract (152 g). The extract was applied to silica gel (150-200

mesh) column chromatography, eluting with CHCl_3 -MeOH gradients (20:1, 9:1, 8:2, 7:3, 6:4, 5:5) to afford fractions A-F. Further separation of fraction B (35.4 g) by silica gel (300-400 mesh), eluted with CHCl_3 -CO(CH₃)₂ (9:1-1:2), yielded fractions B1-B7. Fraction B3 (5.72 g), upon further separation on silica gel using petroleum ether-CH₃COOC₂H₅ and semi-preparative HPLC (55 % MeOH-H₂O, flow rate 12 mL/min), afforded **1** (18.2 mg), **2** (16.7 mg), **3** (21.4 mg) **4** (28.2 mg) and **5** (29.7 mg).

Gramniphénol H (1): Red gum; $[\alpha]_D^{25.0}$ -16.8 (c 0.25, CH₃OH); UV (CH₃OH) λ_{max} (log ϵ) 210 (3.72), 267 (3.88), 320 (2.52), 346 (2.71) nm; IR (KBr, ν_{max} , cm⁻¹): 3325, 2970, 2892, 1694, 1612, 1547, 1455, 1184, 1119; ¹C and ¹³H NMR data (C₅D₅N, 500 and 125 MHz) see Table-1; negative ESIMS m/z 327 [M-H]⁻; negative HRESIMS m/z 327.0862 [M-H]⁻ (calcd. (%) for C₁₈H₁₅O₆, 327.0869).

TABLE-1

¹H AND ¹³C NMR DATA OF COMPOUND **1** (δ IN ppm, IN C₅D₅N)

Position	Compound 2	
	δ_C	δ_H (J in Hz)
1	150.8 s	—
2	118.2 d	6.71 d (8.8)
3	129.5 d	7.06 d (8.8)
4	144.4 s	—
5	156.6 s	—
6	106.9 d	6.91 s
7	154.5 s	—
8	108.1 d	6.77 s
9	192.9 s	—
4a	125.0 s	—
4b	120.6 s	—
8a	137.1 s	—
9a	116.8 s	—
1'	71.0 t	3.97 dd (6.9, 10.0)
—	—	4.13 dd (9.5, 4.0)
2'	74.1 d	4.36 m
3'	145.8 s	—
4'	113.4 t	4.89 s
—	—	5.18 s
5'	19.0 q	1.68 s
Ar-OH-1	—	11.02 s
Ar-OH-2	—	10.90 s
Ar-OH-4	—	10.50 s

RESULTS AND DISCUSSION

Gramniphénols H (**1**) and four known fluorenone derivatives (**2-5**) (Fig. 1) were isolated from the 70 % aqueous acetone extract of *A. grammifolia*. Known compounds were identified as 1,4,5-trihydroxy-7-methoxy-9H-fluoren-9-one (**2**)¹¹, dendroflorin (**3**)¹², Dengibsin (**4**)¹³, Denchrysan A (**5**)¹⁴ by comparison of spectroscopic data with the literature.

Fig. 1. Structures of derivatives fluorenone from *A. grammifolia*

Gramniphénol H (**1**) was obtained as red gum. The molecular formula C₁₈H₁₅O₆ was inferred by HRESI-MS at m/z 327.0862 [M-H]⁻ (calcd. (%) for C₁₈H₁₅O₆, 327.0869). The IR absorption bands indicated the presence of hydroxy (3325 cm⁻¹), carbonyl (1694 cm⁻¹) and aromatic ring (1612, 1547, 1455 cm⁻¹) and UV absorptions at 267, 320, 346 nm suggested a conjugated aromatic ring system. The ¹H and ¹³C NMR spectra of **1** (Table-1) displayed signals for all **18** carbons and **16** protons, suggesting the presence of an 1,4,5,7-oxygenated fluorenone moiety¹¹ (H-2, H-3, H-6, H-8 and C-1-C-9a), three phenolic protons (δ_H 10.50, 10.90 and 11.02) and a 2-hydroxy-3-methylbut-3-enyloxy unit [-OCH₂CH(OH)C(CH₃)(CH₃)]¹⁰ (H-1', H-2', H-4', H-5' and C-1'-C-5'). The location of the 2-hydroxy-3-methylbut-3-enyloxy unit at C-5 position was supported by the HMBC correlation observed between H-1' (δ_H 3.97 and 4.13) and C-5 (δ_C 156.6). The three phenolic groups were assigned to C-1, C-4 and C-7 positions on the basis of HMBC correlations (Fig. 2). The S configuration at C-2' was assigned by a comparison of NMR and optical rotation data with those of stachyline A¹⁰, of which absolute configuration was established by the Mosher method.

Fig. 2. Selected HMBC (—) correlations of compound **1**

Since certain of the phenolic compounds exhibit potential cytotoxicity¹⁵⁻¹⁷. Compound **1** was tested for their cytotoxicity against five human tumor cell lines (NB4, A549, SHSY5Y, PC3 and MCF7) using the MTT method as reported previously¹⁸. Taxol was used as the positive control. The results revealed that compound **1** showed low active (IC₅₀ values >10 μ M) for NB4, A549 and MCF7 tumor cell and showed modest cytotoxicity against PC3 and SHSY5Y cell with IC₅₀ values of 6.18 and 4.25 μ M, respectively.

ACKNOWLEDGEMENTS

This project was supported financially by the Excellent Scientific and Technological Team of Yunnan Higher Education (2010CI08) and the Yunnan University of Nationalities Green Chemistry and Functional Materials Research for Provincial Innovation Team (2011HC008).

REFERENCES

- D.Y. Hong, Y. Lian, S. Shen and L.D., Flora of China; Chinese Science Press: Beijing, Vol. 73, p. 320 (1983).
- K. Xiao, H.J. Zhang, L.J. Xuan, J. Zhang, Y.M. Xu and D.L. Bai, *Studies Nat. Prod. Chem.*, **34**, 453 (2008).
- M.F. Liu, Y. Ding and D.M. Zhang, *Chin. J. Chin. Mater. Med.*, **30**, 353 (2005).
- M.F. Liu, Y. Han, D.M. Xing, Y. Shi, L.Z. Xu, L.J. Du and Y. Ding, *J. Asian Nat. Prod. Res.*, **6**, 229 (2004).

5. M.F. Liu, Y. Ding and L. Du, *J. Chin. Trad. Herb Drug*, **38**, 676 (2007).
6. H. Zhu and Q.S. Song, *Nat. Prod. Res. Dev.*, **20**, 5 (2008).
7. A.S.C. Wan, R.T. Aixel and H.J. Nicholas, *Phytochemistry*, **10**, 2267 (1971).
8. P.L. Majumder and S. Ghosal, *J. Indian Chem. Soc.*, **68**, 88 (1991).
9. Z.R. Gao, S.T. Xu, J. Wei, H.L. Shi, Z. Li and Q.F. Hu, *Asian J. Chem.*, **25**, 2747 (2013).
10. Q.F. Hu, B. Zhou, J.M. Huang, X.M. Gao, L.D. Shu, G.Y. Yang and C.T. Che, *J. Nat. Prod.*, **75**, 684 (2013).
11. Y.G. Chen, Y.P. Li, C. Qing, Y.L. Zhang, L.Q. Wang and Y. Liu, *Food Chem.*, **108**, 973 (2008).
12. C.O. Fan, W. Wang, Y.P. Wang, G. Qin and W.M. Zhao, *Phytochemistry*, **57**, 1255 (2001).
13. S.K. Talapatra, S. Bose, A.K. Mallik and B. Talapatra, *Tetrahedron*, **41**, 2765 (1985).
14. Q.H. Ye, W.M. Zhao and G.W. Qin, *Nat. Prod. Res.*, **17**, 201 (2003).
15. Y.L. Lu, X.S. Li, H.X. Mu, H.T. Huang, G.P. Li and Q.F. Hu, *J. Braz. Chem. Soc.*, **23**, 656 (2012).
16. X.M. Gao, Y.Q. Shen, L.Y. Yang, L.D. Shu, G.P. Li, and Q.F. Hu, *J. Braz. Chem. Soc.*, **23**, 1274 (2012).
17. Q.F. Hu, X.S. Li, H.T. Huang, H.X. Mu, P.F. Tu and G.P. Li, *Helv. Chim. Acta*, **95**, 349 (2012).
19. T. Mosmann, *J. Immunol. Methods*, **65**, 55 (1983).

THE INTERNATIONAL CONFERENCE ON PURE AND APPLIED CHEMISTRY (ICPAC-2014)

23 — 27 JUNE, 2014

HOTEL SOFITEL MAURITIUS IMPERIAL RESORT & SPA, MAURITIUS

(<http://sites.uom.ac.mu/icpac/>)

Theme of ICPAC-2014 is Crystallising Ideas: The Role of Chemistry

For abstract submission: <http://sites.uom.ac.mu/icpac/index.php/abstract>

*Full papers (optional) will be peer reviewed and accepted papers will be collected in the book of proceedings to be published by Springer.

Deadlines

Workshop/Symposium proposal	31st December 2013
Abstract submission (oral or poster)	15th March 2014
Booking for accommodation	1st April 2014
Early Registration	15th April 2014
Full paper (optional)	1st October 2014