Synthesis of Novel γ -Ketoesters from Succinic Anhydride

MUHAMMAD IQBAL*, IMAM BAKHSH BALOCH and MUSA KALEEM BALOCH

Department of Chemistry, Gomal University, Dera Ismail Khan-29050, Khyber Pakhtunkhwa, Pakistan

*Corresponding author: E-mail: miqbal_44@yahoo.com

(Received: 21 January 2013;

Accepted: 16 October 2013)

AJC-14257

Four alkyl γ -ketoheptanoates (**3a-3d**) have been prepared from succinic anhydride employing a three step reaction strategy. In the first step using *p*-toluene sulfonic acid as catalyst, the ring of succinic anhydride was opened with isopropyl, isobutyl, isopentyl and benzyl alcohols, respectively to form alkyl hydrogen succinates (**1a-1d**). In the 2nd step these alkyl hydrogen succinates on treatment with SOCl_2 yielded 4-alkoxy-4-ketobutanoyl chlorides (**2a-2d**). The acid halides thus obtained, on reaction with diethyl cadmium led to require γ -ketoesters. All the synthesised compounds were characterized by recording and analyzing ^1H , ^{13}C NMR, IR spectra and mass measurements.

Key Words: γ -Ketoesters, Isopentyl γ -ketoheptanoate, Succinic anhydride, Isobutyl alcohol.

INTRODUCTION

γ -Keto-esters are used in the preparation of lactam antibiotics¹, isoquinolines¹, lactonic sex pheromones^{1,2}, furans^{1,3}, benzotropolones⁴, alkoxythiophenes⁵ and other heterocycles⁶. They are used as intermediates in the preparation of cyclopentenone³, natural flavor² and useful building blocks of pharmaceutical interest³. Moreover, they are converted to butyrolactone, its dimer⁷, chiral γ -hydroxyester^{8,9}, γ -hydroxycarboxylic acids⁶ and chiral auxiliaries in asymmetric synthesis^{10,11}. Although many methods have been developed for γ -ketoesters synthesis¹²⁻²⁴, they often have disadvantages, such as poor yields, expensive starting materials, sensitivity of the esters to hydrolysis in acidic or basic medium and use of expensive catalysts like NaAuCl_4 ²³, $\text{Co}(\text{OAc})_2$ ¹ and $\text{Rh}_2(\text{OAc})_4$ ². Our objective was thus to develop a new route for the syntheses of γ -ketoesters (**3a-3d**) using inexpensive substrates, solvents and catalysts and to optimize the yield of the products.

EXPERIMENTAL

All the chemicals were of analytical grade, purchased from E. Merck, Darmstadt, Germany and used as acquired; toluene was dried and stored over sodium metal before use. The purity of the alcohols and succinic anhydride was checked by NMR and IR whereas the purity of thionyl chloride was checked by IR. All the reactions were carried out in an efficient fume hood under dry nitrogen. The chemical splash goggles, Nomex gloves and lab coat were used to minimize the exposure to diethyl cadmium. Melting points were determined using a Gallenkamp digital melting point apparatus and are uncor-

rected. The UV spectrum was recorded in absolute methanol on a IRMECO UV/VIS Model U-2020 spectrophotometer. IR spectra were obtained on a TENSOR 27 FT-IR spectrophotometer supplied by Bruker, Ettlingen, Germany. ^1H and ^{13}C NMR (1D, 2D, homo/heteronuclear) spectra were acquired in CDCl_3 on a Bruker Biospin, AMX 300 MHz FT NMR (300 MHz for ^1H , 75 MHz for ^{13}C) spectrometer. EI-MS were determined with a direct insertion probe on a double-focusing Finnigan MAT 112 at 70 eV. HR-EI-MS measurements were carried out on a JEOL HX 110 spectrometer. Elemental analyses were obtained by Midwest Microlab, Indianapolis, IN. Column chromatography was performed using silica gel (PF₂₅₄, mesh size 60-70), E. Merck, Darmstadt, Germany and thin layer chromatography on pre-coated silica gel plates (20 cm $\times$ 20 cm, 0.2 mm thickness) with UV fluorescence (PF₂₅₄), E. Merck, Darmstadt, Germany; eluting with hexane-EtOAc.

Alkyl hydrogen succinates (1a-1d). General procedure: To a single-necked round-bottom flask (100 mL), fitted with a Dean-Stark trap and a reflux condenser and containing a stir bar, succinic anhydride (3.0 g, 30 mmol), anhydrous *p*-toluenesulfonic acid (25 mg) and toluene (15 mL) under nitrogen were added 30 mmol of the alcohols. After 24 h at reflux, the solution was left to cool at 25 °C then poured into a saturated aqueous NaHCO_3 solution (25 mL) and the aqueous layer was extracted with hexane (3 $\times$ 25 mL). The combined organic (toluene and hexane) layers were washed with brine (20 mL), dried over anhydrous Na_2SO_4 and the solvent was removed under vacuum to give a resinous product, which was subjected to column chromatography to afford the pure monoesters.

TABLE-1
 YIELD, MELTING POINTS AND ELEMENTAL ANALYSIS OF COMPOUNDS 1-2

Compound	Yield (%)	m.p. (°C)	b.p./18 mm (°C)	Elemental analysis (%) found (calcd.)	
				C	H
1a	76	64-65	—	52.49 (52.52)	7.55 (7.56)
1b	77	69-71	—	55.16 (55.10)	8.10 (8.11)
1c	76	75-76	—	57.43 (57.45)	8.57 (8.58)
1d	76	115-116 (Lit.117-119) ²⁵	—	63.45 (63.47)	5.81 (5.75)
2a	78	—	94-95	—	—
2b	78	—	101-103	—	—
2c	72	—	108-109	—	—
2d	71	—	110-112	—	—

4-Alkoxy-4-ketobutanoyl Chlorides (2a-2d). Typical

procedure: Compound **1a** (3.6 g, 22.5 mmol) was mixed with thionyl chloride (3.26 mL, 45 mmol) and warmed at 30-40 °C for 3 h. Then the excess of thionyl chloride was removed on a steam bath under reduced pressure to afford a colourless resin (**2a**) which was purified by distillation at reduced pressure and characterized by physical and spectroscopic measurements (Tables 1 and 3).

Alkyl γ -ketoheptanoates (3a-3d). Typical procedure:

Compound **3a** was synthesized by addition of 10 mmol of **2a** (1.78 g) over a period of 10 min to a solution of diethyl cadmium in diethyl ether²⁷⁻²⁹. The mixture was refluxed for six hours and allowed to stand overnight stirring. The reaction mixture was then poured into a beaker (500 mL) containing crushed ice (150 g) and 10 % aqueous H₂SO₄ (30 mL). It was vigorously stirred using a magnetic stirrer for 5 min. The organic layer was extracted with diethyl ether (3 × 20 mL), washed with 50 % aqueous NaHCO₃ solution (40 mL) and dried over anhydrous Na₂SO₄. After removal of the solvent, the obtained material was subjected to column chromatography

 TABLE-2
 YIELD AND BPS OF COMPOUNDS 3a-d FOR COMPARISON

Compound	Yield (%)			b.p/18mm (°C)
	Diethyl cadmium	Gilman reagent	Grignard reagent	
3a	77	29	22	106-107
3b	74	28	21	112-113
3c	72	27	18	118-119
3d ²⁶	71	23	18	116-117

using an ethyl acetate/hexane [(1:9) → (2:8) → (3:7)] as eluent. The desired product **3a** was obtained after elution with ethyl acetate/hexane (3:7).

RESULTS AND DISCUSSION

Reaction of succinic anhydride with alcohols led to alkyl hydrogen succinates **1a-1d** which upon treatment with thionyl chloride afforded the corresponding 4-alkoxy-4-ketobutanoyl chlorides **2a-2d** (Scheme-I). Reaction of **2a-2d** with diethyl cadmium, ethylmagnesium bromide or the corresponding

 TABLE-3
 SPECTROSCOPIC DATA OF COMPOUNDS 1-3

Comp.	IR (cm ⁻¹)	¹ H NMR (δ)	¹³ C NMR (DEPT) (δ)	HR-EI-MS (m/z)	
				Calcd.	Found
1a	3417 (OH) 1728 (CO ₂ H), 1755 (CO ₂ R)	2.55 (t, 6.7), 2.72 (t, 6.7), 4.45 (septet, 7.5), 1.1 (d, 7.5)	173.3 (s), 28.2 (t), 36.6 (t), 180 (s), 68.3 (d) 21.2 (q)	160.17 C ₇ H ₁₂ O ₄	160.15
1b	3438 (OH) 1725 (CO ₂ H), 1759 (CO ₂ R)	2.55 (t, 6.7), 2.72 (t, 6.7) 4.01 (d, 7.5), 1.18 (septet, 7.5), 1.1 (d, 7.5)	173.3 (s), 28.2 (t), 36.6 (t), 180 (s), 68.3 (t), 26.7 (d), 21.2 (q)	174.19 C ₈ H ₁₄ O ₄	174.18
1c	3396 (OH), 1722 (CO ₂ H), 1765 (CO ₂ R)	2.55 (t, 6.7), 2.72 (t, 6.7) 4.45 (d, 7.5), 1.48 (m), 1.77 (m), 1.1 (d, 7.5)	173.3 (s), 28.2 (t), 36.6 (t), 180 (s), 68.3 (t), 36.7 (t), 23.6 (d), 21.2 (q)	188.22 C ₉ H ₁₆ O ₄	188.24
1d	3324 (OH), 3014 (Ar-H), 1748 (CO ₂ R), 1708 (CO)	2.57 (t, 6.7), 2.69 (t, 6.7) 5.2 (s), 7.2-7.3 (Ph)	171.7 (s), 28.3 (t), 35.5 (t), 203.2 (s), 79.1 (t), 128-137, (1C, s, 5C, d)	208.21 C ₁₁ H ₁₂ O ₄	208.19
2a	1810 (COCl), 1755 (CO ₂ R), 741 (C-Cl)	2.53 (t, 6.7), 2.73 (t, 6.7) 4.5 (septet, 7.5), 1.12 (d, 7.5)	171.4 (s), 28.4 (t), 37.5 (t), 172.6 (s), 66.2 (d), 20.1 (q)	178.61 C ₇ H ₁₁ ClO ₃	178.62
2b	1802 (COCl) 1740 (CO ₂ R), 734 (C-Cl)	2.56 (t, 6.7), 2.70 (t, 6.7) 4.05 (d, 7.5), 1.47 (m), 1.16 (d, 7.5)	172.2 (s), 27.0 (t), 37.8 (t), 205.2 (s) 64.5 (t), 31.7 (d), 22.1 (q)	192.64 C ₈ H ₁₃ ClO ₃	192.65
2c	1807 (COCl) 1739 (CO ₂ R), 739 (C-Cl)	2.58 (t, 6.7), 2.74 (t, 6.7) 4.1 (t, 7.5), 1.6 (dt, 7.5), 1.5 (m), 1.1 (d, 7.5)	171.1 (s), 27.5 (t) 36.9 (t), 172.3 (s), 59.9 (t), 37.2 (t) 25.5 (d), 22.2 (q)	206.67 C ₁₁ H ₂₀ O ₃	206.66
2d	3094 (Ar-H), 1787 (COCl), 1748 (CO ₂ R) 734 (C-Cl)	2.57 (t, 6.7), 2.69 (t, 6.7), 5.2 (s), 7.2-7.3 (Ph)	171.7 (s), 28.3 (t) 35.5 (t), 176.7 (s), 79.1(t), 128-137 (1C, s, 5C, d)	226.66 C ₁₁ H ₁₁ ClO ₃	226. 67
3a	1745 (CO ₂ R), 1729 (CO), 1210 (C-O)	2.57 (t, 6.7), 2.71 (t, 6.7) 2.49 (t, 7.3), 1.08 (t, 7.3), 4.52 (septet, 7.5), 1.15 (d, 5)	171.2 (s), 27.9 (t), 35.6 (t), 204.5 (s), 34.7 (t), 7.9 (q) 63.7 (d), 22.6 (q)	172.22 C ₉ H ₁₆ O ₃	172.21
3b	1740 (CO ₂ R), 1722 (C=O), 1220 (C-O)	2.56 (t, 6.7), 2.70 (t, 6.7) 2.47 (t, 7.3), 1.07 (t, 7.3), 4.05 (d, 7.5), 1.47(m), 1.16 (d, 7.5)	172.3 (s), 27.0 (t), 37.8 (t), 205.2 (s), 35.2 (t), 8.3 (q), 64.5 (t), 31.7 (d)	186.23 C ₁₀ H ₁₈ O ₃	186.24
3c	1739 (CO ₂ R), 1717 (C=O), 1225 (C-O)	2.58 (t, 6.7), 2.74 (t, 6.7), 2.51 (q, 7.3), 1.09 (t, 7.3), 4.1 (t, 7.5), 1.6 (dt, 7.5), 1.5 (septet, 7.5), 1.1 (d, 7.5)	173.2 (s), 27.5 (t), 36.9 (t), 202.3 (s) 35.9 (t), 7.7 (q), 59.9 (t), 37.2 (t), 25.5 (d), 22.2 (q)	200.27 C ₁₁ H ₂₀ O ₃	200. 26
3d ²⁶	1736 (CO ₂ R), 1718 (C=O), 1230 (C-O)	1.05 (t, J = 7.3 Hz, 3 H), 2.48 (q, J = 7.3 Hz, 2 H), 2.65 (t, J = 6.7 Hz, 2 H), 2.74 (t, J = 6.7 Hz, 2 H), 5.15 (s, 2 H), 7.33–7.35 (m, 5 H)	171.7 (s), 28.3 (t), 35.5 (t), 208.2 (s), 36.1 (t), 7.9 (q), 66.1 (t), 128-137, (1C, s, 5C, d)	220.26 C ₁₃ H ₁₆ O ₃	220.24

Gilman reagent converted these compounds to alkyl γ -ketoheptanoates **3a-3d** (Scheme-I, Table-1).

Scheme-I: Preparation of compounds **1-3**

Comparison of the results with the three organometallic reagents is illustrated in Table-2. The yield was highest utilizing the cadmium reagent and no side products were formed. The EI-MS spectrum of the compound showed peak at m/z 57 and was attributed to the loss of keto group from the parent molecule.

The ¹H NMR spectrum of **3a** displayed two new peaks at δ 1.08 (t, J = 7.3 Hz, 3H, Me) and at δ 2.49 (q, J = 7.3 Hz, 2H, CH₂CO). Its ¹³C NMR spectrum exhibited two extra peaks as compared to **2a** that were identified as CH₃ and CH₂ by DEPT-135 ¹³C NMR. The 2D NMR also confirmed the formation of γ -ketoester (**3a**). The physical, spectroscopic and the literature data revealed that isobutyl hydrogen succinate (**1b**) and isopentyl hydrogen succinate (**1c**) have been identified in wine³⁰. Benzyl hydrogen succinate (**1d**), m.p. 117-119 °C²⁵ has been synthesized in 52 % yield by exposure of benzyl β -formyl propionate to air for 30 days. In the series **3a-d** only benzyl γ -ketoheptanoate (**3d**) is known²⁶. This compound was prepared²⁶ in low yield (24 %) by reaction of benzyl acrylate with excess propanoyl chloride in the presence of Mg and in DMF as solvent. Our methodology gave **3d** in 71 % overall yield. The spectroscopic data of **3d** are identical with that reported in the literature²⁶.

REFERENCES

1. D. Huang, M. Yan, W.-J. Zhao and Q. Shen, *Synth. Commun.*, **35**, 745 (2005) and reference cited therein.
2. E.V. Starodubtseva, O.V. Turova, M.G. Vinogradov, L.S. Gorshkova, V.A. Ferapontov and M.I. Struchkova, *Tetrahedron*, **64**, 11713 (2008) and references cited therein.

3. I. Ryu, K. Matsumoto, M. Ando, S. Murai and N. Sonoda, *Tetrahedron Lett.*, **21**, 4283 (1980).
4. A. Kamal, M. Sandbhor and A.A. Shaik, *Tetrahedron: Asymm.*, **14**, 1575 (2003).
5. V.M. Sonpatki, M.R. Herbert, L.M. Sandvoss and A.J. Seed, *J. Org. Chem.*, **66**, 7283 (2001).
6. F. Csende and G. Stajer, *Heterocycles*, **53**, 1379 (2000).
7. D.B.G. Williams, K. Blann, J. Caddy and C.W. Holzappel, *Synth. Commun.*, **32**, 3755 (2002).
8. T. Benincori, S. Rizzo, T. Pilati, A. Ponti, M. Sada, E. Pagliarini, S. Ratti, C. Giuseppe, L. de Ferraf and F. Sanniccolo, *Tetrahedron: Asymm.*, **15**, 2289 (2004).
9. D.B.G. Williams, K. Blann and C.W. Holzappel, *Synth. Commun.*, **31**, 203 (2001).
10. D. Romo and A.I. Meyers, *Tetrahedron*, **47**, 9503 (1991).
11. M. Utaka, H. Watabu and A. Takeda, *J. Org. Chem.*, **52**, 4363 (1987).
12. A. Giardina, E. Marcantoni, T. Mecozzi and M. Petrini, *Eur. J. Org. Chem.*, **2001**, 713 (2001).
13. H. Stetter, *Angew. Chem. Int. Ed. Engl.*, **15**, 639 (1976).
14. C. Kashima, Y. Shirahata and Y. Tsukamoto, *Heterocycles*, **54**, 309 (2001).
15. M. Yan, W.-J. Zhao, D. Huang and S.-J. Ji, *Tetrahedron Lett.*, **45**, 6365 (2004).
16. R. Ballini and M. Petrini, *Tetrahedron*, **60**, 1017 (2004).
17. A.P. Cowling and J. Mann, *J. Chem. Soc. Chem. Commun.*, 1006 (1978).
18. R.C. Ronald and C.J. Wheeler, *J. Org. Chem.*, **48**, 138 (1983).
19. W.J. Coates and A. McKillop, *Synthesis*, 334 (1993).
20. W.D. Woessner and P.S. Solera, *Synth. Commun.*, **8**, 279 (1978) and references cited therein.
21. P.A. Wehrli and V. Chu, *J. Org. Chem.*, **38**, 3436 (1973).
22. R.P. Singh, J.M. Leitch, B. Twamley and J.M. Shreeve, *J. Org. Chem.*, **66**, 1436 (2001).
23. W. Wang, B. Xu and G.B. Hammond, *J. Org. Chem.*, **74**, 1640 (2009).
24. A.K. Ghosh and L. Swanson, *J. Org. Chem.*, **68**, 9823 (2003).
25. J.G. Cannon and J.E. Garst, *J. Org. Chem.*, **40**, 182 (1975).
26. M. Matsunami, N. Sakai, T. Morimoto, H. Maekawa and I. Nishiguchi, *Synlett*, 769 (2007).
27. D. Buser, H. Pauling, A. Thum and W. Bonrath, *Molecules*, **7**, 341 (2002).
28. H.O. Hankovszky, K. Hideg, L. Lex, G. Kulcsar and H.A. Halasz, *Can. J. Chem.*, **60**, 1432 (1982).
29. H. Gilman and F. Schulze, *J. Am. Chem. Soc.*, **47**, 2002 (1925).
30. L. Nynaken and H. Suomalainen, *Aroma of Beer, Wine and Distilled Alcoholic Beverages*, ISBN 90-277-1553-X, Kluwer Academic Publishers, p. 208 (1983).