Electrochemical Generation of Complexes Containing Heterometallic Trinuclear Triangular Metal-Sulphur-Cuboidal $\{M'M_2S_4\}^{4+}$ Core (where M and M' = Mo and W): Synthesis and Study of Physico-chemical Behaviour

ANAMIKA¹, RAMJEE SAH² and ASHWINI KUMAR^{3,*}

¹Department of Chemistry, Mahila College, Chaibasa-833 201, India

²University Department of Chemistry, B.R.A. Bihar University, Muzaffarpur-842 001, India

³Chemistry Department, Shyam Nandan Sahay College, Muzaffarpur-842 001, India

*Corresponding author: E-mail: ashwinikumar.chem@gmail.com

(Received: 5 February 2013;

Accepted: 14 November 2013)

AJC-14386

Chromatographic separation of mixture leads to the isolation of molybdenum-tungsten mixed-metal clusters $[MoW_2S_4(H_2O)_9]^{4+}$ and $[W_2MoS_4(H_2O)_9]^{4+}$. From the aqua ion the corresponding oxalato-derivatives as $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{2-}$ and $[W_2MoS_4(C_2O_4)_3(H_2O)_3]^{2-}$ are isolated. The peak positions of the two big bands in each infrared spectrum of the four aqua clusters, $[Mo_3S_4(H_2O)_9]^{4+}$, $[W_3S_4(H_2O)_9]^{4+}$, $[MoW_2S_4(H_2O)_9]^{4+}$ and $[W_2MoS_4(H_2O)_9]^{4+}$ in the 550-400 cm^{-1} region shift to lower wave number when the molybdenum atom is replaced by tungsten. The cyclic voltammograms of $Cs_2[MoW_2S_4(C_2O_4)_3(H_2O)_3]$ (**1**) and $Cs_2[W_2MoS_4(C_2O_4)_3(H_2O)_3]$ (**2**) shows three consecutive one-electron reduction steps in the aqueous medium with supporting electrolyte, KCl at -0.84, -1.40, -1.78 and -0.73, -1.22, -1.66 V, respectively. These steps correspond to the change of oxidation states of three metals in each cluster with the following electronic configuration $\{M_3\}^6 \rightarrow \{M_3\}^7 \rightarrow \{M_3\}^8 \rightarrow \{M_3\}^9$. The half wave potential $E_{1/2}$ are significantly dependent on the cluster metals. In all the reduction processes the $[Mo_nW_{3-n}S_4(C_2O_4)_3(H_2O)_3]^{2-}$ clusters ($n = 0-3$) are easily reduced with the increase in the number of Mo(n) in the clusters. The proceeding reduction centre of the molybdenum-tungsten mixed metal clusters is mainly on the Mo atom(s) rather than W atoms.

Key Words: Electrochemistry, Sulphur-bridged, Incomplete Cubanes-type, Heterometallic cuboidal, Core.

INTRODUCTION

The structure and function of the various molybdoenzyme continues to much attention. Recent years have seen a dramatic expansion in the synthetic and structural molybdenum-sulphur as well as tungsten-sulphur chemistry. It is well known that molybdenum has an extraordinary proclivity to form binuclear and cluster compounds containing terminal μ_2 - or μ_3 -sulphur atoms, some of which are of biological importance¹. Multinuclear transition metal-sulphur complexes are of well-established significance in bio-chemistry, while extended-lattice transition metal-sulphur solids *i.e.*, MoS_2 are ubiquitous in industrial hydro treating catalysis. Many reports have been made on the trinuclear triangular tungsten and also molybdenum compounds in which metal atoms are bridged by several kinds of atoms or ligands like O, S, S, NH and $Cl^{2,3}$. Molybdenum and tungsten sulphur compounds have attracted much attention and a large number of sulphur-bridged molybdenum and tungsten compounds have appeared.

It has recently been reported that the reaction of the incomplete cubane-type Mo_3S_4 aqua ion with metallic iron,

cobalt or nickel gives the corresponding heterometallic cubane-type $[M'Mo_3S_4]^{4+}$ aqua ion where $M' = Fe, Co, Ni$. However, the number of triangular trinuclear heterometallic molybdenum-tungsten mixed metal compounds are limited⁴⁻⁷. $[Mo_2WS_4(C_5(CH_3)_3)]$ reported by Wachter and co-workers⁸, is the only cluster to have the molybdenum-tungsten mixed-metal core, though some-metal clusters with incomplete cubane-type $M_2M'S_4$ cores ($M = Mo$ or W ; $M' =$ other metal) are reported⁹⁻¹⁵.

The author reports the synthesis and electrochemical behaviour of incomplete cubane-type molybdenum(IV)-tungsten(IV) mixed-metal aqua cluster compounds and their corresponding oxalato derivatives, *i.e.*, $Cs_2[MoW_2S_4(C_2O_4)_3(H_2O)_3]$ and $Cs_2[W_2MoS_4(C_2O_4)_3(H_2O)_3]$.

Investigation have been made to find out the interaction between molybdenum and tungsten and also to compare the character of the elements in the incomplete cubane-type clusters with $[W_nMo_{3-n}S_4]^{4+}$ cores ($n = 0-3$).

EXPERIMENTAL

IR spectra of these complexes recorded on Perkin-Elmer model 580 infrared spectrophotometer. The samples were

prepared as CsI pellets and referenced to polystyrene bands. The electronic spectra of the complexes were recorded on Shimadzu. UV-visible 190 and Perkin-Elmer λ_2 double beam spectrophotometers using matched quartz cell.

The cyclic voltammograms were recorded on a Bioanalytical system CV-27 voltammogram, in the connection with a C1B-Cell stand. All the experiments were performed in a standard three electrode configuration using an auxiliary Pt- electrode, a glassy carbon working electrode and a saturated calomel electrode under dry nitrogen atmosphere. All the results were collected at 298 K and referenced to saturated calomel electrode (SCE) was used as supporting electrolyte for aqueous medium. The reported values were uncorrected for junction potential where voltammograms were recorded in different scan speeds but with the scan speed at 100mV/s was used for discussion.

All the reagents and solvents were of A.R. grade and were used without further purification. Water was distilled in quick-fit apparatus over alkaline $KMnO_4$ and H_2SO_4 . The electric conductance of distilled water varied between 7×10^{-7} to $9 \times 10^{-9} W^{-1} cm^{-1}$. Elemental analyses were performed by EA-1108 elemental analyzer. Sulphur and molybdenum were analyzed by the oxidizing the compound by peroxide fusion as $BaSO_4$ and molybdenum oxinate, respectively. *p*-Toluene sulfonic acid (HPTS) was used after recrystallization. Sodium borohydride, oxalic acid, CsCl as well as most other reagents were commercial grade and were used as received. $(NH_4)_2MoS_4$ and $(NH_4)_2WS_4$ were synthesized by published procedure.

Synthesis of compounds

$Cs_2[MoW_2S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$ (1): $(NH_4)_2WS_4$ (3.48 g, 10 mmol) was dissolved in water (200 mL). $(NH_4)_2S_2O_8$ (1.14 g, 5 mmol) was added to the solution and the reaction mixture was allowed to stir constantly for an hour at room temperatures. A prepared solution dissolving $(NH_4)_2MoS_4$ (1.3 g, 5 mmol) in water (50 mL) was mixed to the constantly stirred reaction mixture of $(NH_4)_2WS_4$. The entire reaction mixture was allowed to reduce heterogeneously by adding $NaBH_4$ (5 g, 0.125 mol) portionwise (*ca.* 0.5 g at each interval) and 50 mL of 6 M HCl acid (using 5 mL at each interval) alternatively. The reduction of reaction mixture was continued for 2 h. After the consumption of reducing agent an additional amount 250 mL of 2 M HCl acid was added to the solution, which was heated at 90 °C for 6 h with constant introduction of an air stream. During the course of heating 1 M HCl was added occasionally to keep the volume of the solution constant (300 mL). The solution was allowed to cool at room temperature and filtered under suction.

The filtrate was subjected for chromatographic separation using Sephadex G-15 column (4.0 cm $\times$ 100 cm, 1 M HCl). The gray coloured bands were collected and concentrated by the use of a cation exchange, Dowex 50 W-X₈ (2 M HCl). Sephadex G-15 column chromatography (2.0 cm $\times$ 150 cm, 1 M HCl) was applied again for purification. This purification for concentration process was repeated again. The dark grey band appeared at the top of column was eluted finally by using oxalic acid solution (350 mg in 20 mL water). To this elute, cesium chloride (168 mL, 1 mmol) was added and then the resultant solution was allowed to evaporate under conc. H_2SO_4

acid desiccators. The initial white precipitate was filtered and from the mother liquor dark black plate shaped crystals were separated out, that was filtered, washed with minimum amount of cold water, then with methanol and finally with ether and then dried under vacuum to yield with 75 % of the product based on the starting material. Anal: found (calcd. (%)) for $C_6H_{10}O_{17}S_4MoW_2Cs_2$: C: 6.02 (5.94); H: 0.84 (0.8250); S: 10.12(10.56); Mo: 7.80 (7.92); W: 30.40 (30.36).

$Cs_2[WMo_2S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$ (2): The entire synthetic procedure was followed exactly similar to that used for the preparation of (1) was employed using $(NH_4)_2MoS_4$ (2.6 g, 10 mmol) and $(NH_4)_2WS_4$ (1.74 g, 5 mmol) instead of using the ratio of Mo:W as 1:2. The final product was isolated as dark green needle shaped crystal with 70 % yield based on the starting material. Anal: found. (calcd. (%)) for $C_6H_{10}O_{17}S_4Mo_2WCs_2$: C: 6.56 (6.40); H: 0.92 (0.89); S: 11.68 (11.38); W: 16.42 (16.37); Mo: 17.34(17.08).

RESULTS AND DISCUSSION

Synthesis and properties of molybdenum(IV)-Tungsten(IV) mixed-metal clusters with $\{MoW_2S_4\}^{4+}$ and $\{WMo_2S_4\}^{4+}$ cores: Two types of reaction for the synthesis of the mixed metal clusters have been tried during the study. Type-1, reduction of a mixture of Mo(VI) and W(V) or a mixture of W(VI) and Mo(V) by $NaBH_4$ in acidic medium, Type-2, the reduction of a mixture of Mo(VI) and W(V) or W(VI) and Mo(V) by Mo(III) or W(III) compounds. The method described in the experimental section employs a reaction of type 1 as described in **Scheme-I**.

The combination of the starting materials gave better yields of $[MoW_2S_4(H_2O)_9]^{4+}$ and $[WMo_2S_4(H_2O)_9]^{4+}$ than any other methods. The oxidation states of molybdenum and tungsten in the mixed-metal clusters are four. The mixed-metal clusters are stable towards air-oxidation and the absorption spectra of the clusters in 2 M HCl or in the freshly prepared 2 M *p*-toluene sulfonic acid solutions do not change for several days in the air. The use of $(NH_4)_2S_2O_8$ as an external oxidant for $(NH_4)_2WS_4$ or $(NH_4)_2MoS_4$ leads to the quantitative generation of mixed-metal trimeric core. It has been observed that a solution containing the mixture of $(NH_4)_2WS_4$ and $(NH_4)_2MoS_4$ the either in the stoichiometric ratio of 2:1, 1:2 or 1:1 without use of $(NH_4)_2S_2O_8$ when subject for reaction under the similar condition produced a gray-coloured solution which on chromatographic separation resulted a separation of two distinct bands, blue-violet band at top ($\lambda_{max} = 570$ nm) and green band at bottom ($\lambda_{max} = 609$ nm) responsible for $[W_3S_4(H_2O)_9]^{4+}$ and $[Mo_3S_4(H_2O)_9]^{4+}$, respectively. The suggests about the crucial role of pentavalent thio-metallate(V) in the generation of mixed-metal trimeric core.

The resultant oxalato complexes were analyzed to give S/W = 2.02 ± 0.05 and 3.98 ± 0.05 (four determination) for

$\{\text{MoW}_2\text{S}_4\}^{4+}$ and $\{\text{WMo}_2\text{S}_4\}^{4+}$ core, respectively. Sulphur was determined gravimetrically as BaSO_4 and tungsten by the thiocyanate photometric method). The change of the ion was estimated to be 4+ on the basis of its behaviour similar to that of the $\text{Mo}_3\text{S}_4^{4+}$ or $\text{W}_3\text{S}_4^{4+}$ aqua ion on the ion exchanger.

The proposed structure for both derivatives has schematically been represented in the following Fig. 1.

Fig. 1.

Electronic spectral study: Electronic spectra of $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and $[\text{WMo}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ along with their corresponding oxalato derivatives have been depicted in the Figs. 2 and 3, respectively in the UV-visible region.

Fig. 2. Electronic spectra of $[\text{Mo}_{3-n}\text{W}_n\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ clusters ($n = 1-2$) in 2 M HPTS: (—) $[\text{Mo}_2\text{WS}_4(\text{H}_2\text{O})_9]^{4+}$ and (---) $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$

The peak positions (nm) and ϵ -value are given in Table-1. A distinct splittings of absorption peaks in the visible regions are observed in the mixed metal aqua ions and the corresponding oxalato derivatives. The longer peak wave length positions for the four aqua ions and of the four oxalato-derivatives shift to the longer wavelength, respectively, when the tungsten atom is replaced by molybdenum.

It was demonstrated by the following two experiments that the clusters $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and $[\text{WMo}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ are

Fig. 3. Electronic spectra of $[\text{Mo}_{3-n}\text{W}_n\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3]^{2-}$ clusters ($n = 1-2$) in 0.2 M KCl: (—) $[\text{Mo}_2\text{WS}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3]^{2-}$ and (---) $[\text{MoW}_2\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3]^{2-}$

not a mixture of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$; as: (1) The mixture of $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ is clearly separated into two bands by Sephadex G-15 column chromatography, where as the $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ or $[\text{WMo}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ cluster is not separated by the method and (2) the electronic spectrum of the $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ cluster is clearly different from that of the mixture of 33 % $\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ and 66 % $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ clusters. The same is true for the $[\text{WMo}_2\text{S}_4(\text{H}_2\text{O})_9]^{4+}$ cluster. Elemental analyses for both complexes also support the formula as proposed.

Infrared spectra: Muller and co-workers have extensively studied on molybdenum-sulphur compounds and assigned the band at 460 cm^{-1} for $\nu\text{Mo}_3\text{-S}$ vibration in $(\text{NH}_4)_2[\text{Mo}_3(\mu_3\text{-S})(\mu_2\text{-S}_2)_3(\text{S}_2)_3]$. Infrared spectra of the four aqua clusters, $[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$, $[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$, $[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$ and $[\text{WMo}_2\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$ shows $\nu\text{M}_3\text{-S}$ vibration in the region of $550\text{-}400 \text{ cm}^{-1}$. Each spectrum was two large absorption bands in the region. The peak position shift to the lower wave number when the molybdenum atom is replaced by the tungsten atom. The bands are tentatively assigned to $\nu\text{M-OH}_2$ at the higher wave number and $\nu\text{M}_3\text{-S}$ at lower wave numbers.

Electrochemical study: Both the molybdenum-tungsten mixed-metal clusters $\text{Cs}_2[\text{WMo}_2\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$ and $\text{Cs}_2[\text{MoW}_2\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$ show the similar three reduction processes as do $\text{Cs}_2[\text{Mo}_3\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$ and $\text{Cs}_2[\text{W}_3\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$.

The cyclic-voltammogram of $\text{Cs}_2[\text{WMo}_2\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$ shows two electrochemically quasi-reversible and one chemically reversible redox couples appear at -0.6 and -1.2 V , respectively, although the third reduction peak disappears because of the very large catalytic H_2 -evolution. Sharp redox peak at -0.75 V due to the strong adsorption of

TABLE-1
ELECTRONIC SPECTRAL DATA FOR TRIMER WITH $\{\text{W}_n\text{Mo}_{3-n}\text{S}_4\}^{4+}$ CORES

No.	Compound	λ_{max} nm ($\text{M}^{-1}, \text{m}^{-1}$)	Ref.
1	$[\text{Mo}_3\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$	602(351), 367(5190)	Present work
2	$[\text{W}_3\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$	560(490), 430sh(330), 314(7040)	Present work
3	$[\text{MoW}_2\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$	572(380), 500(340) 400sh(1400), 340(6500)	Present work
4	$[\text{WMo}_2\text{S}_4(\text{H}_2\text{O})_9](\text{PTS})_4$	600(340), 500 Sh(300) 440sh(920), 350(4900)	Present work
5	$\text{Cs}_2[\text{Mo}_3\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$	622(252), 366(3623) 256(9210)	To be published
6	$\text{Cs}_2[\text{W}_3\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$	586(572), 361(1556) 305(8265), 253(12470)	To be published
7	$\text{Cs}_2[\text{MoW}_2\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$	590(480), 515(425) 400Sh (1630), 340(6000), 235sh(24000)	Present work
8	$\text{Cs}_2[\text{WMo}_2\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$	620(430), 500Sh(410) 425sh(1500), 360 (5200) 245sh(19200)	Present work

$[WMo_2S_4(C_2O_4)_3(H_2O)_3]^{2-}$ on the electrode overlap the peaks of the first reduction process.

Thus, the electrolysis yields the one-electron reduction product $[WMo_2S_4(C_2O_4)_3(H_2O)_3]^{3-}$ which is stable in the inert atmosphere only but is readily re-oxidized to $[WMo_2S_4(C_2O_4)_3(H_2O)_3]^{2-}$ when exposed to air. The reversible conversion of the spectra of the clusters was confirmed through several cycles of electrolytic reduction and air-oxidation, indicating the stability of cores. The reduction product which is a mixed-metal and mixed valance cluster, has absorption maxima at 780, 610 and 350 nm.

Another molybdenum-tungsten mixed-metal cluster complex $Cs_2[Mo_2W_2S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$ shows three consecutive one electron reduction processes in the cyclic voltammetric measurements. The first reduction process at the mercury electrode, is also overstepped by the absorption redox couple at $0 > V$. The electrolysis is coulometrically confirmed to be a one electron reduction process to give the mixed-metal and mixed-valance cluster $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{3-}$ that has absorption maxima at 720(sh), 580, 340 nm.

The cyclic voltammogram of $Cs_2[MoW_2S_4(C_2O_4)_3(H_2O)_3]$ and $Cs_2[WMo_2S_4(C_2O_4)_3(H_2O)_3]$ have been shown in Fig. 4a-b.

Table-2 summarizes the half-wave potentials and the wave slopes of the three consecutive one electron reduction precesses of the molybdenum-tungsten mixed-metal $Cs_2[MoW_2S_4(C_2O_4)_3(H_2O)_3]$ and $Cs_2[WMo_2S_4(C_2O_4)_3(H_2O)_3]$. The potential of the absorption redox couple of the clusters on the mercury electrode is almost independent of the metals in the cluster, indicating that the specific treaction with mercury is through the sulphur atom is the clusters with the same order. The catalytic hydrogen evolution wave accompanying the third reduction process depands markedly on the cluster, the catalytic reduction rate is in the order of $[W_3S_4(C_2O_4)_3(H_2O)_3]^{2-} > [MoW_2S_4(C_2O_4)_3(H_2O)_3]^{2-} > [WMo_2S_4(C_2O_4)_3(H_2O)_3]^{2-} > [Mo_3S_4(C_2O_4)_3(H_2O)_3]^{2-}$ while $E_{1/2}$ potentials for IV/III/III $\rightarrow$ III/III/III become less negative in the reverse order.

The reduction potentials of the mixed-metal clusters lie between those of $[Mo_3S_4(C_2O_4)_3(H_2O)_3(C_2O_4)_3]^{2-}$ and $[W_3S_4(C_2O_4)_3(H_2O)_3]^{2-}$ but are not proportionally distributed for the first reduction process, the potential difference between $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{2-/3-}$ and $[Mo_2W_2S_4(C_2O_4)_3(H_2O)_3]^{2-/3-}$ (90 mV) and that between $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{3-/4-}$ and $[WMo_2S_4(C_2O_4)_3(H_2O)_3]^{3-/4-}$ (110 mV). The largest potential difference for the second and third processes lies between $[Mo_2WS_4(C_2O_4)_3(H_2O)_3]^{3-/4-}$ and $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{3-/4-}$ and $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{4-/5-}$ and $[W_3S_4(C_2O_4)_3(H_2O)_3]^{4-/5-}$, respectively.

These result suggest that the proceeding reduction centre of the molybdenum-tungsten mixed-metal clusters is mainly on the Mo atoms rather than the Watoms. Thus the electrode reactions are formally described as

Fig. 4. (a-b) Cyclic voltammogram of $[MoW_2S_4(C_2O_4)_3(H_2O)_3]^{2-}$ and $[Mo_2WS_4(C_2O_4)_3(H_2O)_3]^{2-}$

The spectroscopic measurements involving ESR spectroscopy may help to elucidate in more detail about the electronic states of the reduced clusters especially the mixed-metal mixed-valence cluster, but due to the instability of reduced species, the corresponding clusters were not isolated successfully for their further spectroscopic study. However, the experimental work on the synthetic procedure is being under progress for further study. In these oxalato complexes, the molybdenum system, $Cs_2[Mo_3S_4(C_2O_4)_3(H_2O)_3]$, shows as reversible one electrode reduction with $E_{1/2} = -0.689$ V vs. Ag-AgCl with the appearance of cathodic peak potential at -0.733 V along with another ill-defined reduction centered at -1.034 V. The corres-

TABLE-2
COMPARISON OF $E_{1/2}$ FOR THE REDUCTION OF CLUSTERS WITH $\{W_nMo_{3-n}S_4\}^{4+}$ CORES

Compounds containing homo/hetero core	$E_{1/2}$		
	IV, IV, IV/IV, IV, III (V)	IV, IV, III/IV, III, III (V)	IV, III, III/III, III, III (V)
$Cs_2[Mo_2S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$	-0.69	-1.034	-1.39
$Cs_2[WMo_2S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$	-0.73	1.08	-1.39
$Cs_2[MoW_2S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$	-0.84	-1.40	-1.78
$Cs_2[W_3S_4(C_2O_4)_3(H_2O)_3] \cdot 2H_2O$	-1.23	-1.41	-1.88

ponding tungsten complex shows an irreversible reduction at -1.23 V. In the oxalato complex, We could not locate any anodic peak for oxidation up to +0.6 V because of the limitation of potential window of solvent water.

ACKNOWLEDGEMENTS

The authors expressed their sincere gratitude to Prof. S. Sarkar, Head, Chemistry Department, Indian Institute of Technology, Kanpur for his valuable suggestions and guidance.

REFERENCES

1. T. Shibahara. *Coord. Chem. Rev.*, **123**, 73 (1993); T. Shibahara, *Adv. Inorg. Chem.*, **37**, 143 (1991).
2. T. Shibahara, M. Yamasaki, G. Sakane, K. Minami, T. Yabuki and A. Rhimura, *Inorg. Chem.*, **31**, 640 (1992); Q.-T. Liu, J.-X. Lu and A.G. Sykes, *Inorg. Chim. Acta*, **200**, 623 (1992).
3. F.A. Cotton and S.K. Mandal, *Inorg. Chim. Acta*, **192**, 71 (1992).
4. F.A. Cotton and W. Schwotzer, *Angew. Chem.*, **94**, 652 (1982).
5. M.H. Chisholer, K. Folting, J.C. Huffman and E.M. Kober, *Inorg. Chem.*, **24**, 241 (1985).
6. B. Wang, Y. Sasaki, A. Nagasawa and T. Ito, *J. Am. Chem. Soc.*, **108**, 6059 (1986); B. Wang, Y. Sasaki, S. Ikari, K. Kimura and T. Ito, *Chem. Lett.*, 1955 (1987).
7. A. Patel and D.T. Richers, *J. Chem. Soc. Chem. Commun.*, 274 (1990).
8. H. Brunner, H. Kanermann and J. Wachter, *J. Organomet. Chem.*, **265**, 189 (1984).
9. N.-Y. Zhu, Y.-F. Zhing and X.-T. Wu, *Inorg. Chem.*, **29**, 2705 (1990); N.-Y. Zhu, Y.-F. Zhing and X.-T. Wu, *Polyhedron*, **10**, 2743 (1991).
10. N.-Y. Zhu, X.-T. Wu and J.-X. Lu, *J. Chem. Soc. Chem. Commun.*, 235 (1991).
11. T. Shibahara, M. Yamasaki, T. Watase and A. Ichimura, *Inorg. Chem.*, **33**, 292 (1994); A. Mullier, V.P. Feddin, E. Dienmann, H. Bogge, E. Krickemeyer, D. Solter, A.M. Guuliani, R. Barbieri and P. Adler, *Inorg. Chem.*, **33**, 2243 (1994).
12. Y. Ide and T. Shibahara, *Inorg. Chem.*, **46**, 357 (2007); T. Yamauchi, H. Takagi, Shibahara and H. Akashi, *Inorg. Chem.*, **45**, 5429 (2006).
13. H. Akashi, K. Isobe and T. Shibahara, *Inorg. Chem.*, **44**, 3494 (2005); Y. Ide, M. Sasaki, M. Maeyama and T. Shibahara, *Inorg. Chem.*, **43**, 602 (2004).
14. T. Shibahara, M. Tsuboi, S. Nakaska and Y. Ide, *Inorg. Chem.*, **42**, 935 (2003); S.-B. Yu, M. Droege, B. Segal, S.-H. Kim, T. Sanderson and A.D. Watson, *Inorg. Chem.*, **39**, 1325 (2000).
15. S. Sarkar, R. Sah, P.K. Chaudhary, R. Maiti and S.K. Das, *Proc. Indian Acad. Sci. (Chem. Sci.)*, **105**, 355 (1995).