Inhibitory Activity of the Extract from *Broussonetia papyrifera* Fruits to Cellular Lipid Peroxidation *in vitro*

YANG LI¹, XIAOYA SHANG¹, WEINING NIU¹, CHUNLAN XU¹ and CHUANGUANG QIN^{1,2,*}

¹School of Life Sciences, Northwestern Polytechnical University, Xi'an 710072, P.R. China

²Key Laboratory on Space Applied Physics and Chemistry of Ministry of Education; Key Laboratory on Macromolecular Science & Technology of Shaanxi Province, Department of Applied Chemistry, School of Natural and Applied Sciences, Northwestern Polytechnical University, Xi'an, 710129 P.R. China

*Corresponding author: Fax: +86 29 88460332; Tel: +86 13991392191; E-mail: qinchg@nwpu.edu.cn

Received: 21 March 2013;

Accepted: 11 June 2013;

Published online: 26 December 2013;

AJC-14504

On the basis of our previous work on the antioxidant and anticancer activities of the extract of the *Broussonetia papyrifera* fruits, *in vitro* inhibitory activity of the extract to cellular lipid peroxidation was under study in present work. Inhibitory activity of the extract was evaluated by determining the ability to inhibit H₂O₂-induced erythrocyte hemolysis and lipid peroxidation of liver homogenate, to inhibit mitochondrial swelling and to scavenge hydroxyl radical. The total flavonoid content was found to be 75.46 ± 1.91 mg rutin equivalents g⁻¹ lyophilized extract. The extract was a potent inhibitor of erythrocyte hemolysis, lipid peroxidation and mitochondrial swelling and was a scavenger of hydroxyl radical. The inhibitory activity of the extract showed a dose-dependent increase in the assays. This study suggested the extract was a potent antioxidant and might be developed as a potential botanical medicine or functional food.

Keywords: *Broussonetia papyrifera* fruit, Ethanolic extract, Lipid peroxidation, Antioxidant activity, Flavonoid compound.

INTRODUCTION

Reactive oxygen species (ROS) are generated during normal metabolism in the human body, which could attack various biomolecules including proteins, lipoproteins, lipids and DNA readily to induce oxidative damage. It is believed that several chronic human diseases, including cardiovascular diseases, rheumatism, diabetes mellitus and cancer, have a very close relationship with oxidative damage¹. Therefore, the usefulness of antioxidants in protection against these diseases captures the attention.

Numerous natural antioxidants have already been isolated from plants such as fruits, vegetables, seeds and roots. The phenolic compounds, comprising at least 8000 different known structures including simple phenols, phenolic acids, flavonoids, are the important ones². It has been stated that phenolic compounds are multifunctional antioxidants acting as scavenging free radicals³, reducing agents⁴, hydrogen-donating antioxidants⁵ and improving the endogenous antioxidant system⁶. The phenolic compounds could also influence cellular oxidation and reduction environment to regulate proliferation and/or be a novel redox drug for cancer^{7,8}. Therefore, as development activity increases in this area, it is not surprising that phenolic compounds have gained tremendous interest.

Broussonetia papyrifera (Moraceae) grows naturally in Pacific countries such as China, Burma, Polynesia and USA⁹. The fruits, barks and roots are all used in traditional Chinese medicines. For example, the fruits have been used for treatment of ophthalmic and Alzheimer's disease in Chinese folk^{10,11}. The compounds extracted from different parts of *Broussonetia papyrifera* have been extensively studied and shown to inhibit PTP1B enzyme¹² and lipid peroxidation¹³, antihyperglycemic¹⁴ and antihepatotoxic¹⁵. However, study of ethanolic extract from the fruits has been limited.

According to our previous investigation, the extract had antioxidant activity and could inhibit cancer cells proliferation¹⁶. Therefore, the objective of this study was to investigate inhibitory activity of the extract to cellular lipid peroxidation *in vitro*. This work has provided some valuable data for future study of the extract.

EXPERIMENTAL

Fresh *Broussonetia papyrifera* fruits were collected from Beilin district of Xi'an, China during July 2011 and identified at the School of Life Sciences, Northwestern Polytechnical University. A voucher specimen is preserved in the same place. The fruits were placed in polyethylene bags and stored at -20 °C until use.

Preparation of the extract: *Broussonetia papyrifera* fruits were extracted at 40 °C in 80 % ethanol for 2 h. The extract was filtered through filter paper and evaporated at 40 °C in rotary evaporator (EYELA, N1001, Japan). Subsequently, the concentrated solution was passed through a 2 g sorbent weight C₁₈ SPE cartridge (AGELA, China), previously activated with methanol followed by deionized water. 10 mL of deionized water and ethyl acetate were used to wash the SPE cartridge and then the extract was eluted with methanol. The extract was dried by rotary evaporator at 40 °C, redissolved in deionized water and then lyophilized in a lyophilizer (CHRIST, Alpha 1-2, Germany). The extract was placed in a plastic bottle and stored at room temperature.

Determination of total flavonoid content: Total flavonoid content was determined according to the aluminum chloride colorimetric method¹⁷. Briefly, 1 mL of 1.6 mg mL⁻¹ the extract was placed in a 10 mL volumetric flask and 0.4 mL of 5 % sodium nitrite was added. After 6 min, 0.4 mL of 10 % aluminum chloride was added. 4 mL of 4 % sodium hydroxide and 4.2 mL of 60 % ethanol were added after 6 min. The solution was mixed well and incubated for 15 min at room temperature and then the absorbance was measured at 510 nm by a UV-visible spectrophotometer (HITACHI, F2300, Japan). The measurement was compared to a standard curve of rutin and expressed as milligram rutin equivalents (RE) g⁻¹ lyophilized extract.

Preparation of erythrocytes and liver homogenate: Kunming mice, body weight of 18-22 g, were purchased from the Experimental Animal Center of College of Medicine, Xi'an Jiaotong University. The mice were housed in polypropylene cages and given standard food and water *ad libitum*. After the blood was collected by enucleating eyes in heparinized tubes, the mice were sacrificed by cervical dislocation and the liver was excised quickly. Erythrocytes were isolated and stored according to the method of Yuan *et al.*¹⁸ and Yang *et al.*¹⁹ with some modifications. Blood samples were washed three times by centrifugation (2000 rpm, 5 min) in 10 volumes of phosphate buffer saline (PBS: 150 mmol L⁻¹ NaCl, 8.1 mmol L⁻¹ Na₂HPO₄, 1.9 mmol L⁻¹ NaH₂PO₄, 10 mmol L⁻¹ EDTA, pH 7.4). The supernatant and buffy coats were carefully removed after each centrifugation and then the compacted erythrocytes were obtained. The erythrocytes were stored at 4 °C and used within 6 h.

On the other hand, the liver was washed, weighed and then homogenized with ice-cold PBS in an ice bath to yield a 10 % (w/v) homogenate by means of a Potter-Elvehjem homogenizer. The liver homogenate was stored at 4 °C.

Inhibition of erythrocyte hemolysis: The inhibition of erythrocyte hemolysis was determined according to the method of Tedesco *et al.*²⁰ with some modifications. To 1 mL of 0.5 % (v/v) suspension of erythrocytes in PBS, 0.1 mL of the extract at different concentrations and 0.1 mL of 100 mmol L⁻¹ H₂O₂ were added. The mixture was incubated in water bath at 37 °C for 1 h. After that, 4 mL PBS was added to the mixture and centrifuged at 2000 rpm for 5 min. The absorbance of the supernatant was measured at 415 nm to determine the hemolysis. The inhibitory effect of the extract was compared with positive control of ascorbic acid. The inhibitive rate of hemolysis was calculated as follows:

$$\text{Inhibition (\%)} = \frac{[A_{415(\text{Control})} - A_{415(\text{Sample})}]}{A_{415(\text{Control})}} \times 100 \%$$

The hemolysis induced by the extract was also evaluated. As described above, erythrocytes and 10 µg of the extract were incubated at 37 °C for 1 h and the absorbance was measured.

Lipid peroxidation inhibitory activity: Lipid peroxidation was determined by the measurement of malondialdehyde (MDA) according to the method of Ohkawa *et al.*²¹ with slight modifications. 0.2 mL of 10 % liver homogenate and 0.1 mL of the extract at different concentrations were mixed and then treated with 0.1 mL of 100 mmol L⁻¹ H₂O₂ at 37 °C for 1 h. Subsequently, the mixture was combined with 0.2 mL of 8.1 % SDS, 1.5 mL of 15 % trichloroacetic acid and 1.5 mL of 0.8 % thiobarbituric acid. The mixture was heated to 95 °C for 1 h. After cooling in an ice bath, the mixture was centrifuged at 3000 rpm for 10 min. The absorbance of the supernatant fraction was measured at 532 nm on a spectrophotometer. Rutin was used as positive control. The inhibition of lipid peroxidation was calculated using the following formula:

$$\text{Inhibition (\%)} = \frac{[A_{532(\text{Control})} - A_{532(\text{Sample})}]}{A_{532(\text{Control})}} \times 100 \%$$

Mitochondrial swelling assay: Mitochondrial swelling assay was performed according to the procedures described by Song *et al.*²². The liver homogenate was centrifuged at 1000 rpm at 4 °C for 20 min and then the mitochondria were isolated by centrifugation of the supernatant at 10000 rpm at 4 °C for 20 min. The pellet was washed with ice-cold PBS by centrifuged at 10000 rpm at 4 °C for 20 min. The mitochondria were resuspended in ice-cold PBS to a concentration of 0.5 mg mL⁻¹ of protein by following the Bradford method, using BSA as standard and stored at 4 °C. To a mixture of 0.2 mL of the extract at different concentrations and 2 mL of mitochondrial suspension, 0.2 mL of 0.5 mmol L⁻¹ ferrous sulfate and 0.2 mL of 0.5 mmol L⁻¹ ascorbic acid were added. The mixture was then incubated at 37 °C and the absorbance of the mixture was measured at 520 nm at intervals. A decrease of the absorbance indicates an increase of mitochondrial swelling. PBS was used instead of the extract, ferrous sulfate and ascorbic acid in the control group.

Hydroxyl radical scavenging assay: The assay was determined using the methods of Wang *et al.*²³ with some modifications. The OH[•] in aqueous media is generated through the Fenton reaction. To a 1.5 mL tube, 67 µL of 2.2 mmol L⁻¹ H₂O₂, 67 µL of 2.25 mmol L⁻¹ salicylic acid, 67 µL of 2.25 mmol L⁻¹ FeSO₄ and 0.1 mL of the extract at different concentrations were added, mixed and stood for 5 min in the dark at room temperature. The absorbance was measured at 510 nm. Ascorbic acid was used as positive control. The free radical scavenging activity was calculated as follows:

$$\text{Inhibition (\%)} = \frac{[A_{510(\text{Control})} - A_{510(\text{Sample})}]}{A_{510(\text{Control})}} \times 100 \%$$

Statistical analysis: Results were expressed as the mean ± SD of three independent experiments. Differences between sample group and control group were analyzed with One-way ANOVA procedures (Prism 5, GraphPad Software, USA). A value of *p* < 0.05 was regarded as being significant. IC₅₀ values were calculated by regression analysis.

RESULTS AND DISCUSSION

Ethanol extract of *Broussonetia papyrifera* fruits: As described above, the extract was purified by washing the C₁₈ cartridge and the sugars, acids and other water soluble compounds were removed. The extract was the pigment of fresh *Broussonetia papyrifera* fruit and showed orange colour. By phytochemical screening, it was found that the extract contained flavonoids. This may be the reason for its antioxidant activity. The total flavonoid content was 75.46 ± 1.91 mg rutin equivalent g⁻¹ lyophilized extract, which was higher than that of mulberry leaves of 19 varieties of species¹⁷.

Inhibition of erythrocyte hemolysis: Because of the high concentration of polyunsaturated fatty acids in the membrane and the O₂ transport associated with redox active hemoglobin molecules, which are potent promoters of ROS, erythrocytes are considered as prime targets for free radical attack²⁴. Therefore, erythrocytes have been used as a cellular model to investigate oxidative damage in biomembranes.

Hemolysis of erythrocytes induced by the extract was determined. The percentage of hemolysis was found at 4.11 ± 0.35 %, which was not significantly different from control of 3.91 ± 0.53 %. This result showed the extract had no harmful effect on erythrocytes.

The inhibitory effect of the extract on erythrocyte hemolysis was evaluated by oxidative stress induced by H₂O₂. Fig. 1 showed that lysis of erythrocytes caused by H₂O₂ was significantly inhibited by the extract. The inhibitory activity to erythrocyte hemolysis was dose-dependent over the range of tested concentrations. The IC₅₀ value of the extract was $12.84 \mu\text{g mL}^{-1}$, which was lower than that of positive control ($31.77 \mu\text{g mL}^{-1}$). The results indicated the extract could provide stronger protection for erythrocytes against damage caused by H₂O₂ than ascorbic acid.

Fig. 1. Inhibition of *Broussonetia papyrifera* fruit extract against H₂O₂ induced erythrocyte hemolysis. Data is expressed as mean $\pm$ SD (n = 3). Values in different groups are significantly different from control, $p < 0.05$. (Δ) *Broussonetia papyrifera* fruit extract; (□) ascorbic acid

Inhibitory activity to lipid peroxidation: Lipid peroxidation is an oxidative alteration of polyunsaturated fatty acids,

which is a common process in the biological systems. Malondialdehyde one of the major products of lipid peroxidation, has been extensively used as an index of lipid peroxidation and as a marker of oxidative stress²⁵. The measure of malondialdehyde formation has been widely used for determination of anti-lipid peroxidation activity of natural extracts.

Compared to the control, addition of H₂O₂ to the liver homogenate for 1 h significantly increased malondialdehyde formation. However, the extract ($12.5\text{--}400 \mu\text{g mL}^{-1}$) added to the reaction system significantly reduced malondialdehyde formation, indicating significant anti-lipid peroxidation activity. Fig. 2 showed that inhibition of lipid peroxidation increased with increasing concentrations of the extract. The IC₅₀ value of the extract for lipid peroxidation inhibition on liver homogenate was $155.7 \mu\text{g mL}^{-1}$, which seemed to have roughly the same inhibitory effect as rutin ($147 \mu\text{g mL}^{-1}$).

Fig. 2. Inhibitory activity of *Broussonetia papyrifera* fruit extract to lipid peroxidation. Data is expressed as mean $\pm$ SD (n = 3). Values of samples at $12.5\text{--}400 \mu\text{g mL}^{-1}$ in different groups are significantly different from control, $p < 0.05$. (Δ) *Broussonetia papyrifera* fruit extract; (□) rutin

Inhibitory effect on mitochondrial swelling: Reaction system of ferrous sulfate and ascorbic acid produced H₂O₂ to initiate lipid peroxidation and destroy the structure of mitochondrial membrane, which was the reason for mitochondrial swelling²⁶. The results were shown in Fig. 3. A decrease of absorbance at 520 nm represented the swelling of mitochondrial increased and the integrity of mitochondrial was damaged. Compared to the control group, absorbance of damage groups (with ferrous sulfate and ascorbic acid) decreased considerably with time. However, the swelling of mitochondrial decreased present in the extract. This result indicated the extract inhibited mitochondrial swelling and the inhibitory activity to mitochondrial swelling increased as the concentration increased. Rutin showed stronger inhibitory effect than that of the extract at the same concentration of $200 \mu\text{g mL}^{-1}$.

Hydroxyl radical scavenging activity: Hydroxyl radical, known to be the most biologically active free radical, is a major ROS in living organisms and is being continuously formed in a process of reduction of oxygen to water². It can attack and

Fig. 3. Inhibitory effect of *Broussonetia papyrifera* fruit extract on mitochondrial swelling. Data is expressed as mean $\pm$ SD ($n = 3$). Values in different groups are significantly different from damage group, $p < 0.05$. (□) control group; (◇) damage group; (▽) 50 $\mu\text{g mL}^{-1}$ of the extract; (△) 200 $\mu\text{g mL}^{-1}$ of the extract; (○) 800 $\mu\text{g mL}^{-1}$ of the extract; (◁) 200 $\mu\text{g mL}^{-1}$ of rutin

damage virtually any molecule including macromolecules such as DNA, proteins, membrane lipids and carbohydrates²⁷. According to the results obtained above, the extract could inhibit H_2O_2 -induced cellular lipid peroxidation. Thus, we conjectured the extract had the ability of scavenging hydroxyl radical. In this assay, the extract was evaluated for this ability. As shown in Fig. 4, a concentration dependent inhibition was observed. At final concentration of 750 $\mu\text{g mL}^{-1}$, the scavenging ability of the extract was $86.23 \pm 0.02\%$. The extract exhibited low IC_{50} value of 725.1 $\mu\text{g mL}^{-1}$ as compared to ascorbic acid at 1640.9 $\mu\text{g mL}^{-1}$. The scavenging activity of the extract compared with the positive control suggested that the extract was a greater potent scavenger of hydroxyl radical than ascorbic acid.

Fig. 4. Hydroxyl radical scavenging activity of *Broussonetia papyrifera* fruit extract. Data is expressed as mean $\pm$ SD ($n = 3$). Values in different groups are significantly different from control, $p < 0.05$. (△) *Broussonetia papyrifera* fruit extract; (□) ascorbic acid

Conclusion

This study demonstrated that the extract from *Broussonetia papyrifera* fruits had inhibitory activity to cellular lipid peroxidation. As an antioxidant, the isolation and characterization of the active compounds and the determination of biological activity *in vivo* must be processed continuously. In addition to being a potential botanical medicine, the extract can also be developed to functional food in the future due to its properties of beautiful colour, edible and antioxidant activity.

ACKNOWLEDGEMENTS

This work was financially supported by the National Natural Science Foundation of China (31001012, 31101304, 20802057, 20672086) and the Natural Science Basic Research Plan in Shaanxi Province of China (2012JZ2002).

REFERENCES

1. K. Pong, *Expert Opin. Biol. Ther.*, **3**, 127 (2003).
2. B. Lipinski, *Oxid. Med. Cell Longev.*, **2011**, 809696 (2011).
3. J. Bhagat, R. Lobo, V. Parmar and M. Ballal, *Chin. J. Nat. Med.*, **9**, 456 (2011).
4. S. Arabshahi-Delouee and A. Urooj, *Food Chem.*, **102**, 1233 (2007).
5. C.A. Rice-Evans, N.J. Miller and G. Paganga, *Free Radic. Biol. Med.*, **20**, 933 (1996).
6. H.F. Al-Azzawie and M.S. Alhamdani, *Life Sci.*, **78**, 1371 (2006).
7. E.H. Sarsour, M.G. Kumar, L. Chaudhuri, A.L. Kalen and P.C. Goswami, *Antioxid. Redox Signal.*, **11**, 2985 (2009).
8. G.T. Wondrak, *Antioxid. Redox Signal.*, **11**, 3013 (2009).
9. D. Seelenfreund, R. Pina, K.Y. Ho, S. Lobos, X. Moncada and A. Seelenfreund, *New Zeal. J. Bot.*, **49**, 413 (2011).
10. H. Matsuda, H. Cai, M. Kubo, H. Tosa and M. Iinuma, *Biol. Pharm. Bull.*, **18**, 463 (1995).
11. R.Q. Mei, Y.H. Wang, G.H. Du, G.M. Liu, L. Zhang and Y.X. Cheng, *J. Nat. Prod.*, **72**, 621 (2009).
12. R.M. Chen, L.H. Hu, T.Y. An, J. Li and Q. Shen, *Bioorg. Med. Chem. Lett.*, **12**, 3387 (2002).
13. H.H. Ko, S.M. Yu, F.N. Ko, C.M. Teng and C.N. Lin, *J. Nat. Prod.*, **60**, 1008 (1997).
14. J.Y. Cha, Y.T. Kim, H.S. Kim and Y.S. Cho, *J. Med. Food*, **11**, 499 (2008).
15. S.Y. Kim, J.H. Kim, S.K. Kim, M.J. Oh and M.Y. Jung, *J. Am. Oil Chem. Soc.*, **71**, 633 (1994).
16. Y. Li, H.L. Li, Y. Zhang, L. Li and C.G. Qin, *Asian J. Chem.*, **25**, 5453 (2013).
17. J. Zhishen, T. Mengcheng and W. Jianming, *Food Chem.*, **64**, 555 (1999).
18. X.P. Yuan, J. Wang, H.Y. Yao and F. Chen, *LWT-Food Sci. Technol.*, **38**, 877 (2005).
19. H.L. Yang, S.C. Chen, N.W. Chang, J.M. Chang, M.L. Lee, P.C. Tsai, H.H. Fu, W.W. Kao, H.C. Chiang, H.H. Wang and Y.C. Hseu, *Food Chem. Toxicol.*, **44**, 1513 (2006).
20. I. Tedesco, M. Russo, P. Russo, G. Iacomino, G.L. Russo, A. Carraturo, C. Faruolo, L. Moio and R. Palumbo, *J. Nutr. Biochem.*, **11**, 114 (2000).
21. H. Ohkawa, N. Ohishi and K. Yagi, *Anal. Biochem.*, **95**, 351 (1979).
22. Y.Y.N. Song, J. Hui, W. Kou, R. Xin, F. Jia, N. Wang, F.Q. Hu, H.L. Zhang and H.S. Liu, *Curr. Microbiol.*, **57**, 454 (2008).
23. S.Y. Wang and H.J. Jiao, *J. Agric. Food Chem.*, **48**, 5677 (2000).
24. S.M. Sadrzadeh, E. Graf, S.S. Panter, P.E. Hallaway and J.W. Eaton, *J. Biol. Chem.*, **259**, 14354 (1984).
25. D.R. Janero, *Free Radic. Biol. Med.*, **9**, 515 (1990).
26. Z.B. Chen, J. Lv, F. Chen and L. Lin, *J. Mol. Catal. B: Enzym.*, **55**, 99 (2008).
27. B. Halliwell, *Adv. Pharmacol.*, **38**, 3 (1997).