A New Cytotoxic Xanthone from *Hypericum chinense*

WENXIU XU¹, YANQING YE², HAIYIN YANG², XUEMEI GAO², QIUFEN HU² and YADONG GUO^{1,*}

¹School of Pharmaceutical Science & Yunnan Key Laboratory of Pharmacology for Natural Products, Kunming Medical University, Kunming 650500, P.R. China

²Key Laboratory of Chemistry in Ethnic Medicinal Resources, State Ethnic Affairs Commission & Ministry of Education, Yunnan University of Nationalities, Kunming 650031, P.R. China

*Corresponding author: E-mail: yadong_guo@yahoo.com.cn

Received: 14 September 2013;

Accepted: 19 December 2013;

Published online: 16 September 2014;

AJC-15919

A new xanthone, 1,8-dihydroxy-4-(2-hydroxyethyl)-3-methoxy-9*H*-xanthen-9-one (**1**) was isolated from the leaves and stems of *Hypericum chinense*. Its structure was elucidated by spectroscopic methods, including extensive 1D- and 2D NMR techniques. Compound **1** was tested for its cytotoxicity against five human tumor cell lines (NB4, A549, SHSY5Y, PC3 and MCF7) and it showed high cytotoxicity against PC3 and SHSY5Y cell with IC₅₀ values of 3.5 and 2.4 μM, respectively.

Keywords: Cytotoxic, Xanthone, *Hypericum chinense*.

INTRODUCTION

The family Clusiaceae is a rich source of xanthones, which show various bioactivities^{1,2}. These xanthones show various bioactivities, including, anti-hepatitis B virus³, anti-tobacco mosaic virus⁴, antibacterial^{5,6}, antioxidant^{7,8}, antiinflammatory⁹, tumor-promoting inhibition¹⁰ and cytotoxicity^{11,12}. The genus *hypericum* belonging to clusiaceae is distributed widely in temperate regions and has been used for traditional medicines in various parts of the world. In China, *Hypericum chinense* is used as a folk medicine for treatment of female disorders¹³. Previous phytochemical investigations on *H. chinense* resulted in the isolation of xanthones¹², acylphloroglucinols¹⁴, lactones¹⁵ and norlignans¹⁶ from this species.

With the aim of multipurpose utilization of herb plants and identify bioactive natural products from this genus, the phytochemical investigation on *H. chinense* was carried out. As a result, a new xanthone (**1**) was isolated from this plant. Its structure was elucidated on the basis of spectroscopic methods, including extensive 1D- and 2D NMR techniques. In addition, the cytotoxicity of compound **1** was evaluated. The details of the isolation, structure elucidation and cytotoxicity of this new compound are reported in this article.

EXPERIMENTAL

UV spectra were obtained using a Shimadzu UV-2401A spectrophotometer. IR spectra were obtained in KBr disc on a Bio-Rad Wininfrad spectrophotometer. ESI-MS were measured

on a VG Auto Spec-3000 MS spectrometer. ¹H, ¹³C and 2D NMR spectra were recorded on Bruker DRX-500 instrument with TMS as internal standard. Column chromatography was performed on silica gel (200-300 mesh), or on silica gel H (10-40 mm, Qingdao Marine Chemical Inc., China). Second separate was used an Agilent 1100 HPLC equipped with ZORBAX-C18 (21.2 mm × 250 mm, 7.0 mm) column and DAD detector.

The leaves and stems of *Hypericum chinense* L. were collected in Honghe Prefecture, Yunnan Province, People's Republic of China, in September 2010. The identification of the plant material was verified by Prof. Ren P.Y. (Xishuangbanna Botanical Garden). A voucher specimen (YNNI-2010-9-23) has been deposited in our laboratory.

Extraction and isolation: The air-dried and powdered leaves and stems of *H. chinense* (2 kg) were extracted 4 times with 70 % acetone (4 × 3 L) at room temperature and filtered. The crude extract (136 g) was applied to silica gel (200-300 mesh) column chromatography, eluting with a CHCl₃-acetone gradient system (9:1, 8:2, 7:3, 6:4, 5:5), to give five fractions A-E. The further separation of fraction A (9:1, 16.8 g) by silica gel column chromatography, eluted with petroleum ether-EtOAc (9:1, 8:2, 7:3, 6:4, 1:1), yielded mixtures A1-A5. The subfraction A3 (7:3, 4.21 g) was subjected to preparative HPLC (55 % MeOH, flow rate 12 mL/min) to give compound **1** (11.4 mg).

1,8-Dihydroxy-4-(2-hydroxyethyl)-3-methoxy-9*H*-xanthen-9-one (1**):** Obtained as a yellow gum; UV (MeOH)

$\lambda_{\max}$ (log ϵ) 210 (4.13), 242 (3.42), 305 (3.87) nm; IR (KBr, $\nu_{\max}$, cm^{-1}) 3438, 3075, 2942, 2880, 1652, 1600, 1542, 1465, 1370, 1183, 1062, 882, 768; ESIMS m/z (positive ion mode) 325 $[\text{M} + \text{Na}]^+$; HRESIMS (positive ion mode) m/z 325.0682 $[\text{M} + \text{Na}]^+$ (calcd. $\text{C}_{16}\text{H}_{14}\text{NaO}_6$ for 325.0688).

RESULTS AND DISCUSSION

A 70 % aq. acetone extract prepared from the leaves and stems of *H. chinense* was subjected repeatedly to column chromatography on Silic gel, Sephadex LH-20, RP-18 and Preparative HPLC to afford compound **1**. The structure of **1** was shown in Fig. 1 and its ^1H and ^{13}C NMR data were listed in Table-1.

Fig. 1. Structure of compound **1**

Compound **1** was isolated as a yellow gum. The HRESIMS of **1** gave the pseudomolecular $[\text{M} + \text{Na}]^+$ ion at m/z 325.0682, corresponding to a molecular formula of $\text{C}_{16}\text{H}_{14}\text{O}_6$. Its UV spectrum showed the maximum absorption at 305, 242 and 210 nm. Strong absorption bands accounting for $\nu(\text{OH})$ (3438 cm^{-1}), $\nu(\text{C}=\text{O})$ (1652 cm^{-1}) and aromatic groups (1600, 1542, 1465 cm^{-1}) could also be observed in its IR spectrum. The ^1H - and ^{13}C NMR spectrum (Table-1) displayed signals for all 16 carbons and 14 protons, including a xanthenes skeleton¹⁷ (C-1 - C-9, C-4a, C-8a - C-10a; H-2, H-5 - H-7), one methoxy group (δ_{C} 55.9 q, δ_{H} 3.80 s), a hydroxyethyl unit¹¹ [δ_{C} 34.3 t, 63.2 t; δ_{H} 2.50 t (7.2), 3.62 t (7.2)] and two phenolic hydroxy groups (δ_{H} 13.46 s and 13.14 s). The HMBC correlation (Fig. 2) of the methoxy proton signal (δ_{H} 3.80) with C-3 (δ_{C} 161.4) showed that the methoxy group was located at C-3. The long-range correlations of H_2 -1' (δ_{H} 2.50) to C-3 (δ_{C} 161.4), C-4 (δ_{C} 108.3) and C-4a (δ_{C} 155) were observed in **1**. This led us to conclude that the hydroxyethyl unit was located at C-4. Finally, HMBC correlations between the hydroxy proton (δ_{H} 13.46) and C-1 (δ_{C} 161.5), C-2 (δ_{C} 98.6) and C-9a (δ_{C} 104.6), as well as those between the other hydroxy proton (δ_{H} 13.14) and C-7

(δ_{C} 108.6), C-8 (δ_{C} 162) and C-8a (δ_{C} 109.2), led to the assignment of the phenolic groups at C-1 and C-8. The typical proton signals of ring A [δ_{H} 6.82 d (8.3), 7.49 t (8.3), 6.68 d (8.3)] and ring B (δ_{H} 6.62 s) also supported that **1** should be a 1,3,4,8-tetrasubstituted xanthone¹⁷. Thus, compound **1** was assigned as 1,8-dihydroxy-4-(2-hydroxyethyl)-3-methoxy-9H-xanthen-9-one.

Fig. 2. Key HMBC (→) correlations of **1**

Since xanthenes are known to exhibit potential cytotoxicity^{2,11,12}. Compound **1** was tested for its cytotoxicity against five human tumor cell lines (NB4, A549, SHSY5Y, PC3 and MCF7) using the MTT method as reported previously¹⁸. Taxol was used as the positive control. The results showed that compound **1** exhibited high cytotoxicity against PC3 and SHSY5Y cell with IC_{50} values of 3.5 and 2.4 μM , respectively.

ACKNOWLEDGEMENTS

This research was supported by the National Natural Science Foundation of China (21362044), the Excellent Scientific and Technological Team of Yunnan High School (2010CI08), the Yunnan University of Nationalities Green Chemistry and Functional Materials Research for Provincial Innovation Team (2011HC008), the National Undergraduates Innovating Experimentation Project (2011HX18) and start-up funds of Yunnan University of Nationalities.

REFERENCES

- M.M. Pinto, M.E. Sousa and M.S. Nascimento, *Curr. Med. Chem.*, **12**, 2517 (2005).
- K.S. Masters and S. Bräse, *Chem. Rev.*, **112**, 3717 (2012).
- C.M. Cao, H. Zhang, R.J. Gallagher and B.N. Timmermann, *Planta Med.*, **79**, 697 (2013).
- Y.P. Wu, W. Zhao, Z.Y. Xia, G.H. Kong, X.P. Lu, Q.F. Hu and X.M. Gao, *Molecules*, **18**, 9663 (2013).
- S. Klaiiklay, Y. Sukpondma, V. Rukachaisirikul and S. Phongpaichit, *Phytochemistry*, **85**, 161 (2013).

TABLE-1
 ^1H AND ^{13}C NMR DATA OF COMPOUND **1** (δ IN ppm, 500 AND 125 MHz, IN $\text{C}_5\text{D}_5\text{N}$)

No.	δ_{C} (m)	δ_{H} (m, J, Hz)	No.	δ_{C} (m)	δ_{H} (m, J, Hz)
1	161.5 s		4a	155.0 s	
2	98.6 d	6.62 s	8a	109.2 s	
3	161.4 s		9a	104.6 s	
4	108.3 s		10a	158.4 s	
5	110.5 d	6.82 d (8.3)	1'	34.3 t	2.50 t (7.2)
6	135.2 d	7.49 t (8.3)	2'	63.2 t	3.62 t (7.2)
7	108.6 d	6.68 d (8.3)	3-OMe	55.9 q	3.80 s
8	162.0 s		1-Ar-OH		13.46 s
9	182.5 s		8-Ar-OH		13.14 s

6. H.R. Dharmaratne, Y. Sakagami, K.G. Piyasena and V. Thevanesam, *Nat. Prod. Res.*, **27**, 938 (2013).
7. S. Udomchotphruet, P. Phuwapraisirisan, J. Sichaem and S. Tip-Pyang, *Phytochemistry*, **73**, 148 (2012).
8. C. Uvarani, K. Chandraprakash, M. Sankaran, A. Ata and P.S. Mohan, *Nat. Prod. Res.*, **26**, 1265 (2012).
9. M. Ali, M. Arfan, M. Ahmad, K. Singh, I. Anis, H. Ahmad, M.I. Choudhary and M.R. Shah, *Planta Med.*, **77**, 2013 (2011).
10. Q.B. Han, N.Y. Yang, H.L. Tian, C.F. Qiao, J.Z. Song, D.C. Chang, S.L. Chen, K.Q. Luo and H.X. Xu, *Phytochemistry*, **69**, 2187 (2008).
11. Q. Hu, X. Gao, D. Niu, X. Li, Y. Qin, Z. Yang, G. Zhao, Z. Yang and Z. Chen, *Heterocycles*, **87**, 1127 (2013).
12. N. Tanaka, Y. Kashiwada, S.Y. Kim, M. Sekiya, Y. Ikeshiro and Y. Takaishi, *Phytochemistry*, **70**, 1456 (2009).
13. Z.Y. Xiao and Q. Mu, *Nat. Prod. Res. Dev.*, **19**, 344 (2007).
14. S. Abe, N. Tanaka and J. Kobayashi, *J. Nat. Prod.*, **75**, 484 (2012).
15. N. Tanaka, S. Abe, K. Hasegawa, M. Shiro and J. Kobayashi, *Org. Lett.*, **13**, 5488 (2011).
16. W. Wang, Y.H. Zeng, K. Osman, K. Shinde, M. Rahman, S. Gibbons and Q. Mu, *J. Nat. Prod.*, **73**, 1815 (2010).
17. Y.P. Wu, W. Zhao, Z.Y. Xia, G.H. Kong, X.P. Lu, Q.F. Hu and X.M. Gao, *Phytochem. Lett.*, **6**, 629 (2013).
18. X.M. Gao, R.R. Wang, D.Y. Niu, C.Y. Meng, L.M. Yang, Y.T. Zheng, G.Y. Yang, Q.F. Hu, H.D. Sun and W.L. Xiao, *J. Nat. Prod.*, **76**, 1052 (2013).