NOTE

A New Phenolic Amide from Stems of *Cassia fistula* and Their Anti-Tobacco Mosaic Virus Activities

GUIYOU LIU, JUANXIA YANG, HUAN WANG, JIE LOU, LIMEI LI, XUEMEI GAO, QIUFEN HU and YANQING YE*

Key Laboratory of Chemistry in Ethnic Medicinal Resources, State Ethnic Affairs Commission & Ministry of Education, Yunnan University of Nationalities, Kunming 650031, P.R. China

*Corresponding author: E-mail: yey-qing@163.com

Received: 30 October 2013;

Accepted: 25 March 2014;

Published online: 1 September 2014;

AJC-15905

A new phenolic amide, (E)-3-(3,4-dihydroxyphenyl)-N-[2-(4-methoxyphenyl)-2-oxoethyl]-prop-2-enamide (**1**) was isolated from the stems of *Cassia fistula*. Its structure was elucidated by spectroscopic methods, including extensive 1D- and 2D-NMR techniques. Compound **1** was also evaluated for its anti-tobacco mosaic virus (anti-TMV) activity and it exhibit potential anti-TMV activity with inhibition rates of 16.2 %.

Keywords: *Cassia fistula*, Phenolic amide, Anti-tobacco mosaic virus activity.

Cassia fistula L., (Leguminosae) is an ornamental tree with beautiful yellow flowers¹. In China, it has been used as traditional Chinese medicine by people of Dai nationality, who lived in Xishuangbanna, Yunnan province for treatment of diarrhea, gastritis, ringworm and fungal skin infections^{2,3}. Previous phytochemical studies of *C. fistula* have shown the presence of anthraquinones^{4,5}, steroids⁶, chromones^{7,8} and flavonoids⁹. Motivated by a search for new bioactive metabolites from this plant, our research group has investigated the chemical constituents of the stem of *C. fistula*, which led to the isolation and characterization of a new phenolic amide. This paper deals with the isolation, structural characterization and the anti-tobacco mosaic virus (anti-TMV) activity of this compound.

UV spectra were obtained on a Shimadzu UV-2401A spectrophotometer. A Tenor 27 spectrophotometer was used for scanning IR spectroscopy with KBr pellets. 1D and 2D NMR spectra were recorded on a DRX-500 NMR spectrometer with TMS as internal standard. Unless otherwise specified, chemical shifts (δ) are expressed in ppm with reference to the solvent signals. HRESIMS was performed on a VG Autospec-3000 spectrometer. Semipreparative HPLC was performed on a Shimadzu LC-8A preparative liquid chromatograph with Zorbax PrepHT GF (21.2 mm $\times$ 25 cm) or Venusil MP C₁₈ (20 mm $\times$ 25 cm) columns. Column chromatography was performed using silica gel (200-300 mesh, Qing-dao Marine Chemical, Inc., Qingdao, P.R. China), Lichroprep RP-18 gel (40-63 μ m, Merck, Darmstadt, Germany) and MCI gel

(75-150 μ m, Mitsubishi Chemical Corporation, Tokyo, Japan). The fractions were monitored by TLC and spots were visualized by heating silica gel plates sprayed with 5 % H₂SO₄ in EtOH.

The stems of *C. siamea* were collected in Honghe prefecture of Yunnan Province, People's Republic of China, in September 2012. The identification of plant material was verified by Prof. Ning Yuan. A voucher specimen (Ynni-12-09-64) has been deposited in our Laboratory.

Extraction and isolation: The air-dried and powdered *C. siamea* (2.2 kg) were extracted four times with methanol (4 $\times$ 3 L) at room temperature and filtered. The filtrate was evaporated under reduced pressure and the crude extract (63.5 g) was decolorized by MCI. The 90 % methanol part (31.2 g) was chromatographed on a silica gel column eluting with a chloroform-acetone gradient system (20:1, 9:1, 8:2, 7:3, 6:4, 5:5), to give six fractions A-F. The further separation of fraction C (8:2, 5.47 g) by silica gel column chromatography, eluted with petroleum ether-acetone (9:1-1:2), yielded mixtures C1-C6. Fraction C3 (7:3, 0.82 g) was subjected to silica gel column chromatography using petroleum ether-acetone and semi-preparative HPLC (48 % MeOH-H₂O, flow rate 12 mL/min) to give **1** (8.5 mg).

(E)-3-(3,4-Dihydroxyphenyl)-N-[2-(4-methoxyphenyl)-2-oxoethyl]-prop-2-enamide (1): Obtained as white powder; IR (KBr, $\nu_{\max}$, cm⁻¹): 3410, 3312, 2915, 2586, 1675, 1638, 1527, 1455, 1232, 1168, 1121, 862, 793; UV (MeOH), $\lambda_{\max}$ (log ϵ): 215 (4.20), 245 (2.56), 292 (4.08), 322 (3.46); ¹H and

^{13}C NMR spectroscopic data (500 and 125 MHz, CDCl_3), see Table-1; ESI-MS (positive mode) m/z 350 $[\text{M} + \text{Na}]^+$; HRESIMS (positive mode) m/z 350.1011 $[\text{M} + \text{Na}]^+$ (Calcd. for $\text{C}_{18}\text{H}_{17}\text{NNaO}_5$, 350.1004).

The stems of *C. siamea* were extracted with methanol. The extract was subjected repeatedly to column chromatography on silica gel, RP-18 and semi-preparative RP-HPLC separation to afford compound **1**. Its structure was shown in Fig. 1. The ^1H and ^{13}C NMR data of the compound **1** was listed in Table-1.

Fig. 1. Structure of compound **1**

Table-1 ^1H AND ^{13}C NMR DATA OF COMPOUND 1 (δ in ppm, in CDCl_3 , 500 AND 125 MHz)					
No.	δ_{C}	δ_{H} Mult., (J Hz)	No.	δ_{C}	δ_{H} Mult., (J Hz)
1	127.5 s		2',6'	130.3 d	7.90 d (8.8)
2	114.8 d	6.92 d (1.8)	3',5'	115.2 d	6.76 d (8.8)
3	145.6 s		4'	163.8 s	
4	148.2 s		7'	193.3 s	
5	116.6 d	6.78 d (8.6)	8'	45.6 t	4.65 d (5.8)
6	121.5 d	6.85 d (8.6, 1.8)	-NH		8.29 t (5.8)
7	140.8 d	7.25 d (16.0)	-OMe	55.9 q	3.76 s
8	118.2 d	6.52 d (16.0)	Ar-OH-3		9.62 s
9	166.5 s		Ar-OH-4		9.53 s
1'	127.6 s				

Compound (**1**) was isolated as a white powder. Its molecular formula $\text{C}_{18}\text{H}_{17}\text{NO}_5$ was determined from the quasi-molecular ion peak observed using electrospray ionization (ESI)-MS and HRESI-MS measurement at m/z 350.1011 $[\text{M} + \text{Na}]^+$ (Calcd. for $\text{C}_{18}\text{H}_{17}\text{NNaO}_5$, 350.1004), suggesting a 11 degrees of unsaturation. Strong absorption bands accounting for hydroxy (3410 cm^{-1}), amino (3312 cm^{-1}), carbonyl (1675 cm^{-1}) and aromatic group ($1638, 1527, 1455\text{ cm}^{-1}$) could be observed in its IR spectrum. The UV spectrum of **1** showed absorption maxima at 215, 245, 292 and 322 confirmed the existence of the aromatic function. The ^1H and ^{13}C NMR spectrum of **1** revealed an caffeic acid moiety (C-1- C-9; H-2, H-5, H-6, H-7 and H-8) and an 2-amino-1-(4-methoxyphenyl) ethanone moiety (C-1'-C-8'; H-2',6', H-3',5', H-8' and NH) (Table-1). The NMR data of **1** were similar to those of tribulusamide D¹⁰, the major difference being the replacement of a hydroxy proton signal in tribulusamide D by a methoxy signal (δ_{C} 55.9, δ_{H} 3.76) in **1**. The HMBC correlation (Fig. 2) of the methoxy proton (δ_{H} 3.76) with C-4' (δ_{C} 163.8) indicated that the methoxy group was located at C-4'. Compound **1** is therefore the 4'-methoxy derivative of tribulusamide D.

Fig. 2. HMBC correlations () of **1**

Since certain of the phenolic compounds exhibit potential anti-TMV activities¹¹⁻¹³, compound **1** was tested for its anti-TMV activity. The inhibitory activity of compound **1** against TMV replication were tested using the half-leaf method^{12,13}. Ningnanmycin (a commercial product for plant disease in China, with inhibition rate of 29.5 %) was used as a positive control. The result showed that compound **1** exhibited potential anti-TMV activity with inhibition rate in the range of 16.2 %.

ACKNOWLEDGEMENTS

This research was supported by the National Natural Science Foundation of China (No. 21302164), the Excellent Scientific and Technological Team of Yunnan High School (2010CI08), the Yunnan University of Nationalities Green Chemistry and Functional Materials Research for Provincial Innovation Team (2011HC008) and Open Research Fund Program of Key Laboratory of Ethnic Medicine Resource Chemistry (Yunnan University of Nationalities) (2010XY08).

REFERENCES

- V. Duraipandiyan and S. Ignacimuthu, *J. Ethnopharmacol.*, **112**, 590 (2007).
- S. Rajan, D.S. Baburaj, M. Sethuraman and S. Parimala, *Ethnobotany*, **6**, 19 (2001).
- J. Ma, L.X. Zhang and Y.H. Guan, *Chin. J. Ethnomed. Ethnopharm.*, **5**, 178 (2004).
- K.A. Abo, A.A. Adeyemi and A.O. Sobowale, *Afr. J. Med. Med. Sci.*, **30**, 9 (2001).
- S. Aurapa and G. Wandee, *Int. J. Biomed. Pharm. Sci.*, **3**, 42 (2009).
- P. Sartorelli, S.P. Andrade, M.S. Melhem, F.O. Prado and A.G. Tempone, *Phytother. Res.*, **21**, 644 (2007).
- Y.H. Kuo, P.H. Lee and Y.S. Wein, *J. Nat. Prod.*, **65**, 1165 (2002).
- S.L. Jothy, Z. Zakaria, Y. Chen, Y.L. Lau, L.Y. Latha, L.N. Shin and S. Sasidharan, *Molecules*, **16**, 7583 (2011).
- W. Zhao, X.Y. Zeng, T. Zhang, L. Wang, G.Y. Yang, Y.K. Chen, Q.F. Hu and M.M. Miao, *Phytochem. Lett.*, **6**, 179 (2013).
- S.S. Hong, W. Jeong, J.G. Kwon, Y.H. Choi, E.K. Ahn, H.J. Ko, D.W. Seo and J.S. Oh, *Bull. Korean Chem. Soc.*, **34**, 3105 (2013).
- Q.-F. Hu, G. Yang, X. Li, X. Yang, H. Mu, Y. Chen and X.-M. Gao, *Heterocycles*, **85**, 147 (2012).
- Q.F. Hu, B. Zhou, X.M. Gao, L.Y. Yang, L.D. Shu, Y.Q. Shen, G.P. Li, C.T. Che and G.Y. Yang, *J. Nat. Prod.*, **75**, 1909 (2012).
- Q.F. Hu, B. Zhou, J.M. Huang, X.M. Gao, L.D. Shu, G.Y. Yang and C.T. Che, *J. Nat. Prod.*, **76**, 292 (2013).