Synthesis and Characterization of Methoxy-Substituted Salamo-Type Bisoximes Based on *Bis*(aminoxy)alkane and 3-Methoxy-2-hydroxybenzaldehyde

ZONG-LI REN¹, SU-XIA GAO², YANG ZHANG¹, LI WANG¹ and XIU-YAN DONG^{1,*}

¹School of Chemical and Biological Engineering, Lanzhou Jiaotong University, Lanzhou 730070, P.R. China

²School of Environmental and Municipal Engineering, Lanzhou Jiaotong University, Lanzhou 730070, P.R. China

*Corresponding author: E-mail: dxy568@163.com

Received: 25 February 2014;

Accepted: 5 May 2014;

Published online: 25 September 2014;

AJC-16044

Four new methoxy-substituted Salamo-type bisoxime compounds 6,6'-dimethoxy-2,2'-[(propylene-1,3-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L¹**), 6,6'-dimethoxy-2,2'-[(butylene-1,4-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L²**), 6,6'-dimethoxy-2,2'-[(propane-1,5-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L³**) and 6,6'-dimethoxy-2,2'-[(hexane-1,6-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L⁴**) have been synthesized through condensation reactions under hot ethanol medium, respectively and characterized by elemental analyses, IR, UV-visible and ¹H NMR spectra.

Keywords: Synthesis, Salamo-type bisoxime compound, Characterization.

INTRODUCTION

Salen-type compounds are a kind of linear tetradentate, N₂O₂-coordinating ligands which are obtained by condensation reaction of ketones or aldehydes with diamine. They are used extensively in the catalysis activity, biological systems, sterilization and antivirus in the last century¹⁻⁵. Recently, a preferable class of Salen-type bisoxime ligands was reported, using an *O*-alkyloxime unit [-CH=N-O-(CH₂)_n-O-N=CH-] instead of the [-CH=N-(CH₂)_n-N=CH-] group. The large electronegativity of oxygen atoms is expected to affect strongly the electronic properties of the N₂O₂ coordination sphere, which can lead to different and novel properties and structures of the resulting complexes⁶⁻⁹. The new Salen-type compounds and their complexes play an important role in the development of coordinating chemistry¹⁰⁻¹². In this article, we report the synthesis of four new Salen-type bisoxime compounds from 3-methoxy-2-hydroxy-benzaldehyde and several *bis*(aminoxy)alkane with different long-chain alkane molecules, 6,6'-dimethoxy-2,2'-[(propylene-1,3-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L¹**), 6,6'-dimethoxy-2,2'-[(butylene-1,4-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L²**), 6,6'-dimethoxy-2,2'-[(propane-1,5-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L³**) and 6,6'-dimethoxy-2,2'-[(hexane-1,6-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (**H₂L⁴**) and their characterizations.

EXPERIMENTAL

3-Methoxy-2-hydroxy-benzaldehyde (≥98 %), 1,3-dibromopropane, 1,4-dibromobutane, 1,5-dibromopentane and

1,6-dibromohexane were purchased from Alfa Aesar and used without further purification. The other reagents and solvents were analytical grade reagents from Tianjin Chemical Reagent Factory. The physico-chemical measurements are the same as previous literature¹³.

Synthetic route to Salamo-type bisoxime compounds **H₂L¹**-**H₂L⁴** are shown in Fig. 1.

1,3-*bis*(Aminoxy)propane, 1,4-*bis*(aminoxy)butane, 1,5-*bis*(aminoxy)pentane and 1,6-*bis*(aminoxy)hexane were synthesized according to an analogous method reported earlier¹³.

Preparation of 6,6'-dimethoxy-2,2'-[(propylene-1,3-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (H₂L¹**):** To an ethanolic solution (5 mL) of 3-methoxy-2-hydroxy-benzaldehyde (608.6 mg, 4 mmol) was added an ethanolic solution (5 mL) of 1,3-*bis*(aminoxy)propane (212.2 mg, 2 mmol). After the solution had been stirred at 55 °C for 3 h, then cooled to room temperature. The formed white precipitate was separated by filtration and washed successively with ethanol and ether. The product was dried under vacuum to yield the compound **H₂L¹**, Yield, 87.7 %, m.p. 437-438 K.

Preparation of 6,6'-dimethoxy-2,2'-[(butylene-1,4-diylldioxy)*bis*(nitrilomethylidyne)]diphenol (H₂L²**):** To an ethanolic solution (5 mL) of 3-methoxy-2-hydroxybenzaldehyde (608.6 mg, 4 mmol) was added an ethanolic solution (5 mL) of 1,4-*bis*(aminoxy)butane (240.3 mg, 2 mmol). After the solution had been stirred at 55 °C for 3 h, then cooled to room temperature. The formed white precipitate was separated by filtration and washed successively with ethanol and ether.

Fig. 1. Synthetic route to Salamo-type bisoxime compounds H_2L^1 - H_2L^4 ($n = 2-5$)

The product was dried under vacuum to yield the compound H_2L^2 , Yield, 68.8 %, m.p. 387-388 K.

Preparation of 6,6'-dimethoxy-2,2'-[(propane-1,5-diylidyoxy)bis(nitrilome-thylidyne)]diphenol (H_2L^3): To a ethanolic solution (5 mL) of 3-methoxy-2-hydroxybenzaldehyde (152.2 mg, 1 mmol) was added an ethanolic solution (2 mL) of 1,5-bis(aminooxy)pentane (67.1 mg, 0.50 mmol). After the solution had been stirred at 55 °C for 4 h, then cooled to room temperature. The light yellow precipitate was filtered and washed successively with a mixture of ethanol and *n*-hexane (*v:v* = 1:4) and *n*-hexane, respectively. The product was dried under vacuum to yield the compound H_2L^3 , Yield, 63.2 %, m.p. 372-373 K.

Preparation of 6,6'-dimethoxy-2,2'-[(hexane-1,6-diylidyoxy)bis(nitrilomethylidyne)]diphenol (H_2L^4): To a hot ethanolic solution (5 mL) of 3-methoxy-2-hydroxybenzaldehyde (152.2 mg, 1 mmol) was added an ethanolic solution (2 mL) of 1,6-bis(aminooxy)hexane (74.1 mg, 0.50 mmol). After the solution had been stirred at 55 °C for 4 h, then cooled to room temperature, the white precipitate was filtered and washed successively with a mixture of ethanol and *n*-hexane (*v:v* = 1:4) and *n*-hexane, respectively. The product was dried under vacuum to yield the compound H_2L^4 , Yield, 82.6 %, m.p. 359-360 K.

RESULTS AND DISCUSSION

A series of new Salamo-type bisoxime compounds H_2L^1 - H_2L^4 have been synthesized and the composition are confirmed by elemental analyses, IR, UV-visible and 1H NMR spectra.

Physical and chemistry properties: The colour, yields, melting points and elemental analytical results of the synthesized Salamo-type bisoxime compounds H_2L^1 - H_2L^4 are presented in Table-1.

Their compositions agree with the suggested molecular formula. Compounds H_2L^1 - H_2L^4 are either white or light yellow

microcrystalline solid. All the compounds are stable in air and soluble in acetone, methanol, ethanol, chloroform, dichloromethane, tetrahydrofuran, ethyl acetate, DMF and DMSO, insoluble in water, ether, acetonitrile and *n*-hexane.

IR spectra: The key IR spectral bands for Salamo-type bisoxime compounds H_2L^1 - H_2L^4 are given in Table-2.

TABLE-2
KEY IR SPECTRAL BANDS (cm^{-1}) FOR SALAMO-TYPE BISOXIME COMPOUNDS H_2L^1 - H_2L^4

Comp.	$\nu(O-H)$	$\nu(CH_2)$	$\nu(C=N)$	$\nu(C=C)$	$\nu(Ar-O)$
H_2L^1	3443	2942, 2887	1606	1586, 1473, 1429	1255
H_2L^2	3438	2936, 2885	1605	1582, 1474, 1427	1259
H_2L^3	3441	2941, 2868	1607	1596, 1475, 1425	1263
H_2L^4	3433	2947, 2885	1605	1593, 1474, 1428	1259

In the IR spectra of Salamo-type bisoxime compounds H_2L^1 - H_2L^4 , the O-H stretching band of Salamo-type bisoxime compounds H_2L^1 - H_2L^4 is expected in the 3800-3300 cm^{-1} region. However, this frequency is generally displaced to about 3433 cm^{-1} due to the internal hydrogen bond $OH \cdots N=C^{14}$. Thus, each unit of the polymeric ligands behaves as a dibasic tetradentate ONNO donor. As the hydrogen bond becomes stronger, the band width increases and this band sometimes is not detected. Electron-donating groups on the phenolic ring increase the electron density on the hydroxyl oxygen making the H-O bond stronger, the absorption usually appears as a broad band in the FT-IR spectrum. Meanwhile, the characteristic $C=N$ stretching bands of Salamo-type bisoxime compounds H_2L^1 - H_2L^4 appear at 1608-1605 cm^{-1} , respectively, indicating that 3-methoxy-2-hydroxybenzaldehyde has been condensed with 1,3-bis(aminooxy)propane, 1,4-bis(aminooxy)butane, 1,5-bis(aminooxy)pentane and 1,6-bis(aminooxy)hexane, respectively and formed new Salamo-type bisoxime^{13,14} compounds H_2L^1 - H_2L^4 . In addition, the Ar-O stretching bands occur at 1263-1255 cm^{-1} as reported for similar bisoxime compounds¹³.

TABLE-1
COLOUR, YIELDS, MELTING POINTS AND ANALYTICAL DATA OF SALAMO-TYPE BISOXIME COMPOUNDS H_2L^1 - H_2L^4

Compounds	Colour	m.p. (K)	Yield (%)	m.f. (m.w.)	Elemental analysis (%): Found (Calcd.)		
					C	H	N
H_2L^1	White	437-438	87.7	$C_{19}H_{22}N_2O_6$ (374.3)	60.44 (60.95)	6.07 (5.92)	7.12 (7.48)
H_2L^2	White	387-388	68.8	$C_{20}H_{24}N_2O_6$ (388.4)	61.74 (61.84)	6.53 (6.23)	7.16 (7.21)
H_2L^3	White	372-373	63.2	$C_{21}H_{26}N_2O_6$ (402.4)	63.87 (63.99)	7.46 (7.52)	6.48 (6.42)
H_2L^4	White	359-362	82.6	$C_{22}H_{28}N_2O_6$ (416.5)	64.55 (64.56)	7.67 (7.65)	6.27 (6.26)

UV-visible spectra: The UV-visible absorption spectra of Salamo-type bisoxime compounds $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$ (Table-3) in diluted dichloromethane solution show that the spectra of Salamo-type bisoxime compounds $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$ are similar to each other. Salamo-type bisoxime compounds $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$ exhibit two intense peaks at around 273 and 318 nm. The former absorption peaks at about 273 nm can be assigned to the $\pi\text{-}\pi^*$ transition of the benzene rings, while the latter absorption peaks at about 318 nm can be attributed to the intra-ligand $\pi\text{-}\pi^*$ transition of the $\text{C}=\text{N}$ bonds¹⁵. It is of note that there was no absorption around 400 nm, which are seen in the corresponding Salen derivatives. The absorption is ascribed to the quinoid form of $\text{H}_2\text{salen}^{16}$.

TABLE-3
UV-VISIBLE SPECTRAL DATA FOR THE
SALAMO-TYPE BISOXIME COMPOUNDS $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$

Comp.	C ($\times 10^{-5}$ mol L ⁻¹)	First band	Second band
		λ_{max1} (nm)	λ_{max2} (nm)
H_2L^1	5.00	273	318
H_2L^2	5.00	275	318
H_2L^3	5.00	279	322
H_2L^4	5.00	282	325

^1H NMR spectra: The ^1H NMR spectra of the title compounds $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$ in $\text{DMSO-}d_6$ are shown in Table-4. The ^1H NMR spectra showed a singlet at about 8.22-8.28 ppm indicating the existence of oxime bonds¹⁶.

TABLE-4
 ^1H NMR DATA FOR SALAMO-TYPE BISOXIMES $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$

Comp.	^1H NMR (400 MHz, $\text{DMSO-}d_6$, δ/ppm)
H_2L^1	2.43-2.54 (m, 2H, CH_2), 3.89 (s, 6H, CH_3), 4.47 (s, 4H, $\text{CH}_2\text{-O}$), 6.80 (dd, $J = 7.9$, 2.0 Hz, 2H, PhH), 6.85 (t, $J = 7.7$ Hz, 2H, PhH), 6.90 (dd, $J = 7.8$, 1.6 Hz, 2H, PhH), 8.22 (s, 2H, $\text{N}=\text{CH}$), 9.70 (s, 2H, OH)
H_2L^2	2.44-2.54 (m, 4H, CH_2), 3.90 (s, 6H, CH_3), 4.48 (s, 4H, $\text{CH}_2\text{-O}$), 6.81 (dd, $J = 7.6$, 1.6 Hz, 2H, PhH), 6.86 (t, $J = 7.8$ Hz, 2H, PhH), 6.91 (dd, $J = 7.6$, 2.0 Hz, 2H, PhH), 8.26 (s, 2H, $\text{N}=\text{CH}$), 9.71 (s, 2H, OH)
H_2L^3	2.45-2.55 (m, 6H, CH_2), 3.90 (s, 6H, CH_3), 4.47 (s, 4H, $\text{CH}_2\text{-O}$), 6.80 (dd, $J = 7.6$, 1.7 Hz, 2H, PhH), 6.86 (t, $J = 8.2$ Hz, 2H, PhH), 6.92 (dd, $J = 7.7$, 1.7 Hz, 2H, PhH), 8.25 (s, 2H, $\text{N}=\text{CH}$), 9.72 (s, 2H, OH)
H_2L^4	2.45-2.56 (m, 8H, CH_2), 3.91 (s, 6H, CH_3), 4.50 (s, 4H, $\text{CH}_2\text{-O}$), 6.80 (dd, $J = 7.6$, 1.8 Hz, 2H, PhH), 6.85 (t, $J = 8.0$ Hz, 2H, PhH), 6.90 (dd, $J = 7.6$, 1.8 Hz, 2H, PhH), 8.28 (s, 2H, $\text{N}=\text{CH}$), 9.76 (s, 2H, OH)

Conclusion

In this work, a series of methoxy-substituted Salamo-type bisoxime compounds $\text{H}_2\text{L}^1\text{-H}_2\text{L}^4$ that have two oxime bonds instead of imine bonds have been designed and synthesized by the reaction of 2 equivalents of 3-methoxy-2-hydroxybenzaldehyde with 1,3-bis(aminooxy)propane, 1,4-bis(aminooxy)-butane, 1,5-bis(aminooxy)pentane and 1,6-bis(aminooxy)-hexane under hot ethanolic medium, respectively. It is shown that a bisoxime moiety is much more useful to assemble supramolecular systems than a Schiff base moiety. Further investigation on the synthesis and structures of methoxy-substituted Salamo-type complexes with transition metal ions, are now in progress.

ACKNOWLEDGEMENTS

This work was supported by the Fundamental Research Funds for the Gansu Province Universities (212086) and the science and technology support funds of Lanzhou Jiaotong University (ZC2012003), which are gratefully acknowledged.

REFERENCES

1. E.N. Jacobsen, *Acc. Chem. Res.*, **33**, 421 (2000).
2. D.J. Zhou, Q. Li, C.H. Huang, G.Q. Yao, U. Shigeo, M. Masakazu, L.M. Ying, A.C. Yu and X.S. Zhao, *Polyhedron*, **16**, 1381 (1997).
3. T.K. Ronson, H. Adams and M.D. Ward, *Inorg. Chim. Acta*, **358**, 1943 (2005).
4. N.S. Venkataramanan, G. Kuppuraj and S. Rajagopal, *Coord. Chem. Rev.*, **249**, 1249 (2005).
5. M. Sasaki, K. Manseki, H. Horiuchi, M. Kumagai, M. Sakamoto, H. Sakiyama, Y. Nishida, M. Sakai, Y. Sadaoka, M. Ohba and H. Okawa, *J. Chem. Soc., Dalton Trans.*, **259** (2000).
6. S.S. Sun, C.L. Stern, S.T. Nguyen and J.T. Hupp, *J. Am. Chem. Soc.*, **126**, 6314 (2004).
7. J.C.G. Bünzli and C. Piguet, *Chem. Rev.*, **102**, 1897 (2002).
8. S. Akine, T. Taniguchi and T. Nabeshima, *Chem. Lett.*, **30**, 682 (2001).
9. W.K. Dong, J.Y. Shi, J.K. Zhong, Y.Q. Tian and J.G. Duan, *Chinese J. Inorg. Chem.*, **24**, 10 (2008).
10. P.G. Lacroix, *Eur. J. Inorg. Chem.*, 339 (2001).
11. S. Akine, W.K. Dong and T. Nabeshima, *Inorg. Chem.*, **45**, 4677 (2006).
12. S. Akine, Y. Morita, F. Utsuno and T. Nabeshima, *Inorg. Chem.*, **48**, 10670 (2009).
13. (a) W.K. Dong, J.G. Duan, Y.H. Guan, J.Y. Shi and C.Y. Zhao, *Inorg. Chim. Acta*, **362**, 1129 (2009); (b) W.K. Dong, X.N. He, H.B. Yan, Z.W. Lv, X. Chen, C.Y. Zhao and X.L. Tang, *Polyhedron*, **28**, 1419 (2009); (c) W.K. Dong, L. Li, C.F. Li, L. Xu and J.G. Duan, *Spectrochim. Acta A*, **71**, 650 (2008); (d) W.K. Dong, Y.X. Sun, C.Y. Zhao, X.Y. Dong and L. Xu, *Polyhedron*, **29**, 2087 (2010).
14. J.A. Faniran, K.S. Patel and J.C. Bailar Jr., *J. Inorg. Nucl. Chem.*, **36**, 1547 (1974).
15. T. Ghosh, B. Mondal, T. Ghosh, M. Sutradhar, G. Mukherjee and M. Drew, *Inorg. Chim. Acta*, **360**, 1753 (2007).
16. H.E. Smith, *Chem. Rev.*, **83**, 359 (1983).