Microwave-Assisted Acetylation of Phenols without Catalyst Under Solvent Free Condition

YU-QUAN ZHANG^{1,*}, YAN-CHUN LI¹, JI-HUA ZHU¹, ZHI-FENG LI², GUO-ZHE GUO¹ and DONG-PING CHEN¹

¹College of Chemistry and Chemical Engineering, Longdong University, Lanzhou Road 54 Xifeng, Qingyang 745000, P.R. China

²College of Life Science and Chemistry, Tianshui Normal University, Tianshui 741001, P.R. China

*Corresponding author: E-mail: zhyq118@sohu.com

Received: 2 April 2014;

Accepted: 6 June 2014;

Published online: 6 November 2014;

AJC-16226

Etherification between phenols with acetic anhydride was tested under different conditions. Phenols were efficiently acylated with acetic anhydride to give phenol acetate derivatives in good high yields without catalyst and solventless conditions under microwave irradiation.

Keywords: Acetylation, Phenols, Microwave irradiation.

INTRODUCTION

The acylation of hydroxyl groups is one of the most frequently used transformations in organic synthesis. Among the various protecting groups used for the hydroxyl function, acetyl is the most common group in view of its easy introduction, being stable to the acidic reaction conditions and also easily removable by mild alkaline hydrolysis¹. Typically, acylation of alcohols and phenols is performed under homogeneous catalysis with acetic acid or acetyl chloride or anhydride, in the presence of a convenient basic catalyst such as triethylamine or pyridine². In addition, 4-(dimethylamino)pyridine, 4-pyrrolidinopyridine³, Tetramethylethylenediamine⁴, tertiary phosphines⁵, carbon tetrabromide in ethyl acetate⁶, *p*-toluenesulfonic acid⁷, zinc chloride⁸, iodine⁹, sulfamic acid¹⁰, magnesium bromide¹¹, cobalt chloride¹², tantalum chloride¹³, La(OPr-*i*)₃¹⁴, silica gel-supported sodium hydrogen sulfate¹⁵, montmorillonite K-10 and KSF¹⁶, alumina¹⁷, yttria zirconia with acetic acid¹⁸, Ac₂O-pyridine/basic alumina under microwave irradiation¹⁹, KF-Al₂O₃ with Ac₂O/AcCl²⁰, vanadyl(V) acetate²¹, distannoxane²², zeolite HSZ-360²³, Pseudo-monas cepacia PS lipase adsorbed on Celite²⁴, and twisted amides²⁵ have also been applied for the acetylation of alcohols and phenols. Another important development has been the introduction of trifluoromethanesulfonate derivatives, such as TMSOTf²⁶, Yb-(OTf)₃²⁷, Cu(OTf)₂²⁸, Sc(OTf)₃²⁹, In(OTf)₃³⁰, Bi(OTf)₃³¹, cerium(III)³² and (IV)³³ salts as efficient catalysts in acetylation with acetic acid, or anhydride or acetyl chloride. All these ways lead to wastes as well as some reactions involving solvents, often toxic, polluting and longer time.

There is an increasing interest in the use of environmentally benign reagents and conditions. Particularly to solvent free

procedures. Avoiding organic solvent during the reactions in organic synthesis leads to a clean, efficient and economical technology: safety is largely increased. Working is considerably simplified, cost is reduced increased amount of reactants, which can be used³⁴. Microwave irradiation has become increasingly popular in recent years to improve the yield and shorten reaction times in a variety of reactions³⁵⁻³⁷. Reactive and selective sometimes are enhanced. Some microwave irradiation assisted solventless acetylation were described using catalyses³⁸. Here we report acetylation of phenols using acetic anhydride as acetylating agent without catalyst and solventless conditions under microwave irradiation, as shown in **Scheme-I**:

Scheme-I

EXPERIMENTAL

All the reagents employed in the synthesis were of analytical reagent grade. The materials were used as purchased and used directly without any additional purification. Melting points were determined on an XT-4 electro thermal micro melting point apparatus and are uncorrected. Infrared spectra were measured as KBr discs (or liquid film) using an Alpha centaurspectrometer. ^1H NMR spectra (200 MHz) were recorded in CDCl_3 using a BRUKER PT 200 spectrometer. Mass spectra were obtained on a Qp-1000A GC-MS spectrometer using the electron impact mode (70 or 20 eV). Microwave irradiation was carried out with a domestic microwave oven GlanzWP 750B at 2450 Hz. Sonication was performed in Shanghai SK-250LH ultrasonic cleaner with the frequency of

59 kHz. $\text{KF/Al}_2\text{O}_3$ was prepared according to reference²⁰. The remaining chemicals were obtained from commercial sources.

General procedure: To a mixture of phenols (5 mmol) and acetic anhydride (6 mmol) were added in a bottle (50 mL) and irradiated at 675 W for 10 min in microwave oven. After cooling to room temperature, then reaction mixture was diluted with water (50 mL) and extracted with ethyl acetate (2×50 mL). The organic layer was washed with satd NH_4Cl solution (30 mL) and NaHCO_3 solution (25 mL) and brine (40 mL), respectively. The organic layer dried over MgSO_4 and concentrated to give the product.

RESULTS AND DISCUSSION

The results for a variety of esterification compounds are summarized in the Table-1, various types of phenols with electron donating and withdrawing groups were rapidly acylated with acetic anhydride and afforded the corresponding phenol acetate in excellent yields under microwave irradiation without catalyst and solventless conditions.

TABLE-1
RESULTS FOR A VARIETY OF ETHERIFICATION
COMPOUNDS WITHOUT CATALYSES AND COLVENTFREE
CONDITIONS UNDER MICROWAVE IRRADIATION^a

Product ^b	Phenols	Yields (%) ^c	m.p. °C/Lit
3a	Phenol	96	Oil ⁴¹
3b	4-Methyl phenol	95	Oil ⁴²
3c	3-Nitro phenol	90	52-53(53-54 ⁴¹)
3d	1,2-Catechol	94	62-63°C (62-63 °C ⁴³)
3e	1,4-Quinol	94	121-122(120-121 ⁴³)
3f	1,3,5-Triphenol	88	101-103(102-103 ⁴³)
3g	1-Naphthol	90	45-46(44-45 ⁴¹)
3h	2-Naphthol	92	68-69(67-69 ⁴¹)
3i	4-Nitro phenol	93	77-79(77-78 ⁴¹)

^aConditions: Aryl phenols (5 mmol), acetic anhydride (6 mmol), 675 W for 10 min. ^bAll products gave satisfactory ¹H NMR, Mass spectra and IR. ^cYield of isolated product

To explore the scope and versatility of this method, different reaction conditions were investigated. Highlighted in Table-2 for 2-naphthol acetate (**3h**), for example, is the influence of catalysts, reaction time, mole ratios, ultrasound wave and microwave on the reaction yield.

Since $\text{KF/Al}_2\text{O}_3$ ³⁹ is a new kind of highly active catalytic catalysts that can catalyze many reactions under mild conditions, to give high yields, the high efficacy and ease of product

isolation, a large number of ultrasonic reactions⁴⁰ can be carried out in higher yield, shorter reaction time or milder conditions, this prompted us to investigate its use for acetylation purposes.

To demonstrate the catalysis of $\text{KF/Al}_2\text{O}_3$, the reaction was carried out with high speed stirring, ultrasonic irradiation and microwave irradiation conditions in the absence of $\text{KF/Al}_2\text{O}_3$ catalyst, it was concluded that the acetylation of 2-naphthol with acetic anhydrides could not happen with high speed stirring at 135 °C and under ultrasonic irradiation. However, it was observed that compound **3h** (Table-2 entry 7) was obtained in 92 % yield in 10 min. When $\text{KF/Al}_2\text{O}_3$ was used to catalyze the reaction of 2-naphthol with acetic anhydrides, the acetylation of 2-naphthol with acetic anhydrides could not happen at 135 °C with high speed stirring and ultrasonic irradiation, while in the presence of $\text{KF/Al}_2\text{O}_3$ catalyst and microwave, compound **3h** was also obtained in 92 % yields in 10 min (Table-2 entry 2,3,4,5,6), these result revealed that microwave can effectively accelerate the reaction, $\text{KF/Al}_2\text{O}_3$ was no effective for acetylation of 2-naphthol with acetic anhydrides.

The different mole ratios of esterification of 2-naphthol with acetic anhydrides was studied under microwave irradiation, the results show the best mole ratio is 2-naphthol: acetic anhydrides = 1:1.2.

We investigated the effects of irradiation power and time on the reaction. It was found that the highest yield of compounds is obtained at a power level of 675 W for 10 min continuous irradiation.

Conclusion

In summary, a simple, safe and eco-efficient method for acetylation of phenols has been developed. The method has advantages in terms of yields, short reaction times, ease of operation and compatibility with other protecting groups. We believe that the present methodology could be an important addition to the existing methodologies.

ACKNOWLEDGEMENTS

The authors thank Scientific and Research Projects of Department of Education of Gansu Province for 2012 and the Longdong University Science Technology Development Foundation (XYZK0504) for financial support.

TABLE-2
OPTIMIZATION OF THE REACTION CONDITION

Entry	Catalyst (mmol %) $\text{KF/Al}_2\text{O}_3$	Method	Time (min)	^b Yields (%)
1	0.3	MWI	6	60 ^a
2	0.3	MWI	8	78 ^a
3	0.3	MWI	10	92 ^a (0 ^c)
4	0.3	MWI	12	89 ^a
5	0.5	MWI	10	92 ^a
6	0.7	MWI	10	92 ^a
7	0	MWI	10	92 ^a (0 ^c)
8	0.3	Ultrasound	40	(0 ^d)
9	0	Ultrasound	40	(0 ^d)

Conditions: 2-naphthol (5 mmol), acetic anhydride (6 mmol). ^aMicrowave irradiation at 675 W. ^bYield of isolated product. ^cReaction temperature 135 °C, high speed stirring reaction time 5 h, without microwave irradiation. ^dReaction temperature 20-25 °C, ultrasound irradiation at 59 kHz

REFERENCES

1. A.L. Pearson and W.J. Roush, *Handbook of Reagents for Organic Synthesis: Activating Agents and Protecting Groups*; John Wiley & Sons: UK, pp. 9-16 (1999).
2. R.I. Zhdanov and S.M. Zhenodarova, *Synthesis*, 222 (1975).
3. W. Steglich and G. Hofle, *Angew. Chem. Int. Ed. Engl.*, **8**, 981 (1969).
4. T. Sano, K. Ohaschi and T. Oriyama, *Synthesis*, 1141 (1999).
5. E. Vedejs, N.S. Bennett, L.M. Conn, S.T. Diver, M. Gingras, S. Lin, P.A. Oliver and M.J. Peterson, *J. Org. Chem.*, **58**, 7286 (1993).
6. H. Hagiwara, K. Morohashi, H. Sakai, T. Suzuki and M. Ando, *Tetrahedron*, **54**, 5845 (1998).
7. A.C. Cope and E.C. Herrick, *Org. Synth.*, **4**, 304 (1963).
8. R.H. Baker and F.G. Bordwell, *Org. Synth.*, **3**, 141 (1955).
9. R. Borah, N. Deka and J.C. Sarma, *J. Chem. Res. Synop.*, 110 (1997).
10. T.S. Jin, J.R. Ma, Z.H. Zhang and T.S. Li, *Synth. Commun.*, **28**, 3173 (1998).
11. E. Vedejs and O. Daugulis, *J. Org. Chem.*, **61**, 5702 (1996).
12. J. Iqbal and R.R. Srivastava, *J. Org. Chem.*, **57**, 2001 (1992).
13. S. Chandrasekhar, T. Ramachander and M. Takhi, *Tetrahedron Lett.*, **39**, 3263 (1998).
14. T. Okano, K. Miyamoto and K. Kiji, *Chem. Lett.*, **24**, 246 (1995).
15. G.W. Breton, *J. Org. Chem.*, **62**, 8952 (1997).
16. A.X. Li, T.S. Li and T.H. Ding, *Chem. Commun.*, 1389 (1997).
17. S.S. Rana, J.J. Barlow and K.L. Matta, *Tetrahedron Lett.*, **22**, 5007 (1981).
18. P. Kumar, R.K. Pandey, M.S. Bodas and S.P. Dagade, *J. Mol. Catal. Chem.*, **181**, 207 (2002).
19. P. Paul, P. Nanda, R. Gupta and A. Loupy, *Tetrahedron Lett.*, **43**, 4261 (2002).
20. V.K. Yadav, K.G. Babu and M. Mittal, *Tetrahedron*, **57**, 7047 (2001).
21. B.M. Choudary, M.L. Kantam and V. Neeraja, *J. Mol. Catal. Chem.*, **140**, 25 (1999).
22. A. Orita, K. Sakamoto, Y. Hamada, A. Mitsutome and J. Otera, *Tetrahedron*, **55**, 2899 (1999).
23. R. Ballini, G. Bosica, S. Carloni, L. Ciaralli, R. Maggi and G. Sartori, *Tetrahedron Lett.*, **39**, 6049 (1998).
24. P. Allevi, P. Ciuffreda, A. Longo and M. Anastasia, *Tetrahedron Asym.*, **9**, 2915 (1998).
25. S. Yamada, T. Sugaki and K. Matsuzaki, *J. Org. Chem.*, **61**, 5932 (1996).
26. P.A. Procopiou, S.P.D. Baugh, S.S. Flack and G.G.A. Inglis, *Chem. Commun.*, 2625 (1996).
27. A.G.M. Barrett and D. Christopher Braddock, *Chem. Commun.*, 351 (1997).
28. K.L. Chandra, P. Saravanan, R.K. Singh and V.K. Singh, *Tetrahedron*, **58**, 1369 (2002).
29. K. Ishihara, M. Kubota, H. Kurihara and H. Yamamoto, *J. Am. Chem. Soc.*, **117**, 4413 (1995).
30. K.K. Chauhan, C.G. Frost, I. Love and D. Waite, *Synlett*, 1743 (1999).
31. G. Bartoli, M. Bosco, E. Marcantoni, L. Sambri and E. Torregiani, *Synlett*, 209 (1998).
32. G. Bartoli, G. Cupone, R. Dalpozzo, A. De Nino, L. Maiuolo, A. Procopio, L. Sambri and A. Tagarelli, *Tetrahedron Lett.*, **43**, 5945 (2002).
33. K. Tanaka and F. Toda, *Chem. Rev.*, **100**, 1025 (2000).
34. P. Lidström, J. Tierney, B. Wathey and J. Westman, *Tetrahedron*, **57**, 9225 (2001).
35. J. Hamelin, J.-P. Bazureau, F. Texier-Boullet in ed.: A. Loupy, *Microwaves in Organic Synthesis*, Wiley-VCH, Weinheim, pp. 253-294 (2002).
36. (a) Y.-Q. Zhang, *J. Chem. Res.*, 365 (2013); (b) L. Bai, J.-X. Wang and Y.M. Zhang, *Green Chem.*, **5**, 615 (2003).
37. N. Deka, A.-M. Mariotte and A.Ä. Boumendjel, *Green Chem.*, **3**, 263 (2001).
38. T. Ando, J. Yamawaki, T. Kawate, S. Sumi and T. Hanafusa, *Bull. Chem. Soc. Jpn.*, **55**, 2504 (1982).
39. T.J. Mason and D. Peters, *Practical Sonochemistry*, Ellis-Horwood, London (2002).
40. M. Aloe, *Rend. Accad. Sci. Napoli*, **27**, 75 (1921).
41. F.D. Chattaway, *J. Chem. Soc.*, 2495 (1931).
42. J. Buckingham and S.M. Donghy, *Dictionary of Organic Compounds*, Chapman and Hall, New York, edn. 5 (1982).