CASE REPORT

GIANT APPENDICULAR MUCOCELE - A CASE REPORT

Muraliswar Rao J¹, Surabi Karthik²

HOW TO CITE THIS ARTICLE:

Muraliswar Rao J, Surabi Karthik. "Giant appendicular mucocele - A Case Report". Journal of Evolution of Medical and Dental Sciences 2013; Vol2, Issue 32, August 12; Page: 6119-6123.

ABSTRACT: Appendicular mucocele by definition is a cystic dilatation of the appendiceal lumen by mucin accumulation. This is a rare lesion; its prevalence in appendectomy specimens being only 0.2 - 0.3%. Mucocele can result from mucosal hyperplasia, mucinous cystadenoma, or mucinous cystadenocarcinoma. Specific criteria are being proposed for definitive diagnosis and surgical management of appendiceal mucocele. While some neoplasms with malignant potential may be treated definitively by resection. It is known to be associated with pseudomyxoma peritonei resulting from a rupture. It is therefore important to identify the disease process preoperatively and to plan a careful resection also to rule out possibility of malignancy. We report here one case with radiological, surgical and histopathological confirmation.

KEYWORDS: Appendicular mucocele, Mucinous cystadenoma, pseudomyxoma peritonei.

INTRODUCTION: Appendicular mucocele is a rare lesion. Cystic mass resulting from a dilated appendiceal lumen caused by abnormal accumulation of mucus, regardless of its underlying cause. Mucinous cystadenoma and cystadenocarcinoma account for 60 - 70% of all mucocele.^[1] The clinical presentation is usually non-specific with 50% of cases being an incidental finding at surgery.^[1] Symptoms could be an indeterminate abdominal pain or chronic or intermittent abdominal colicky pain.

CASE REPORT: 39yr old male patient presented with chief complaints of intermittent pain in the right lower lumbar region and iliac fossa since 5 months. On clinical examination abdomen was soft and on bimanual palpation in the Rt. Lower abdominal quadrant revealed soft to firm mass with regular borders, measuring 9x5.6cms in diameter which is slightly mobile.

Ultrasonography showed 9.8x5.6cms relatively well defined oblong heterogenous cystic mass with concentric onion peel internal echoes. There was no evidence of calcifications. On color and power doppler scanning no vascular signals were detected.

Single contrast barium enema showed a smooth marginated filling defect noted in the region of ceecal base.

On CT a well defined, well encapsulated mass was seen in the right lower quadrant posterior inferior to ceecal base noted in the region of the appendix with attenuation of around 15-20 HU.

Intraoperatively a tense cystic lesion of size 10x6 cms was found in the position of appendix.

Gross cut section appearance showed jelly like mucoid material with appendicular portion thickened and dilated. Microscopic features at the level of base of the appendix showed hypertrophy, hyperplastic muscle layer with focal areas of epithelium filled with mucinous material and sub-mucosal lymphoid aggregates. Section from the wall showed thickened muscle wall infiltrated by foamy histiocytes and inflammatory cells. Final impression was given as mucinous cystadenoma.

CASE REPORT

DISCUSSION: Mucocele of the appendix is a descriptive term for an appendix distended by mucus, secondary to mucinous cystadenoma (63%), mucosal hyperplasia (25%), mucinous cystadenocarcinoma (11%) and retention cyst.^[2]

Overall, appendiceal mucocele make up about 0.2%–0.3% of appendix specimen.^[3] Clinical presentation may include right lower quadrant pain, change in bowel habits, per rectal bleeding or a palpable mass.^[4] Approximately 23–50% of patients are asymptomatic, with the lesions being discovered incidentally during surgery^[3-5]. The preoperative clinical diagnosis of appendiceal mucocele can therefore be difficult because of this lack of clinical symptomatology.

The initial detection of the lesion may be facilitated by radiological, sonographic or endoscopic means.

On barium enema, the lesion may be seen as a sharply outlined sub mucosal or extrinsic mass indenting the caecum and laterally displacing it.^[3]

CT of the abdomen usually shows a cystic well-encapsulated mass sometimes with mural calcification, in the expected location of the appendix.^[3, 6-8]

Ultrasound findings can be variable. Purely cystic lesions with anechoic fluid, hypoechoic masses with fine internal echoes as well as complex hyperechoic masses can be seen depending on the contents.^[9] The onion skin sign is considered to be specific for mucocele of the appendix.^[10]

Colonoscopic findings include the 'volcano sign', the appendiceal orifice seen in the centre of a firm mound covered by normal mucosa or a yellowish, lipoma-like submucosal mass.^[11]

In the above case report, USG and CT were able to provide a preoperative diagnosis. In our case, the decision for excision of the appendiceal mucocele was made and a need to rule out malignancy. Surgical excision of mucocele of appendix can either be by laparotomy or laparoscopy.

However careful handling of the specimen is recommended as spillage of the contents can lead to pseudomyxoma peritonei. This can be achieved by atraumatic handling of the appendix and use of impermeable bag for removal of the specimen.^[12] Involvement of the caecum or adjacent organs is an indication for right hemi-colectomy and thorough exploration of the gastrointestinal tract and ovaries.^[13]

CONCLUSION: Appendicular mucocele though rare and mostly asymptomatic, is to be considered in the differential diagnosis of a right iliac fossa mass and CT scan is imperative in the correct preoperative diagnosis. This helps the surgeon to be more careful and it reduces the risk of iatrogenic rupture of the mucocele with resultant leakage of its contents into the abdominal cavity causing pseudomyxoma peritonei and to rule out possibility of malignancy as the cause of mucocele.

Image: 1: Single contrast barium enema shows well defined filling defect at the level of caecum.

CASE REPORT

Image: 2: Sonographic image showing well defined hypoechoic lesion with whorled internal echoes.

Image: 3: Axial CECT image shows well defined hypodense lesion posterior-inferior to caecum in the region of appendix.

Image: 4 & 5: Sagittal and coronal CECT images show a well-defined hypodense lesion in the right lower quadrant posterior-inferior to the caecum.

CASE REPORT

Image: 6 & 7: Gross specimen pictures shows soft to firm well defined mass with mucinous contents oozing out at cut sections.

Image: 8: Histopathological image shows tall mucinous epithelium with mucinous material within the lumen with thickened muscle wall infiltrated by foamy histiocytes and inflammatory cells.

REFERENCES:

1. Ekinadoese Juliana Aghahowa, Chandramouli Bharati, Muneera Al- Adwani. Appendicular Mucocoele - A Case Report. Kuwait Medical Journal 2008; 40 : 78-80
2. Higa E, Rosai J, Pizzimbono CA, Wise L. Mucosal hyperplasia, mucinous cystadenoma and mucinous cystadenocarcinoma of the appendix. A re-evaluation of appendiceal "mucocoele". Cancer. 1973; 32:1525-1541.
3. Dachman AH, Lichtenstein JE, Friedman AC. Mucocoele of the appendix and pseudomyxoma peritonei. AJR Am J Roentgenol. 1985;144:923-929
4. Aho AJ, Heinomen R, Laurén P. Benign and malignant mucocoele of the appendix. Histological types and prognosis. Acta Chir Scand. 1973;139:392-400
5. Soweid AM, Clarkston WK, Andrus CH, Jannet CG. Diagnosis and management of appendiceal mucocoele. Dig Dis. 1998; 16:183-186.
6. Kim SH, Lim HK, Lee WJ, Lim JH, Byun JY. Mucocoele of the appendix; ultrasonographic and CT findings. Abdom Imaging. 1998; 23:292-296.
7. Zissin R, Gayer G, Kots E, Apter S, Peri M, Sharipo-Feinberg M. Imaging of mucocoele of the appendix with emphasis on the CT findings: a report of 10 cases. Clin Radiol. 1999; 54:826-832.

CASE REPORT

8. Madwen D, Mindelzun R, Jeffrey RB. Mucocele of the appendix: Imaging findings. AJR Am J Roentgenol. 1992; 159:69–72.
9. Skaane P, Ruud TE, Haffner J. Ultrasonographic features of mucocele of the appendix. J Clin Ultrasound. 1998; 16:584–587.
10. Caspi B, Cassif E, Auslender R, Herman A, Hagay Z, Appelman Z. The onion skin sign: a specific sonographic marker of appendiceal mucocele. J Ultrasound Med. 2004; 23:117–121.
11. Hamilton DL, Stormont JM. The volcano sign of appendiceal mucocele. Gastrointest Endosc. 1989; 35:453–456.
12. Navarra G, Asopa V, Basaglia E, Jones M, Jiao LR, Habib NA. Mucous cystadenoma of the appendix: is it safe to remove it by a laparoscopic approach? Surg Endosc. 2003; 17:833–4.
13. Kahn M, Friedman JH. Mucocele of the appendix: Diagnosis and surgical management. Dis Colon Rectum. 1979; 22:267–269.

AUTHORS:

1. Muraliswar Rao J.
2. Surabi Karthik

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of Radiodiagnosis, ASRAM Medical College.
2. Post Graduate, Department of Radiodiagnosis, ASRAM Medical College.

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Muraliswar Rao J,
Department of Radiodiagnosis,
ASRAM Medical College,
Elluru – 534001, West Godavari (dt),
Andhra Pradesh, India.
Email – muraliradiology@gmail.com

Date of Submission: 02/08/2013.
Date of Peer Review: 05/08/2013.
Date of Acceptance: 06/08/2013.
Date of Publishing: 12/08/2013.