CHRONIC OBSTRUCTIVE PULMONARY DISEASE AND ITS RISK FACTORS: A CASE CONTROL STUDY.

Manoj Kumar Meghwani¹, Harish Chandra Tiwari², Richa Mishra³, Shrawan Kumar⁴

HOW TO CITE THIS ARTICLE:

Manoj Kumar Meghwani, Harish Chandra Tiwari, Richa Mishra, Shrawan Kumar. "Chronic obstructive pulmonary disease and its risk factors". Journal of Evolution of Medical and Dental Sciences 2013; Vol2, Issue 34, August 26; Page: 6382-6386.

ABSTRACT: Chronic obstructive pulmonary disease (COPD) has been defined as a disease characterized by airflow limitation that is not fully reversible. It is a major health problem in India in terms of morbidity and mortality as well. In a resource-poor country like India, it is a challenge to curb the growing burden of COPD. To understand the risk factors that lead to the development of COPD in our community, a case control study was carried out at Rama Medical College, Hospital & Research Center (RMCH & RC) Mandhana, Kanpur. A total of 323 COPD cases & 325 controls were selected among the patients attending the department of tuberculosis & chest disease of this medical college. Diagnosis of COPD was made by using the standard clinical criteria and was further confirmed by spirometry, chest X-ray and Non Contrast Computed Tomography (NCCT) thorax as and when necessary. After obtaining informed consent from each patient a detailed clinical history regarding age, gender, usual place of residence, smoking habits and type of cooking fuel combustion at home was enquired by using predesigned questionnaire. Our studies showed that proportion of male were higher in case group than in controls. Significantly higher proportions of cases of COPD were having history of smoking as compared to controls. COPD patients had smoked higher mean pack-years as compared to controls. Exposures to environment tobacco smoke (ETS) and active smoking both were found as most important risk factor for the development of COPD. Other important risk factors found in this study were exposure to solid fuel smoke at home especially in females.

KEY WORDS: COPD, Smoking, Environment tobacco smoke (ETS).

INTRODUCTION: Chronic obstructive pulmonary disease (COPD) has been defined as a disease characterized by airflow limitation that is not fully reversible. It includes namely emphysema, an anatomically defined condition characterized by destruction & enlargement of lung alveoli; chronic bronchitis, a clinically defined condition with chronic cough & phlegm; and a small airway disease, a condition in which small bronchioles are narrowed¹.

Chronic Obstructive Pulmonary Disease (COPD) is the 4th largest cause of death in the world². According to the World Health Organization, COPD kills more people than HIV-AIDS, Malaria and Tuberculosis all put together in the South East Asian region. COPD is a major health problem in India in term of morbidity and mortality as well. Its prevalence is underestimated due to lack of uniform diagnostic criteria.

According to a report published by the National Centre for Macroeconomics and Health, it was estimated that economic loss due to COPD in India is around Rs 35,000 crores³. In a resource-poor country like India, it is a challenge to curb the growing burden of COPD. Therefore the prevention of development of COPD becomes the most promising option to reduce the burden of COPD.

ORIGINAL ARTICLE

For this, we need to understand the various risk factors that lead to the development of COPD. Keeping all this in mind present study was undertaken to understand the role of risk factors such as the tobacco smoking, biomass fuel combustion and other environmental exposures for occurrence of COPD.

MATERIAL & METHOD: The present case control study was carried out at RMCH & RC Mandhana Kanpur. The COPD cases attending to the department of tuberculosis & chest disease between October 2012 to July 2013 were included in this study. Only patients aged ≥ 35 year & willing to participate were included in the study. Geriatric, obstetric, psychiatric hospitalization and cases of rehabilitation were excluded from our study. COPD was diagnosed by using the standard criteria for chronic bronchitis, i.e. presence of cough with expectoration for more than three months in a year for the past two or more years excluding other causes of chronic cough with expectoration. Diagnosis was also confirmed by Spirometry, chest X-ray P.A. view and Non Contrast Computed Tomography (NCCT) thorax as and when necessary. After suitable matching to the cases, controls were identified from patients visiting to same department with other complaints.

After obtaining informed consent from each patient a detailed history of each patient was taken. The variables used for analysis included the age, gender, usual place of residence, smoking habits and the type of cooking fuel combustion at home. A smoker was defined by the presence of regular smoking of any type i.e. cigarettes, bidis or hookah, for one year or more. Exposure to cooking fuel combustion was considered positive if the individual gave the history of regularly cooking at home. The types of cooking fuels used at home included liquefied petroleum gas (LPG), kerosene, or the solid fuels i.e. coal, dried wood, dung and other products of animal or plant origin (biomass fuels).

Data collected was analyzed by using SPSS software- 17 version. Differences in baseline characteristics between COPD patients and control subjects were analyzed using chi square tests for categorical variables.

RESULTS:

Table 1– Characteristics of cases of COPD patients & control subjects

		Cases		Controls		Total	
		N	%	N	%	N	%
Age group	35-50 Year	124	38.4%	102	31.4%	226	34.8%
	51- 60 Year	107	33.1%	117	36.0%	224	34.6%
	61-70 year	92	28.5%	106	32.6%	198	30.6%
Sex	Male	208	64.4%	176	54%	384	59.2%
	Female	115	35.6 %	149	46%	264	40.8%
Total		323	100%	325	100%	648	100%

In the present study, 323 (49.8%) of COPD cases & 325 (50.2%) of controls were included. All cases as well as controls were of almost same age group with slightly higher percentage of cases (38.4%) were of age group 35-40 year as compared to controls (31.4%). There were significantly higher proportions of male (64.4%) among the cases than controls (54%). (Table 1).

Table 2– History of smoking among COPD cases & control subjects

	Cases		Controls		Total		X ² , P-value
	N	%	N	%	N	%	
Never smoker	56	17.3%	96	29.5%	152	23.4%	X ² = 14.6, df=1, p= <0.001
Ex smoker	96	29.7%	93	28.6%	189	29.1%	
Current smoker	171	52.9%	136	41.8%	307	47.3%	
Mean pack year	26		14		20		
Total	323	100%	325	100%	648	100%	

No history of smoking was found in 17.3% of cases of COPD as compared to 29.5% among controls. Among cases 29.7% were ex- smoker and 52.9% current smokers while among controls 28.6% were ex- smoker and 41.8% current smokers. Significantly higher percentages of cases of COPD were having history of smoking as compared to controls. (X² = 14.6, df=1, p= <0.001). COPD patients had smoked higher mean pack-years (26) as compared to controls (14). (Table 2)

Table 3– History of exposure of dust, environment tobacco smoke (ETS) among COPD cases & Control subjects

		Cases		Controls		Total		X ² , P-value
		N	%	N	%	N	%	
History of ETS	Yes	174	53.8%	118	36.3%	292	45.1%	X ² = 20.2, df=1, p=<0.01
	No	149	46.1%	207	63.7%	356	54.9%	
Current exposure of ETS at home	Yes	109	33.7%	83	25.5%	192	29.6%	X ² = 5.23, df=1, p=<0.05
	No	214	66.3%	242	74.5%	456	71.4%	
Current exposure of ETS at work	Yes	78	24.1%	67	20.6%	145	22.4%	X ² = 1.16, df=1, p=>0.05
	No	245	75.8%	258	79.4%	503	77.6%	
Exposure to solid fuel smoke at home	Yes	153	47.4%	106	32.6%	259	39.9%	X ² = 14.7, df=1, p=<0.01
	No	170	52.6%	219	63.4%	389	60.1%	
Total		323	100%	325	100%	648	100%	

Exposures to environment tobacco smoke (ETS) were significantly higher among COPD patients (53.8%) than control subjects (36.3%), (X²= 20.2, df=1, p=<0.01). Current exposures to ETS, both at home & work were slightly more prevalent among COPD patients than control subjects, although the differences were not statistically significant for current history of ETS at work (X²= 1.16, df=1, p=>0.05). Exposure to solid fuel smoke at home were significantly more prevalent among COPD patients (47.4%) than control subjects (32.6%) (X²= 14.7, df=1, p=<0.01).

DISCUSSION: The present study showed that proportion of male (64.4) was higher in case group than control group (54%) significantly. This is consistent with the study of S. K. Jindal who reported that COPD is distinctly more common amongst men⁴. We found in our study that significantly higher

percentages of cases of COPD were having history of smoking as compared to controls. In our study we found 82.7% of COPD patients were smoker (current/past) while among controls it is 70.5%. COPD patients had smoked higher mean pack-years (26) as compared to controls (14). Our study reveals that smoking is a major risk factor for COPD is consistent with various other studies.^{5, 6, 7}

Exposures to environment tobacco smoke (ETS) were found significantly higher among COPD patients than control subjects in this study. Although it is widely acknowledged that active smoking is by far the most important independent risk factor for the development of COPD but environmental exposure to tobacco has also been shown to be important risk factor for development of COPD in present study. Recent results from the European Community Respiratory Health Survey showed that adult lung function and susceptibility to COPD were partly determined early in life⁸. The results from the present study are in consistent with these observations. Apparently, tobacco smoking's as well as the ETS exposures are important risk factors for COPD prevalence.

Other important risk factor among nonsmoker women was exposure to solid fuel smoke at home was also found in this study. The present study revealed that exposure to solid fluid smoke and dust was significantly more prevalent among COPD patients than control subjects. Similar finding were also reported by Ane Johannessen et al⁹ in their study.

REFERENCES:

1. John J Reilly, Jr., Edwin K Silverman, Steven D Shapiro. Chronic Obstructive Pulmonary Disease. Harrisons Principle of Internal Medicine; 16th edition Vol. II, 1547.
2. Sundeep Salvi, Anurag Agrawal. India Needs a National COPD Prevention and Control Programme. JAPI 2012; 60:4-5.
3. Murthy KJR, Sastry JG. Economic burden of chronic obstructive pulmonary disease: NCMH Background Papers- Burden of Disease in India 2005
4. S K Jindal. COPD: The Unrecognized Epidemic in India. Supplement to JAPAI. February 2012; 60:4-5.
5. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management and prevention of chronic obstructive pulmonary disease. Updated 2003. National Institutes of Health, National Heart, Lung, and Blood Institute.
6. US Department of Health and Human Services. The health consequences of smoking: A Report of the Surgeon General, Atlanta, GA: US Department of Health and Human Services, CDC, Office on Smoking and Health 2004.
7. Reddy KS, Gupta PC. Report on Tobacco Control in India. Ministry of Health and Family Welfare, Government of India. New Delhi 2004.
8. Svanes C, Sunyer J, Plana E et al. Early life origins of chronic obstructive pulmonary disease. Thorax 2010; 65: 14-20.
9. Ane Johannessen. Association of exposure to environmental tobacco smoke in childhood with chronic obstructive pulmonary disease and respiratory symptoms in adults. Respirology (2012) 17, 499-505.

ORIGINAL ARTICLE

AUTHORS:

1. Manoj Kumar Meghwani
2. Harish Chandra Tiwari
3. Richa Mishra
4. Shrawan Kumar

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of Tuberculosis and Chest Diseases, RMCH & RC, Kanpur.
2. Assistant Professor, Department of Community Medicine, RMCH & RC, Kanpur.
3. Senior Resident, Department of Community Medicine, IMS, BHU, Varanasi.
4. Associate Professor, Department of Medicine, RMCH & RC, Kanpur.

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Harish Chandra Tiwari,
Assistant Professor,
Department of Community Medicine,
Rama Medical College, Hospital & Research Center,
G.T. Road, Mandhana,
Kanpur, UP – 209217.
Email – dr.harishchandratiwari@gmail.com

Date of Submission: 16/08/2013.
Date of Peer Review: 17/08/2013.
Date of Acceptance: 19/08/2013.
Date of Publishing: 20/08/2013