CAN MRI REPLACE DIAGNOSTIC ARTHROSCOPY IN EVALUATION OF INTERNAL DERANGEMENT OF KNEE JOINT – A PROSPECTIVE STUDY

Laxman Prasad Ahirwar¹, Chandraprakash Ahirwar²

HOW TO CITE THIS ARTICLE:

Laxman Prasad Ahirwar, Chandraprakash Ahirwar. "Can MRI replace diagnostic arthroscopy in evaluation of internal derangement of knee joint – a prospective study". Journal of Evolution of Medical and Dental Sciences 2013; Vol2, Issue 40, October 07; Page: 7621-7631.

ABSTRACT: OBJECTIVE: To evaluate the role of MRI in patients clinically suspected to have internal derangement of knee joint and to compare the MRI results with Arthroscopy. **METHODS:** This prospective study was done in the Department of Radiodiagnosis M.G.M. Medical College & M.Y. Hospital, Indore. Patients referred from orthopedic OPD with suspected internal derangement (IDK) of the knee were investigated with MRI. Total 100 patients were included in the study. After completion of MRI examination, arthroscopy was performed in cases wherever it was indicated. Arthroscopy was used as a gold standard and findings of these 100 cases were compared with MRI Results: It was found that MRI has a diagnostic accuracy of 90%, 90%, 93.5%, 100% and 74.4% for detecting medial meniscus, lateral meniscus, and anterior cruciate, posterior cruciate and osteochondral injuries respectively. **CONCLUSION:** MRI has a high diagnostic value in the diagnosis of meniscal and cruciate injuries. MRI is highly accurate in diagnosis of complete ACL tear and PCL tear. MRI is less sensitive than arthroscopy in detecting early chondral injuries.

KEYWORDS: MRI knee, internal derangement of knee (IDK), knee arthroscopy.

INTRODUCTION: Internal Derangement of Knee (IDK) is the term used for a group of disorders involving disruption of the normal functioning of the ligaments or cartilages (menisci) of the knee joint. In the setting of injury to knee, it is not always possible to diagnose the meniscal or ligamentous injury clinically as the clinical tests have their limitations.– Arthroscopy, considered as the gold standard, is invasive, expensive and requires day surgery admission. In today's era it has become a therapeutic surgical procedure, rather than a diagnostic tool.

MRI is currently gold standard non-invasive diagnostic imaging modality of choice for nearly all clinical indications concerning the knee. MRI provides excellent visualization of the internal structures of the knee along with soft tissue structures and bone marrow abnormalities.

MATERIALS AND METHODS: This prospective study was done in the Radiodiagnosis Department & SRL Diagnostic centre of Mahatma Gandhi Memorial Medical College & M.Y. Hospital, Indore, and Madhya Pradesh, India in a period between August 2009 to July 2011. A total of 100 patients, who were referred to our department with strong clinical suspicion of internal derangements of knee joint, underwent magnetic resonance imaging evaluation of knee followed by arthroscopy in selected cases, wherever indicated. Patients with neoplasm, inflammatory or infectious disorders and patients who had previously undergone arthroscopy with repair of ligaments & menisci were excluded from study. After completion of MRI examination, arthroscopy was performed in cases wherever it indicated. Arthroscopy was used as a gold standard and MRI findings of these cases were compared with MRI.

ORIGINAL ARTICLE

OBSERVATIONS AND RESULTS: Present study is a prospective clinical hospital based study comprising 100 patients of suspected internal derangements of knee joint (IDK). Arthroscopy evaluation was performed in 31 patients.

Most common age group of patients presenting with IDK was from 21-30 years, constituting 31% with mean age 24.3 years. Males were majority (76 %).

Pain was most common presenting complaint followed by swelling, constituting 79% and 54% respectively. Patients presented with overlapping symptoms in most of the cases. On clinical examination, most common positive clinical test was McMurray's test seen positive in 48% of cases followed by Lachman's Test constituting 26% of patients.

Our study reveal that the meniscal tear was the most common cause of IDK followed by cruciate ligament injuries (Table-1)

S. NO.	LESION	NO. OF CASES	% OF CASES
1.	MEDIAL MENISCAL TEAR	49	49%
2.	LATERAL MENISCAL TEAR	18	18%
3.	ANTERIOR CRUCIATE LIGAMENT TEAR	32	32%
4.	POSTERIOR CRUCIATE LIGAMENT TEAR	7	7%
5.	MEDIAL COLLETERAL LIGAMENT TEAR	11	11%
6.	LATERAL COLLETERAL LIGAMENT TEAR	5	5%
7.	CARTILAGE DEFECT	19	19%

Table 1: Lesions comprising internal derangement of knee joint

MEDIAL MENISCAL INJURES: Out of 100 patients of IDK, 67(67%) patients had meniscal tear. Of these, 49(73%) patients had medial meniscal tear and 18 (27%) patients had lateral meniscal tear. Out of 49 medial meniscal tears, 32(66%) involved posterior horn, 3(5%) involved anterior horn, 6 (12%) involved body and 8 (16%) involved entire meniscus. Out of 49 medial meniscal tears, 8 (22.2%) were classified as grade 1, 8 (10.2%), as grade 2 and 30(73.4%) as grade 3 (Fig.1). Out of 36 grade-3 medial meniscal tears, 9 (25%) tears were classified as horizontal tears , 4(11.1%) as vertical , and 15(41.6%) complex tears , 3(8.3%)-bucket handle , 3(8.3%) radial tear and 2(5.5%) as flap tear (Fig.2) (Table-2).

Arthroscopy was performed in 31 patients. Preoperative MRI of these patients showed grade -3 medial meniscal tears in 17 patients. On arthroscopy medial meniscal tears found in 16 patients. 2 cases diagnosed as tear on MRI found to be normal on arthroscopy (false positive) and 1 case which was normal on MRI found to be torn on arthroscopy (false negative). In our study the positive predictive value , negative predictive value, sensitivity, specificity and accuracy for detecting medial meniscal tears were 88%, 92%, 88%, 86% and 90% respectively.

ORIGINAL ARTICLE

SL NO	TYPE OF TEAR	NO. OF CASES	% OF CASES
1.	HORIZONTAL	9	25%
2.	VERTICAL	4	11.1%
3.	COMPLEX	15	41.6%
4.	RADIAL	3	8.3%
5.	BUCKET HANDLE	3	8.3%
6.	FLAP	2	5.5%
	TOTAL	36	

Table 2: Distribution of morphological types of medial meniscal tears

LATERAL MENISCAL INJURIES: Lateral meniscal tears were reported in 18(18%) out of these, 12 (66.6%) were involved posterior horn, 1(5.5%) anterior horn, 3(16.6%) body and 2 (11.1%) involving entire meniscus. Of the 18 lateral meniscal tears, grade-3 was most common (15 tears, 83.3%) following 2(11.1%) as grade 1 tear and 1 (5.5%) as grade 2. Of the 15 grade -3 lateral meniscal tears, 2 (13.3%) were classified as horizontal, 6(40%) vertical, 3(20%) complex, 2(13.3%) bucket handle tear, 1(6.6%) radial tear and 1(6.6%) as flap tear (Table-3). Arthroscopy was performed in 31 patients. Preoperative MRI of these patients reported grade -3 tears in 6. On arthroscopy, lateral meniscal tears found in 5 patients. 2 cases diagnosed as tear on MRI found converted to normal on arthroscopy (false positive) and 1 case which was normal on MRI found to be torn on arthroscopy (false negative). In our study the positive predictive value , negative predictive value, sensitivity, specificity and accuracy for detecting lateral meniscal tears were 66.6%, 96%, 92%, 88% and 90% respectively.

SLNO	TYPE OF TEAR	NO. OF CASES	% OF CASES
1.	HORIZONTAL	2	13.3%
2.	VERTICAL	6	40%
3.	COMPLEX	3	20%
4.	RADIAL	1	6.6%
5.	BUCKET HANDLE	2	13.3%
6.	FLAP	1	6.6%
	TOTAL	15	

Table 3: Distribution of morphological types of lateral meniscal tears

Cruciate ligament injuries: Out of 100 patients of IDK, 39(39%) patients had cruciate ligament tear. 32 (82%) had ACL tears and only 7 (18%) had PCL tear. Out of 32 ACL tear, 25 (78%) were classified as complete tear and 7 (22%) as partial tear (Fig.3). 11(34.3%) of ACL tears were located in proximal segment, 17 (53.2%) in mid substance and 4 (12.5%) in distal segment of ligament. Arthroscopy was performed in 31 patients. Preoperative MRI reported ACL tear in 14, and 13 patients proven as tear on arthroscopy. 1 was normal on arthroscopy (false positive) and 1 case which were normal on MRI found to be torn on arthroscopy (false negative). Hence PPV, NPV, sensitivity, specificity and accuracy for detecting anterior cruciate ligament tears were 92.8%, 94.1%, 92.8%, 94.1% and 93.5% respectively.

Of the 7 PCL tear, 3(42.8%) were classified as complete ligament tear, 1 (14.4%) as partial ligament tear and 3(42.8%) tibial avulsion (Fig. 4). Arthroscopy was performed in 31 patients. Preoperative MRI revealed posterior cruciate ligament tears in 2 patients, both of them found torn on arthroscopy. The PPV, NPV, sensitivity, specificity and accuracy in our study for detecting PCL tears were 100%.

Chondral injuries: Chondral defect were found in 19/100(19%) patients. Arthroscopy was performed in 31 patients. Preoperative MRI reported chondral defects in 7 patients. On arthroscopy chondral defects found in 9 patients. 3 cases diagnosed as defect on MRI found to be normal on arthroscopy (false positive) and 5 cases which was normal on MRI found to be injured on arthroscopy (false negative). The PPV, NPV, sensitivity, specificity and accuracy for detecting chondral defects were 57%, 79%, 44.4%, 86% and 74.4% respectively.

Collateral ligament injuries: MCL tear (11%) was more common than LCL tear (5%). Out of 11 patients of MCL tear, 4 (36%) classified as grade 1 tear, 5 (45%) as grade 2 tear and 2 (18%) as grade 3 tears (Fig.5). out of 5 LCL tear, 1 classified as grade-1, 1 as grade-2 and 3 tears as grade 3. O' Donoghue's triad (ACL with medial meniscal and MCL tear) was seen in 1 patient.

Other associated findings: Joint effusion (73%) was the most common associated finding with bone contusion/marrow oedema seen in 17 patients followed by fracture of bone seen in 9 patients (Table-4). Muscle injuries were seen in 6 patients. Popliteus muscle was most commonly injured. Patellar retinaculum tears were seen in 4 patients and patellar tendon tears were seen in 2 patients. Meniscal cyst was seen in 2 patients and all they involved the posterior horn of lateral meniscus. 2 patients had popliteal cyst (Baker's cyst). Discoid meniscus was present in 2 patients (Fig.6).

SL NO	LESION	NO. OF CASES	% OF CASES
1.	JOINT EFFUSION	73	73%
2.	FRACTURE	9	9%
3.	MARROW OEDEMA	17	17%
4.	MUSCLE CONTUSION	6	6%
5.	PATELLAR RETINACULUM TEAR	4	4%
6.	PATELLAR TENDON TEAR	2	2%
7.	MUCOID DEGENERATION OF ACL	3	3%
8.	DISCOID MENISCUS	2	2%
9.	CYSTIC LESIONS	4	4%

Table 4: Distribution of other associated Findings

DISCUSSION: Knee joint being the most complex weight bearing joint of the body is subject to damage because of its inherent structural complexity and the various types of forces it is subjected to. Disease processes and injuries that disrupt ligaments, menisci, articular cartilage and other structures of the knee cause significant morbidity and disability. MRI has emerged as the frontline investigation for evaluation of IDK. In this study, we attempt to demonstrate diagnostic value of MRI in IDK including various most common injuries of knee and were results compared with the arthroscopy findings.

This is a hospital based prospective study included 100 patients. Most of the patients were adults (20-40 yrs). Similar observations were seen by Vassilios S Nikolaou et al ¹.

In our study, most common cause of IDK was meniscal tear. The medial meniscus was most commonly found torn, because MM is less mobile, and it bears more force during weight bearing than the LM. Most of the meniscal tear were located in posterior horn. Crues et al ² in their study also found meniscal tears involving the posterior horns which accounts for 57% compared to the 16% involving the anterior horn. Weiss et al ³ also reported meniscal tears involving the posterior horn accounting for 50%-60% and tears involving the anterior horn accounting for 5%- 20%. D Smet et al ⁴ also found same result in their study. Most common morphological meniscal tear was complex type followed by horizontal. This is similar to the study by De Smet et al ⁴, in their study of MM tears in 343 patients found complex type of medial meniscus tear as a most common 116 patients.

In our study, 2 false positive MRI involved the posterior horn of the medial meniscus. On retrospective analysis of MRI it was found that in one case, presence of intra-meniscal tear was not communicating with the articular surface of the meniscus and it was misinterpreted as grade-3 meniscal tear. In second case, the exact cause of the false positives diagnosis of tear was not apparent. It may be attributed to the misinterpretation of normal menisiofemoral ligament as meniscal tear or operator/procedure dependant drawback of arthroscopy. This is similar to the study by McKenzie et al ⁵ who described the four most common reasons for false positive diagnosis; wrong diagnosis due to variable anatomic structures, overestimation of pathology countered as meniscus tear, false negative arthroscopic findings and tears within the meniscus without expansion to the articular surface. On retrospective analysis of MRI in one false negative case it was found that the signal intensity of the tear was misinterpreted to reflect a transverse ligament. This is similar to the observation seen by Mesgarzadeh et al ⁶. Another Study which was conducted by McKenzie et al ⁵ stated that false negative results exclusively occurred from misinterpretation of MRI.

The posterior horn was most commonly found torn in LM tears with vertical morphologically type. This is similar to the study by Naranje S et al ⁷ in their study; they found vertical type of lateral meniscus tear as a most common tear (53%) in all LM tears.

In our study, the two false positive MRI involved the posterior horn of the LM. On retrospective analysis of MRI, it was found that in one case, presence of menisiofemoral ligament was misinterpreted as meniscal tear. In second case, the hiatus of the popliteus tendon was mistaken as the tear. Similar observations were seen by Mesgarzadeh et al ⁶ in their study. On retrospective analysis of MRI in one false negative case, it was found that the signal intensity of the tear was communicating with the articular surface of the meniscus and it was misinterpreted as grade-2 meniscal tear. This is similar to the observation seen by McKenzie et al ⁵.

The ACL was more common ligament found injured than PCL because PCL is thick and strong. This is similar to the study by Vassilios S et al ¹ in their study of 26 patients with cruciate ligament tear; they found ACL tears in 23 (88%) patients and PCL tear in 3 (12%) patients. J. P. Singh et al ⁸ also found same results in their study. In ACL tears complete tear were more common than partial tear. Most of the complete tears were located in mid substance of ligament. This is similar to the study by J. P. Singh et al ⁸ in their study of 78 patients with ACL; they found anterior cruciate ligament tears most commonly located in mid substance of the ligament (67%). In our study, the one false positive case diagnosed on MRI as partial tear of anterior cruciate ligament. On retrospective analysis of MRI abnormal signal intensity was seen within the ligament with intact fibers. Vassilios S et al ¹ observed

that abnormal signal in the ligament, in absence of tear may occur due to intra body mucosal or eosinophilic degeneration of the ACL.

On retrospective analysis of MRI in one false negative case it was found that abnormal signal intensity of the ligament misinterpreted as mucoid degeneration of the ACL, but on arthroscopy chronic degenerated partial tear was seen with in the ligament. Patrice W J Vincken et al ⁹ showed that MRI has relatively less accuracy in diagnosis of partial ACL tear.

In posterior cruciate ligament injuries, tibial avulsion was the most common injury. These results are in concordance with the observations seen by J S Grover et al ¹⁰. In our study the diagnostic accuracy for detecting PCL tears was 100%. This is similar to the study of J S Grover et al ¹⁰ and C W Heron et al ¹¹ who found 100% sensitivity and specificity of MRI in diagnosis of PCL.

Chondral defects were seen in one fifth of the patients; however diagnostic accuracy of MRI was lower for chondral injuries in comparison to the meniscal and cruciate ligament injuries. In our study the diagnostic accuracy for detecting chondral defects was 74.4% which was corresponding to the study of Ochi M et al ¹² and Vassilios S et al ¹ who found accuracy of MRI significantly inferior in the diagnosis of chondral lesions. Heron et al ¹¹ described that MRI can satisfactory reveal the advanced chondral defects as well as damages at the patellar articular cartilage, but is not accurate for smaller injuries like fibrilization or small fissuring in articular hyaline cartilage.

In collateral ligaments tears, in our study, MCL tears (11%) were found to be more common than the LCL tear (5%). All these cases had history of trauma and were associated with multiple injuries. This suggests presence of a single injury should prompt the examiner to look for other subtle associated injuries, which was further confirmed by Mink JH et al ¹³. They observed on MRI and arthroscopy of 11 patients who had tear of ACL, 7 patients had tear of MCL, 4 patients had tear of lateral meniscus and 1 patient had tear of medial meniscus.

Joint effusion was associated with most of the positive cases. Bone marrow edema was the most common associated osseous injury. Pattern of bony injury was associated with specific type of injury in many patients. 7 patients with injury of the posterolateral tibial plateau and 4 patients with lateral femoral condyle found to be associated with ACL tear. This is similar to the observation seen by Robertson et al ¹⁴ in their study of multiple signs of anterior cruciate ligament on MR imaging in 103 patients found that posterolateral tibia bruise associated with ACL tear had 53% sensitivity, 97% specificity and 79% accuracy. The presence of lateral femoral bruise with ACL tear had a sensitivity of 47%, specificity of 97% and an accuracy of 76%.

Overall MRI has high diagnostic value in the diagnosis of meniscal and cruciate injuries (table-5). MRI is advantageous in conditions where arthroscopy is not useful like peripheral meniscal tears and inferior surface tears. MRI is more sensitive in detection of multiple meniscal tears that may be overlooked on arthroscopy. MRI is more sensitive than arthroscopy in detection of grade I and II intrasubstance degeneration, precursors to formation of meniscal tears. Many anatomic variants can mimic tears on MRI and false diagnosis can be made due to misinterpretation of these variant. MRI is highly accurate in diagnosis of complete ACL tear and PCL tear however it is less sensitive than arthroscopy in detecting partial ACL tears and early chondral injuries.

ORIGINAL ARTICLE

Test Accuracy	Medial Meniscus	Lateral meniscus	ACL	PCL	Chondral defect
Positive predictive value	88%	66.6%	92.8%	100%	57%
Negative predictive value	92%	96%	94.1%	100%	79.1%
Accuracy	90%	90%	93.5%	100%	74.15
Sensitivity	88%	80%	92.8%	100%	44.4%
Specificity	86%	92%	94.1%	100%	86%

Table 5: Distribution of diagnostic accuracy of MRI in various injuries

CONCLUSION: MRI is a useful non-invasive modality having high sensitivity, specificity and accuracy in the diagnosis of meniscal and cruciate ligament injuries. MRI should be done in every patient of suspected internal derangement of knee joint, to save a patient from unnecessary arthroscopy. So now diagnostic arthroscopy has no role in the management of internal derangement of knee joint and arthroscopy should be performed only for therapeutic purposes.

REFERENCES:

1. Vassilios S Nikolaou, corresponding author1 Efstathios Chronopoulos, 2 Christianna Savvidou, 2 Spyros Plessas, 2 Peter Giannoudis, 1 Nicolas Efstathopoulos, 2 and Georgios Papachristou, MRI efficacy in diagnosing internal lesions of the knee: a retrospective analysis. Journal of Trauma Management & Outcomes 2008 June 2. doi: 10.1186/1752-2897-2-4
2. Crues J V, Stoller DW, Martin C, Kaplan L and Mink JH – Meniscal tears: pathologic correlation with MR imaging Radiology 1987; 163: 731-735.
3. Weiss CB, Lundberg M, Hamberg P. Non operative treatment of meniscal tears. J Bone Joint Surg Am 1989; 71A: 811-822.
4. D Smet, David H Nathan, Ben K Graf, Benjamin A. Haaland and Jason P. Fine. Clinical and MRI Findings Associated with False-Positive Knee MR Diagnoses of Medial Meniscal Tears. AJR: July 2008; 191:93-99.
5. Mckenzie R, Palmer CR, Lomas DJ, and Dixon AK: Magnetic resonance imaging of the knee: diagnostic performance studies. Clin Radiol 1996, 51(4):251-257.

6. Mesgarzadeh M, Moyer R, Leder DS. MR Imaging of the knee; expanded classification and pitfalls to interpretation of meniscal tears. Radiographics 1993; 13: 489-500.
7. Naranje S, Mittal R et al Arthroscopic and magnetic resonance imaging evaluation of meniscus lesions in the chronic anterior cruciate ligament-deficient knee. Department of Orthopaedics, All India Institute of Medical Sciences, New Delhi, India. 2000 Dec; 103(12):1079-85.
8. Singh J P, Garg L, Shrimali R, Setia V, Gupta V, MR Imaging of knee with arthroscopic correlation in twisting injuries. Indian J Radiol Imaging 2004; 14: 33-40.
9. Patrice W. J. Vincken, MD. Bert P. M. ter Braak, MD. Effectiveness of MR Imaging in Selection of Patients for Arthroscopy of the Knee. Radiology 2002; 223:739-746.
10. Grover JS, Bassett LW, Gross ML, Seeger LL, Finerman GA. Posterior cruciate ligament: MR imaging Radiology 1990; 174(2): 527-530.
11. C W Heron. Three-dimensional gradient-echo MR imaging of the knee: comparison with arthroscopy in 100 patients. June 1992 Radiology, 183,839-844.
12. Ochi M, Sumen Y – The diagnostic value and limitation of magnetic resonance imaging on chondral lesions in the knee joint. Department of Orthopedic Surgery, Hiroshima University School of Medicine, Japan. Arthroscopy.1994 Apr; 10(2):176-83.
13. Mink JH: The cruciate and collateral ligaments, in Mink JH, Reicher MA, Crues JV III, (eds): MRI of the Knee (ed 2). New York, NY, Raven, 1993, pp 141-188.
14. Robertson L, Schweitzer ME, Bartolozzi AR, et al. Anterior cruciate ligament tears:evaluation of multiple signs with MR imaging. Radiology 1994; 193:829-834.

Abbreviation:- MRI- Magnetic resonance imaging, IDK- internal derangement of knee, MM- medial meniscus, LM- lateral meniscus , PCL- posterior longitudinal ligament, ACL- anterior cruciate ligament, LCL- lateral collateral ligament, MCL- medial collateral ligament, CD- chondral defect, PPV- positive predictive value, NPV- negative predictive value.

Fig. 1: MENISCAL TEARS GRADING

Fig. 2: MENISCAL TEARS – MORPHOLOGICAL TYPES

HORIZONTAL TEAR (COR PD FAT SAT)

RADIAL TEAR (SAG PD FAT SAT)

COMPLEX TEAR (SAG PD FAT SAT)

BUCKET HANDLE TEAR (COR PD FAT SAT)

Fig. 3: ACL TEAR

PARTIAL

(SAG PD FAT SAT)

(OBL CORONAL T2WI)

COMPLETE

(SAG PD FAT SAT)

(OBL CORONAL T2WI)

Fig. 4: PCL TEAR

PARTIAL (SAG PD)

COMPLETE (SAG PD FAT SAT)

AVULSION (SAG PD FAT SAT)

Fig. 5: COLLATERAL LIGAMENT TEAR

LCL TEAR (COR PD FAT SAT)

MCL TEAR (COR PD FAT SAT)

Fig.6: DISCOID MENISCUS (CORONAL PD FAT SAT)

AUTHORS:

1. Laxman Prasad Ahirwar
2. Chandraprakash Ahirwar

PARTICULARS OF CONTRIBUTORS:

1. Resident, Department of Radiodiagnosis, MGM Medical College, Indore.
2. Assistant Professor, Department of Radiodiagnosis and Imaging, Gandhi Medical College and Hamidia Hospital, Bhopal, India.

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Chandraprakash Ahirwar,
A-83, Housing board Colony (HBC),
Koh-e-fiza, Bhopal.
Email- drchandraprakashradiologist@gmail.com

Date of Submission: 18/09/2013.
Date of Peer Review: 20/09/2013.
Date of Acceptance: 25/09/2013.
Date of Publishing: 01/10/2013.