ROLE OF FETAL VIBROACOUSTIC STIMULATION TEST AS AN ADJUNCT TO NON-REASSURING CARDIOTOCOGRAPHY DURING LABOUR IN LOW-RESOURCE SETTINGS

Sanjib Dutta¹, Narayan Jana², Bibek M Rakshit³, Pradip Kumar Saha⁴, Arun Singh⁵

HOW TO CITE THIS ARTICLE:

Sanjib Dutta, Narayan Jana, Bibek M Rakshit, Pradip Kumar Saha, Arun Singh. "Role of fetal vibroacoustic stimulation test as an adjunct to non-reassuring cardiotocography during labour in low-resource settings". *Journal of Evolution of Medical and Dental Sciences* 2013; Vol2, Issue 42, October 21; Page: 8013-8020.

ABSTRACT: This randomized controlled trial (RCT) was undertaken to evaluate the effect of vibroacoustic stimulation (VAS) on the fetal heart rate tracing mode of delivery and perinatal outcomes, when cardiotocographs (CTG) showed non-reassuring features during labour. In this RCT, 60 women beyond 30 weeks' gestation in labour having non reassuring CTG for 20 min were recruited in either study group (VAS group) or control group (non-VAS group). Women in the study group received VAS for 3 sec up to 3 occasions at an interval of 1 to 3 min while the control group received no VAS. There were significant improvement of baseline variability and acceleration in both VAS and control group during second 20 min ($P < 0.001$). Although these improvements were more evident in VAS group, the difference between the two groups were not statistically significant ($P > 0.05$). However, in VAS group, there was 3 min less testing time required to achieve reassuring CTG features (2.6 versus 5.6 min), and a tendency to have less cesarean delivery (60% versus 55.2%; $p > 0.05$). Perinatal outcomes were similar in both groups. There is a trend toward less number of caesarean section in VAS group, probably because of early and more acquisition of reassuring CTG features. Therefore, this study suggests that VAS is potentially beneficial in women with suspicious CTG. Considering the procedure simple, safe and less expensive, it would be prudent to use VAS during CTG monitoring, especially in low resource countries, where facilities for fetal blood sampling is not widely available.

KEY WORDS: fetal vibroacoustic stimulation, vibroacoustic stimulation test in cardiotocography, fetal vibroacoustic stimulation in low resource settings

INTRODUCTION: Since its introduction, Cardiotocography [CTG] is widely used for fetal monitoring before its efficacy has been evaluated by Randomised Controlled Trial [RCT]. Cochrane review¹, mostly from developed countries suggests that intrapartum CTG is associated with high rate of caesarean section without significant reduction of perinatal mortality [PNM]. This is implicated due to a high false positive rate of CTG (abnormal CTG findings, when baby is healthy), which often fails to differentiate the cause of non-reassuring CTG – a healthy fetus in quiet sleep from a sick fetus with substantial compromise. Around 15% CTG are non-reassuring during labour.

Both invasive (e.g. Fetal Blood Sampling [FBS] for pH estimation and lactate estimation) and non-invasive (e.g. Vibro-acoustic stimulation [VAS], fetal ECG, pulse oximetry) are used as an adjunct to confirm fetal status - healthy or in jeopardy. As FBS is an invasive and technically challenging procedure and it has many contraindications, recent attempts have been made to use VAST as an adjunct to assess fetal status in labour. This simple, non-invasive and inexpensive procedure, which produces sound wave to awaken a healthy fetus in sleep cycle and as

a consequence produces visible CTG changes in the form of acceleration and increased variability in healthy fetus. Unfortunately there is little evidence from RCT regarding efficacy of this test², especially in lower resource settings where sophisticated costly techniques are not widely available.

Therefore we planned to assess efficacy of VAST in women who showed non-reassuring CTG during labour in setups, where CTG are often used intermittently but not as a continuous fetal monitoring, because of resource constraints.

MATERIALS AND METHODS: This randomized controlled trial (from Aug 2007 to Aug 2008) was conducted in the department of Obstetrics and Gynecology in collaboration with the department of Neonatology, Institute of Postgraduate Medical Education and Research and S.S.K.M. Hospital, Kolkata. Pregnant women beyond 30 weeks' gestation admitted in labour ward were monitored initially with CTG (Series 50 IP-2, Philips; Boeblingen, Germany) according to clinical indication. Before CTG, she was properly counseled about the trial. If CTG showed non-reassuring features for >20 minutes, an informed consent was taken.

In this study, the CTG trace is defined as non-reassuring if it fulfills one of the following criteria in 20 minutes record: 1) No acceleration, 2) Variability <5 beats per minute (bpm), and 3) Baseline 100-109. (RCOG criteria 2001³, FIGO 1987⁴). Women with following conditions were excluded: Multifetal pregnancies; presence of any fetal malformation; women who require immediate delivery such as abruptio placentae, suspected scar dehiscence or cord prolapse; in cases where there is evidence of normal or pathological CTG; and exposure to narcotic drugs within 4 hours.

A total of 60 women have been randomized into either study group (VAS group) or control group (non-VAS group) according to random number table. Women in the study group, after a quiet period of 20 minutes of CTG showing non-reassuring features (RCOG 2001), received fetal VAS for 3 seconds up to 3 occasions at an interval of 1 to 3 minutes with a commercially available battery operated VAS device [Fetal Acoustic Stimulator, EMCO, Bangalore, India] with 75 db sound intensity at 1.0 meter and frequency of 75 Hz. The control group having similar CTG findings received no VAS. Both the groups were further monitored for at least 20 minutes with CTG tracing, provided there was no clinical indication for immediate intervention.

Primary outcome measures included: changes in baseline, variability>5, number of acceleration or deceleration and method of delivery (rate of normal delivery, operative vaginal delivery and caesarean sections) among two groups. Caesarean sections and operative vaginal delivery due to fetal distress were analyzed. Perinatal outcome which included Apgar score <3 and <7 at both 1 minute and 5 minute, requirement for neonatal resuscitation, need for transfer to NICU, evidence of HIE, multi-organ dysfunction and perinatal mortality were compared between two groups. Need for consumable CTG papers between two groups and its average cost were also studied between two groups.

Nominal data have been expressed, as percentage and comparison between two groups were done by Chi-square test with Yates' correction and Fisher's exact test and $P \leq 0.05$ was considered significant. Continuous variables were expressed as mean +/- standard deviation (SD) and comparisons made by Student t-test.

OBSERVATIONS: This was a RCT which included 60 pregnant women beyond 30 weeks' gestation admitted in the labour ward. After randomization, one woman has left against medical

advice before delivery. As the outcome measures of this pregnancy have remained unknown, the final outcome analysis of the study group contains data for 29 women. Figure 1 shows the CONSORT (consolidated standards of reporting trials) flow diagram of participants through the present randomized controlled trial.

The control and study groups were comparable in the age, parity, education, socioeconomic status, nutrition and antenatal complications.

Majority of the women in both control and study group were included at term i.e. beyond 37 weeks of gestation. Majority (52.5%) of these women had cardiotocography because of reduced fetal movement. Other indications include presence of meconium stained liquor, oligohydramnios, postdated pregnancy and other clinical apprehension regarding fetal status. However, indications of CTG were not significantly different between the control and study group. Of this 59 women included in the study 44(74.58%) women had both reduced variability (<5 bpm) and absent acceleration as criteria for inclusion. Only 9 (15.25%) women had absence of acceleration as an entry criterion. At the entry point the cardiotocographic features between the two groups were similar ($p>0.05$). However, in the study group there were significant improvements of variability compared to controls.

Although caesarean section rate was higher in the control group (60.00% versus 55.17%; $p>0.05$), the overall mode of delivery was not significantly different between the study and control group. Furthermore the duration of hospital stay was similar between control and study groups (9.6 ± 8.7 versus 9.4 ± 7.4 days; $p>0.05$).

Most infants had good Apgar score at 1 min (≥ 7), and didn't require any special resuscitative measures. However, one-fourth of the neonates from both groups required admission at NICU. Average stay of the neonates at NICU was around 7 days more in the control group compared to the study group (11.5 Vs 4.7 days). There are three perinatal deaths - one stillbirth and two early neonatal deaths.

DISCUSSION: Fetal ability to respond to sound or vibroacoustic stimulation has been recognized⁵. Ultrasound studies have suggested that the fetal response to sound stimulation is analogous to the newborn startle reflex and that is a consistent phenomenon from the 28th week of pregnancy⁶. It is possible that unlike healthy fetuses, compromised fetuses might not react to sound stimulation with startle movements. The usefulness of fetal acoustic stimulation in the intrapartum period was considered to be clinical application of a screening technique⁷.

Due to its high sensitivity, this test may serve as a rapid screening test in early labour for detecting high-risk cases for intrapartum fetal distress. The positive predictive value is low, therefore, in the absence of response further assessment is necessary such as fetal scalp blood pH with more demands on time and, personnel and greater cost in performing the test.

Some researchers have raised safety concerns of the stress induced by VAS, such as fetal heart rate deceleration⁸ or potential cochlear damage from increased intrauterine sound levels⁹. Other studies¹⁰ have shown no evidence of auditory nerve or brain stem damage. They have shown that for equal sound pressure, sound intensity and sound vibration are about 4000 times less in amniotic fluid, compared to that produced in air. Therefore it appears that the potential clinical benefit of VAS outweighs any theoretical risk currently associated with the technique.

We have performed a Randomized controlled trial (RCT) in a group of 60 pregnant women

to evaluate efficacy of VAS in non-reassuring fetal heart rate trace in labour. In both control and study group majority of the women were recruited in latent phase of labour and around one-third of women was in active phase of labour. The effects of VAS on the change of fetal heart rate trace, mode of delivery and perinatal outcome have been evaluated. The probable costs saving due to shortened testing time of CTG have also been assessed.

Effect of VAS on CTG features: VAS played a crucial role in improvement of variability (80% in the study group; 66.67% in the control group), a hallmark of fetal well-being. This is concordant with the earlier studies¹¹. However frequencies of acceleration (67.86% in study group versus 62.96% in the control group) are almost similar in two groups. In addition achievement of normal features of CTG tracings has been achieved 3 minutes earlier than the control group [5.6 minutes (SD 4.85) in the control group versus 2.6 minutes (SD 2.47) in the study group]. Similar observations that vibroacoustic stimulation decreases testing time during cardiotocography were found in some earlier studies¹². Furthermore these 3 minutes of saving is likely to save minimum 1 unit of CTG paper (10 min of CTG trace at a paper speed of 1cm/ min, and perforations located at 10 min interval), and that will save approximately Indian rupee 40 to 60 (US\$ 1 approximately) per participant in this study.

Effect of VAS on mode of delivery and maternal outcome: Majority of the women (60% in the control group and 55% in the study group) were delivered by LSCS. Although, there is no significant difference between study and control group in respect to mode of delivery ($P=0.73$), this could be due to small sample size. However, in the present study, there is 5% reduction of LSCS in VAS group – which has an important clinical relevance, although it has not reached statistical significance.

Effect of VAS on perinatal outcome: Majority of the women in both control and study group were included at term i.e. beyond 37 weeks of gestation. The mean birth-weight of the control group was around 150 g less compared to study group (2440g Vs 2597g; $P=0.71$) because relatively more women were preterm (34 weeks to < 37 weeks of gestation) in the control group compared to study group (20.51% vs 6.90%). There is no significant difference in the Apgar score between the two groups. Approximately, one-fourth of the neonates from both groups required admission at NICU. Average stay of the neonates at NICU was more in the control group compared to the study group, which could be attributed to more preterm babies in the control group. The perinatal mortality between the control and study group was not significant ($P>0.05$).

CTG was introduced with the objective of reducing perinatal mortality and morbidity due to perinatal asphyxia. In unselected population its value in reducing perinatal mortality is still not realized because the studies were carried out in the western countries where perinatal mortality is very low (8 per 1000 live birth), and a large sample size (>50'000) will be required to address the issue. Here, VAS as an adjunct to CTG will require a bigger sample size to address the issue related to perinatal mortality and morbidity. As there is limited information available on the perinatal outcomes relating VAS², it is extremely difficult to calculate exact sample size in the context of developing countries. Our present study having a small sample size can't address this issue adequately. However, it was evident that average NICU stay of the babies in the Study

group was reduced by 60% (4.7 Vs 11.5 days). This difference could be attributed to more number of preterm babies in the control groups. Therefore, it would be premature to attribute reduction of NICU stay of the newborn to the use of VAS in the study group. Previous studies² highlighted a number of areas where the randomized controlled trials didn't evaluate important issues such as perinatal outcome following VAS. In our RCT, we found insignificant differences between control and study groups in respect to perinatal outcome. Our observation is concordant with earlier studies¹².

REFERENCES:

1. Thacker SB, Stroup D, Chang M. Continuous fetal heart monitoring for fetal assessment in labour. In Cochrane Library, Issue 1. Oxford: Update Software. 2002.
2. East CE, Smyth R, Leader LR, Henshall NE, Colditz PB, Tan KH. Vibroacoustic stimulation for fetal assessment in labour in the presence of a non-reassuring fetal heart rate trace. Cochrane Database of Systematic Reviews, Issue 3, 2008. CD004664.
3. Royal College of Obstetricians and Gynaecologists: Clinical Effectiveness Support unit. The use of electronic fetal monitoring. The use and interpretation of cardiotocography in intrapartum fetal surveillance. Evidence based clinical guideline number 8. London: RCOG Press; 2001.
4. FIGO Subcommittee on standards in Perinatal Medicine. Guidelines for the use of fetal monitoring. Int J Gynecol Obstet 1987; 25:159-67.
5. Smith CV, Phelan JP, Paul RH, Broussard P. Fetal acoustic stimulation testing: A retrospective experience with the fetal acoustic stimulation test. Am J Obstet Gynecol 1985; 153:567-8.
6. Gelman SR, Wood S, Spellacy WN, Abrams RM. Fetal movements in response to sound stimulation. Am J Obstet Gynecol 1982; 143:484-5.
7. Ingemarsson I, Arulkumaran S, Paul RH et al. Fetal acoustic stimulation in early labour in patients screened with the admission test. Am J Obstet Gynecol 1988; 158:70-4.
8. Ingemarsson I, Arulkumaran S. Reactive fetal heart rate response to VAS in fetuses with low scalp blood pH. Br J Obstet Gynecol 1989; 96:562-5.
9. Tan KH, Smyth R. Fetal vibroacoustic stimulation for facilitation of tests of fetal well-being. Cochrane Database of Systematic Reviews 2001, Issue 1. CD002963.
10. Arulkumaran S, Talbert D, Hsu TS, Chua S, Anandakumar C, Ratnam SS. In- utero sound levels when VAS is applied to the maternal abdomen: an assessment of the possibility of cochlea damage in the fetus. Br J Obstet Gynecol 1992; 99:43-5.
11. Gagnon R, Hunse C, Carmichael L, Fellows F, Patrick J. External vibratory acoustic stimulation near term: fetal heart rate and heart variability responses. Am J Obstet Gynecol 1987; 156:323.
12. Sood AK. Vibroacoustic stimulation and modified fetal biophysical profile in high-risk pregnancy. J Obstet Gynecol India 2007; 57: 37-41.

Figure 1: CONSORT flow diagram of participants through randomized controlled trial

*1 woman excluded as she left against medical advice

	Control group (No VAS) N=30	Study group (VAS) N=29
Age (Mean, SD)	25.3, 5.01	25.79, 4.46
Parity		
Nulliparous	21(70)	24(83)
Primiparous	6(20)	5(17)
Multiparous	3(10)	0(0)
Education		
Nil	5(17)	5(17)
Class V or less	3(10)	1(4)
High school or more	22(73)	23(79)
Socioeconomic		
Poor	15(50)	9(31)
Low middle	14(47)	16(55)
Upper middle	1(3)	4(14)
Antenatal Complications		
PIH	2(7)	4(14)
GDM	3(10)	1(3)
Hypothyroid	0(0)	1(3)
Heart disease	1(3)	1(3)
Others	1(3)	0(0)
Nutrition		
Mean weight (Kg, SD)	61.63, 8.3	63.93, 7.1
Mean Hb (G/dl, SD)	10.55, 1.0	10.83, 1.2

Table 1: Characteristics of the women in the control and the study group

Percentages in the parentheses

ORIGINAL ARTICLE

There were no significant differences between two groups ($P>0.05$)

	Control group (No VAS) N=30	Study group (VAS) N=29
Gestational age		
Mean (wks), SD	37.7; 2.78	37.98; 2.57
30 to 34 wks	4(13)	3(10)
>34 to ≤37 wks	8(26)	2(7)
>37 wks	18(61)	24(83)
Birth-weight (g)		
Mean, SD	2440; 742	2597; 604
Birth weight (g)		
<1500	4(13)	2(7)
1500 to <2500	9(30)	8(28)
2500	17(57)	19(65)
APGAR score		
<7 at 1min	3(10)	3(10)
≥7 at 1 min	27(90)	26(90)
<7 at 5 min	3(10)	2(7)
≥7 at 5 min	27(90)	27(93)
NICU admission	8 (26.67)	7 (25)
Average NICU stay in days (Mean, SD)	11.5, 14.95	4.7, 5.79
Reasons for admission		
HIE	3(10)	3(10)
MAS	1(3)	1(3)
Hyperbilirubinemia	3(10)	2(7)
Others	1(3)	1(3)
Perinatal death	2(7)	1(3)

Table 2: Perinatal outcome of the women in the control and the study group.

Percentages in the parentheses

* HIE= Hypoxic ischemic encephalopathy

*MAS= Meconium aspiration syndrome. These babies also developed stage 1 HIE

Disclosure of interests: None.

Contribution to authorship: SD and NJ conceived the idea. SD, BMR, PKS collected the data and followed up cases. SD wrote the initial draft and made the analysis and interpretation of data, which was modified by all others with critical inputs. AS revised the article critically. All authors read and approved the final version of the draft.

Details of ethics approval: Approved by institutional ethics committee.

Funding: There was no funding for this paper.

Acknowledgements: None.

AUTHORS:

1. Sanjib Dutta
2. Narayan Jana
3. Bibek M. Rakshit
4. Pradip Kumar Saha
5. Arun Singh

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Obstetrics and Gynaecology, College of Medicine and Sagore Dutta Hospital, Kamarhati, Kolkata, West Bengal, India.
2. Professor, Department of Obstetrics and Gynaecology, Burdwan Medical College and Hospital, Burdwan, West Bengal, India.
3. Associate Professor, Department of Obstetrics and Gynaecology, Burdwan Medical College and Hospital, Burdwan, West Bengal, India.

4. Assistant Professor, Department of Obstetrics and Gynaecology, Midnapore Medical College and Hospital, Midnapore, West Bengal, India.
5. Professor, Department of Paediatrics, College of Medicine and Sagore Dutta Hospital, Kamarhati, Kolkata, West Bengal, India.

NAME ADDRESS EMAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Sanjib Dutta,
Assistant Professor,
Department of Obstetrics and Gynaecology,
College of Medicine and Sagore Dutta Hospital,
Kamarhati, 578, B I. Road, Kolkata - 58,
West Bengal, India.
Email- dr_sanjib_2007@yahoo.co.in

Date of Submission: 30/08/2013.
Date of Peer Review: 31/08/2013.
Date of Acceptance: 14/09/2013.
Date of Publishing: 15/10/2013