CORRELATION OF PORTAL VEIN DIAMETER, SPLENOMEGALY AND THROMBOCYTOPAENIA WITH GASTRO-OESOPHAGEAL VARICES IN CIRRHOTIC PATIENTS

Krishan Oberoi¹, Upasana²

¹Associate Professor, Department of Medicine, Guru Nanak Dev Hospital, Amritsar.

²Lecturer, Department of Obstetrics & Gynaecology, Guru Nanak Dev Hospital, Amritsar.

ABSTRACT

BACKGROUND

The word cirrhosis comes from the Greek word *kirrhos*, which means orange yellow.¹ Cirrhosis is a chronic disease of the liver in which diffuse destruction and regeneration of hepatic parenchymal cells and diffuse increase in connective tissue has resulted in disorganisation of the lobular architecture, which leads to various complications such as portal hypertension and its sequelae including gastro-oesophageal varices and splenomegaly, thrombocytopaenia, etc.

The aim of this study is to study the correlation of portal vein diameter, splenomegaly and thrombocytopaenia with gastro-oesophageal varices in cirrhotic patients.

MATERIALS AND METHODS

This is an observational study including 100 patients of cirrhosis attending OPD/ admitted in various wards of Guru Nanak Dev Hospital and allied group of hospitals attached to Government Medical College, Amritsar.

RESULTS

Most of the patients in our study had grade II oesophageal varices (29%) and the most common age distribution in patients with grade II varices is 25 - 75 years. In our study, it is seen that portal vein diameter increases with development of gastro-oesophageal varices (p-value < 0.001). Spleen size increases with formation of gastro-oesophageal varices (p-value < 0.001). Platelet count decreases with the development of gastro-oesophageal varices (p-value < 0.001).

CONCLUSION

It is concluded from our study that there is a positive correlation between portal vein diameter and splenomegaly with gastro-oesophageal varices and an inverse relationship between thrombocytopaenia and gastro-oesophageal varices.

KEYWORDS

Cirrhosis, Splenomegaly, Thrombocytopaenia.

HOW TO CITE THIS ARTICLE: Oberoi K, Upasana. Correlation of portal vein diameter, splenomegaly and thrombocytopaenia with gastro-oesophageal varices in cirrhotic patients. J. Evolution Med. Dent. Sci. 2017;6(83):5806-5810, DOI: 10.14260/jemds/2017/1260

BACKGROUND

The definition of cirrhosis remains morphological, described by World Health Organisation (WHO) in 1978 as: "a diffuse process characterised by fibrosis and the conversion of normal liver architecture into structurally abnormal nodules."²

Cirrhosis is a chronic disease of the liver, in which diffuse destruction and regeneration of hepatic parenchymal cells and diffuse increase in connective tissue has resulted in disorganisation of the lobular architecture. The triad of parenchymal necrosis, regeneration and scarring is always present in cirrhosis regardless of individual clinical manifestations.³

Clinical features of cirrhosis derive from morphological alterations and usually reflect severity of hepatic damage rather than the aetiology of the underlying liver disease.

Fibrosis and distorted vasculature lead to portal hypertension and its sequelae including gastro-oesophageal varices and splenomegaly. Ascites and hepatic encephalopathy result from both portal hypertension and hepatocellular insufficiency. Thrombocytopaenia occurs as a result of hypersplenism and in alcoholic patients due to direct bone marrow suppression by ethanol. Diminished protein synthesis can lead to reduced production of factors I, II, V, VII, IX and X. The main causes of cirrhosis are alcoholic liver disease (ALD), hepatitis B (HBV), hepatitis C (HCV), non-alcoholic steatohepatitis (NASH), haemochromatosis, autoimmune hepatitis (AIH), primary biliary cirrhosis (PBC) and primary sclerosing cholangitis (PSC).^{4,5}

The natural history of cirrhosis can be divided into a preclinical and a subsequent clinical phase. The preclinical phase is usually prolonged over several years; once clinical events occur such as ascites, hepatic encephalopathy, variceal bleeding or the development of hepatocellular carcinoma, the remaining course of the disease is much shorter and usually fatal.⁶

Portal hypertension is the most common and lethal complication of chronic liver disease. Portal hypertension is defined as an increase in portal pressure, in which the pressure gradient between the portal vein and inferior vena cava (the portal pressure gradient) is increased above the upper normal limit of 5 mmHg. The clinical significance increases when portal pressure increases above 10 mmHg

'Financial or Other Competing Interest': None.

Submission 09-09-2017, Peer Review 02-10-2017,

Acceptance 09-10-2017, Published 16-10-2017.

Corresponding Author:

Dr. Krishan Oberoi,

#13-C, Mata Kaulan Marg,

Kashmir Avenue,

Amritsar-143001, Punjab.

E-mail: drkrishanoberoi@gmail.com

DOI: 10.14260/jemds/2017/1260

(formation of varices) and above 12 mmHg (variceal bleeding and ascites).^{7,8,9}

Development of oesophageal varices is one of the major complications of portal hypertension. After varices have developed, about one-third of patients die of gastro-oesophageal variceal bleeding. The risk of initial bleeding from varices is 25% - 35% within two years with first bleeding episodes mostly occur within one year after detection of varices.^{10,11,12} The reported mortality from first episode of variceal bleeding ranges from 40% - 70%.¹³

Cirrhosis is associated with both quantitative and qualitative abnormalities of platelets.¹⁴ Approximately, 40% of cirrhotic patients have abnormal prolongation of the bleeding time to values of more than 10 minutes and platelet counts less than 100,000/mm³. Causes of thrombocytopaenia in cirrhosis may be: - 1) Portal hypertension-induced splenic sequestration; 2) Alterations in thrombopoietin; 3) Bone marrow suppression mediated by toxins (e.g. alcohol, hepatitis B and C); 4) Consumptive coagulopathy; 5) Increased blood loss (e.g. haemorrhage).

Recently, the Baveno III consensus conference on portal hypertension recommended that all cirrhotic patients should be screened for the presence of oesophageal varices when liver cirrhosis is diagnosed.¹⁵

Upper gastrointestinal endoscopy is the most commonly used method to detect varices. Endoscopic grading of oesophageal varices is subjective. Various criteria have been used to standardise the reporting of oesophageal varices as those compiled by the Japanese Research Society for Portal Hypertension, which include red colour signs, colour of the varix, form (size) of the varix and location of the varix.^{16,17}

Although, varices may form in any location along the tubular gastrointestinal tract, they most often appear in the distal few centimetres of the oesophagus.^{9,18}

Gastric varices were graded according to classification proposed by Sarin et al¹⁹ for gastro-oesophageal varices (GOV) and isolated gastric varices (IGV).

The current consensus is that all patients with cirrhosis of the liver should be screened for oesophageal varices by endoscopy. In patients in whom no varices are detected on initial endoscopy, endoscopy should be repeated in 2 to 3 years to look for varices. If small varices are detected on the initial endoscopy, endoscopy should be repeated in 1 to 2 years.

Thrombocytopaenia and splenomegaly are independent predictors of large oesophageal varices in Cirrhosis. There is evidence that with increase in portal vein diameter, splenomegaly and thrombocytopaenia there is increase in the chance of formation of gastro-oesophageal varices. Thus, these can be used as non-invasive predictors of presence of oesophageal varices and guide for selecting patients for endoscopic evaluation.

Sarwar et al²⁰ reported that patients with portal vein diameter more than 11 mm are more likely to have oesophageal varices. Another study by Dib et al²¹ showed that oesophageal varices developed when the portal vein diameter exceeded 13 mm.

Sarangapani A et al²² in a study analysed 106 patients with chronic liver disease for prediction of large oesophageal varices. Results showed that incidence of large varices was seen in 41% cases. Sotoudehmanesh R et al²³ in a case control study discussed the role of Endoscopic Ultrasonography in

patients with Chronic Liver disease (CLD). The study concluded that splenic and portal vein dilation and thrombocytopaenia significantly correlated with variceal bleeding ($p < 0.05$).

Elliot Schwarzenberger et al²⁴ conducted a retrospective analysis of 137 patients with cirrhosis over the age of 18 that underwent screening endoscopy for varices. They concluded that the platelet count/ spleen diameter ratio with a cut-off value of 909 may not be sufficiently accurate in predicting the presence of oesophageal varices. Upper GI endoscopy remains the method of choice to screen for the presence of varices.

Aims and Objectives

To find out correlation of portal vein diameter, splenomegaly and thrombocytopaenia with gastro-oesophageal varices in cirrhotic patients.

MATERIALS AND METHODS

This is an observational study including 100 patients of cirrhosis attending OPD/ admitted in various wards of Guru Nanak Dev Hospital and allied group of hospitals attached to Government Medical College, Amritsar. After obtaining informed consent, the patients were subjected to detailed history and clinical examination.

Each patient was subjected to investigations which included platelet count, ultrasound to detect portal vein diameter and spleen size and oesophago-gastro-duodenoscopy to grade gastric and oesophageal varices. The results of this study were systematically collected and statistically analysed.

According to Paquet¹⁸ Grading System Varices were graded as-

Grade 0- No varices.

Grade I- Varices, disappearing with insufflation.

Grade II- Larger, clearly visible, usually straight varices, not disappearing with insufflation.

Grade III- More prominent varices, locally coil-shaped and partly occupying the lumen.

Grade IV- Tortuous, sometimes grape-like varices occupying the oesophageal lumen.

Gastric varices were graded according to classification proposed by Sarin et al¹⁹ for gastro-oesophageal varices (GOV) and isolated gastric varices (IGV).

Inclusion Criteria

- Individuals of age > 18 years.
- Patients of cirrhosis with portal hypertension.

Exclusion Criteria

- Patients with advanced cirrhosis (Child-Pugh Class C).
- Patients with human immunodeficiency virus (HIV) infection.
- Patients with hepatocellular carcinoma.
- Patients with portal vein thrombosis.
- Patients with current alcohol abuse.
- Previous or current treatment with B-blockers, diuretics and other vasoactive drugs.
- Patients with severe or unstable cardiovascular disease and pulmonary disease.
- Patients with clinically significant renal or hepatic disease or dysfunction,
- Patients with haematological disorders.

Upper gastrointestinal endoscopy was done with fibre-optic video endoscope manufactured by Pentax. The results were systematically recorded and analysed. The ANOVA with post-hoc test has been used for statistical analysis. The SPSS software has been used in performance of statistical analysis. After evaluating patients for above parameters, treatment for cirrhosis was initiated according to the need of the patient.

RESULTS

A total of 100 patients were selected for the study. Among them, 74 patients (74%) were males and 26 patients (26%) were females. Most of the patients (29%) had grade II oesophageal varices, 27% had grade III varices, 25% had grade IV varices and 19% had grade I varices. The distribution of patients according to grades of oesophageal varices is shown in Table I.

Grade of Oesophageal Varices	Number of Patients	Percentage (%)
I	19	19
II	29	29
III	27	27
IV	25	25
Total	100	100

Table I. Distribution of Grades of Oesophageal Varices

Mean age of the patients with grade I oesophageal varices was 47.58 ± 11.442 years, mean age of the patients with grade II varices was 46.45 ± 11.44 years, mean age of the

Grade of Oesophageal Varices	Number of Patients	Range (In mm)	Mean Portal Vein Diameter (mm) Mean $\pm$ SD	Correlation Coefficient (r)	Significance (p value)
I	19	12.0-14.0	13.16 $\pm$ 0.61	0.953	p < 0.001(HS)
II	29	13.6-16.0	14.59 $\pm$ 0.56		
III	27	15.0-17.5	16.39 $\pm$ 0.69		
IV	25	17.0-21.0	19.02 $\pm$ 0.99		

Table III. Correlation of Mean Portal Vein Diameter with Grading of Oesophageal Varices

Correlation of Spleen Size with the Grading of Oesophageal Varices

It was observed from the study that the mean spleen size of the patients with grade I oesophageal varices was 12.05 ± 0.67 cm with grade II oesophageal varices was 14.33 ± 0.93 cm with grade III oesophageal varices was 16.35 ± 1.03 cm and with grade IV oesophageal varices was 19.10 ± 1.16 cm.

From statistical analysis, it was also found that there was a positive correlation between spleen size and grading of the oesophageal varices and this correlation was found to be statistically significant (p < 0.001) that concluded that when spleen size increased, oesophageal varices also increased in size as shown in Table IV.

Grade of Oesophageal Varices	Number of Patients	Range (in cm)	Mean Spleen Size (cm) Mean $\pm$ SD	Correlation Coefficient (r)	Significance (p value)
I	19	11.0-13.0	12.05 $\pm$ 0.67	0.941	p < 0.001(S)
II	29	12.0-16.0	14.33 $\pm$ 0.93		
III	27	14.5-20.0	16.35 $\pm$ 1.03		
IV	25	17.0-21.0	19.10 $\pm$ 1.16		

Table IV. Correlation of Mean Spleen Size with Grading of Oesophageal Varices

patients with grade III varices was 47.56 ± 13.12 years and mean age of the patients with grade IV varices was 48.72 ± 15.13 years ranging from 14 to 85 years as shown in Table II.

Grade of Oesophageal Varices	Number of Patients	Range (Years)	Mean Age (Years) (mean $\pm$ SD)	Significance (p value)
I	19	30-65	47.58 $\pm$ 11.44	p = 0.937 (NS)
II	29	25-75	46.45 $\pm$ 11.44	
III	27	25-82	47.56 $\pm$ 13.12	
IV	25	14-85	48.72 $\pm$ 15.13	

Table II. Age Distribution with the Grading of Oesophageal Varices

Correlation of Portal Vein Diameter with the Grading of Oesophageal Varices

It was observed from the study that the mean portal vein diameter of the patients with grade I oesophageal varices was 13.16 ± 0.61 mm with grade II oesophageal varices was 14.59 ± 0.56 mm with grade III oesophageal varices was 16.39 ± 0.69 mm and with grade IV oesophageal varices was 19.02 ± 0.99 mm.

From statistical analysis, it was also found that there was a positive correlation between portal vein diameter and grading of the oesophageal varices and this correlation was found to be statistically significant (p < 0.001) that meant that when portal vein diameter increased oesophageal varices also increased in size as shown in Table III.

Correlation of Thrombocytopaenia with the Grading of Oesophageal Varices

From the above study, it was observed that mean platelet count in grade I oesophageal varices was $142568.42 \pm 6640.369/\text{mm}^3$, in grade II oesophageal varices was $109272.41 \pm 9567.16/\text{mm}^3$, in grade III oesophageal varices was $79288.89 \pm 5506.94/\text{mm}^3$ and in grade IV oesophageal varices was $45156.00 \pm 7032.43/\text{mm}^3$.

Statistical analysis showed an inverse relation between the platelet count and grading of oesophageal varices. It was a statistically significant correlation with p-value < 0.001. It also suggested that as the platelet count decreased, grading of oesophageal varices increased as shown in Table V.

Grade of Oesophageal Varices	Number of Patients	Range (in/mm ³)	Mean Platelet Count (/mm ³) Mean $\pm$ SD	Correlation Coefficient (r)	Significance (p value)
I	19	120000-152000	142568.42 $\pm$ 6640.37	-0.961	P < 0.001(S)
II	29	80000-129000	109272.41 $\pm$ 9567.16		
III	27	70000-90000	79288.89 $\pm$ 5506.94		
IV	25	36000-58000	45156.00 $\pm$ 7032.43		

Table V. Correlation of Mean Platelet Count with Grades of Oesophageal Varices

It was observed that concomitant gastric varices along with oesophageal varices were present only in 2 (2%) patients in our study.

Number	Age (Years)	Gender	Platelet Count (/mm ³)	Spleen size (cm)	Portal Vein Diameter (mm)	Oesophageal Varices (grade)	Gastric Varices (grade)
1	14	Female	41,000	19	19	IV	GOV II
1	45	Male	43,000	21	20	IV	GOV II

Table VI. Table showing Distribution of Concomitant Gastric Varices and Oesophageal Varices

There was no patient of isolated gastric varices in our study.

DISCUSSION

Cirrhosis is the most advanced form of liver disease and variceal haemorrhage is one of its lethal complications. Over half of the patients with cirrhosis will develop varices. The risk of recurrent bleeding is approximately 66% and usually occurs within 6 months of the initial bleeding episode.²⁵

Cirrhotic patients with large oesophageal varices are at a high risk for bleeding, so efforts should be made to identify cirrhotic patients with large varices.²⁵ In 1997, the American College of Gastroenterology (ACG) recommended screening endoscopy for cases with established cirrhosis who were candidates for medical therapy.²⁶

One hundred patients were enrolled for this study and out of these 74 patients (74%) were males and 26 patients (26%) were females. Mean age of the patients was 47.5 years ranging from 14 to 85 years. It was observed that out of 100 patients in 19 patients (19%) grade I oesophageal varices were seen, in 29 patients (29%) grade II oesophageal varices were seen, in 27 patients (27%) grade III oesophageal varices were seen, and in 25 patients (25%) grade IV oesophageal varices were seen. Also, out of 100 patients, in 2 patients concomitant gastric varices were seen. A similar study was done by Abbasi et al²⁷ in 102 patients to find out the correlation of thrombocytopaenia with grading of oesophageal varices in chronic liver disease patients. Out of 102 patients 7 patients had oesophageal varices grade I; 24 had grade II; 35 had grade III and 36 had grade IV. Gastric varices were detected only in 2 patients.

Our study demonstrated that thrombocytopaenia, presence of portal hypertension with splenomegaly and portal vein diameter are strong predictors of developing gastro-oesophageal varices in cirrhotic patients.

Earlier, the pathophysiology of thrombocytopaenia in liver disease was linked to the presence of hypersplenism, but now it has been discovered that thrombopoietin production is dependent on functioning liver cell mass and is responsible for reduced thrombopoiesis and consequently peripheral thrombocytopaenia. On the other hand, the presence of splenomegaly in cirrhotic patients is likely the result of vascular disturbance that are mainly linked to portal hypertension.²⁷

In our study, it was observed that mean platelet count in grade I oesophageal varices was $142568.42 \pm 6640.369/\text{mm}^3$, in grade II oesophageal varices was $109272.41 \pm 9567.16/\text{mm}^3$, in grade III oesophageal varices was $79288.89 \pm 5506.94/\text{mm}^3$ and in grade IV oesophageal varices was $45156.00 \pm 7032.43/\text{mm}^3$. Platelet count showed a highly statistical significant inverse correlation with the grading of oesophageal varices, which is in agreement with Thomopoulos et al²⁴ who reported that platelet count was the only common factor found to be significant predictor of both small and large varices.

Splenomegaly and portal vein diameter are also reliable predictors of the presence of oesophageal varices. In our study, the mean spleen size of the patients with grade I oesophageal varices was 12.05 ± 0.67 cm with grade II oesophageal varices was 14.33 ± 0.93 cm with grade III oesophageal varices was 16.35 ± 1.03 cm and with grade IV oesophageal varices was 19.10 ± 1.16 cm, which showed a statistical significant direct correlation with the presence of oesophageal varices which is in agreement with Chalasani et al²⁸ who reported that splenomegaly is recognised as one of the diagnostic signs of cirrhosis and portal hypertension. In a study carried out by Mandal et al,²⁹ it has been shown that grading of oesophageal varices also correlates with spleen size and portal vein diameter. Sharma and Agarwal³⁰ in their study had noted that clinically palpable spleen was associated with high-grade varices; however, they did not measure the splenic size radiologically. Farooqi et al³¹ also found out that platelet count of $< 65 \times 10^3/\mu\text{L}$, serum albumin < 2.2 g/dL and portal vein diameter of > 13 mm are independent and significant predictors of oesophageal varices on endoscopy.

It was observed in our study that the mean portal vein diameter of the patients with grade I oesophageal varices was 13.16 ± 0.61 mm, with grade II oesophageal varices was 14.59 ± 0.56 mm, with grade III oesophageal varices was 16.39 ± 0.69 mm and with grade IV oesophageal varices was 19.02 ± 0.99 mm, which shows a statistical significant direct correlation between Portal vein diameter and the presence of oesophageal varices which is in accordance with the study done by Sarwar et al²⁰ who postulated that portal vein diameter more than 11 mm on ultrasonography is independently associated with the presence of oesophageal varices.

Hence, it is concluded from our study that gastro-oesophageal varices in cirrhotic patients has a direct correlation with portal vein diameter and splenomegaly and an inverse correlation with thrombocytopaenia.

Hence, the inference is drawn that with increase in size of gastro-oesophageal varices, the size of portal vein and spleen also increases and the patient is likely to develop thrombocytopaenia.

CONCLUSION

It is concluded from our study that,

1. Portal vein diameter increases with development of gastro-oesophageal varices.
2. Spleen size increases with formation of gastro-oesophageal varices.
3. Platelet count decreases with the development of gastro-oesophageal varices.
4. There is a positive correlation between portal vein diameter and splenomegaly with gastro-oesophageal varices and an inverse relationship between thrombocytopaenia and gastro-oesophageal varices.

REFERENCES

- [1] Arey LB, Burrows W, Greenhill JP, et al. Dorland's illustrated medical dictionary. 23rd edn. Philadelphia: WB Saunders Company 1962: p. 286.
- [2] Anthony PP, Ishak KG, Nayak NC, et al. The morphology of cirrhosis. Recommendations on definition, nomenclature, and classification by a working group sponsored by the World Health Organization. *J Clin Pathol* 1978;31(5):395-414.
- [3] Conn HO. Cirrhosis. In: Schiff L. edr. Diseases of the liver. 4th edn. Philadelphia: JB Lippincott Company 1975: p. 833.
- [4] Guha NI, Iredale JP. Clinical and diagnostic aspects of cirrhosis. In: Rodés J, Benhamou JP, Blei A, et al. eds. Textbook of hepatology from basic science to clinical practice. 3rd edn. Oxford: Blackwell Publishing 2007:604-22.
- [5] Podolsky DK, Isselbacher KJ. Cirrhosis and alcoholic liver disease and major complications of cirrhosis. In: Fauci AS, Braunwald E, Isselbacher KJ, et al. eds. Harrison's Principles of internal medicine. 14th edn. McGraw-Hill 1998:1704-17.
- [6] De Franchis R, Primignani M. Natural history of portal hypertension in patients with cirrhosis. *Clin Liver Disease* 2001;5(3):645-63.
- [7] Casado M, Bosch J, Garcia-Pagan JC, et al. Clinical events after transjugular intrahepatic portosystemic shunt: correlation with hemodynamic findings. *Gastroenterology* 1998;114(6):1296-1303.
- [8] Gacia-Tsao G, Groszmann RJ, Fisher RL, et al. Portal pressure, presence of gastroesophageal varices and variceal bleeding. *Hepatology* 1985;5(3):419-24.
- [9] Rigau J, Bosch J, Bordas JM, et al. Endoscopic measurement of variceal pressure in cirrhosis correlation with portal pressure and variceal hemorrhage. *Gastroenterology* 1989;96(3):873-80.
- [10] Gupta TK, Tournier M, Chung MK, et al. Endothelial dysfunction and decreased production of nitric oxide in the intrahepatic microcirculation of cirrhotic rats. *Hepatology* 1998;28(4):926-31.
- [11] De Franchis R, Dell'Era A. Non-invasive diagnosis of cirrhosis and the natural history of its complications. *Best Pract Res Clin Gastroenterol* 2007;21(1):3-18.
- [12] Rigo GP, Merighi A, Chahin NJ, et al. A prospective study of the ability of three endoscopic classifications to predict hemorrhage from esophageal varices. *Gastrointest Endosc* 1992;38(4):425-9.
- [13] Shabestari AA, Nikoukar E, Bakhshandeh H. Hepatic doppler ultrasound in assessment of the severity of esophageal varices in cirrhotic patients. *Iran J Radiol Spring* 2007;4(3):151-8.
- [14] Kajiwaru E, Akagi K, Azuma K, et al. Evidence for an immunological pathogenesis of thrombocytopenia in chronic liver disease. *Am J Gastroenterol* 1995;90(6): 962-6.
- [15] Bosch J, Navasa M, Garcia-Pagan JC. Portal hypertension. *Med Clin North Am* 1989;73:931-53.
- [16] González-Ojeda A, Cervantes-Guevara G, Chávez-Sánchez M, et al. Platelet count/spleen diameter ratio to predict esophageal varices in Mexican patients with hepatic cirrhosis. *WJG* 2014;20(8):2079-84.
- [17] Shah VH, Kamath PS. Portal hypertension and gastrointestinal bleeding. In: Sleisenger and Fordtran's gastrointestinal and liver disease. Vol 2. 9th edn. Elsevier 2009:1496-7.
- [18] Paquet KJ. Prophylactic endoscopic sclerosing treatment of esophageal wall in varices: a prospective controlled trial. *Endoscopy* 1982;14(1):4-5.
- [19] Sarin SK, Preimignani M, Agarwal SR. Gastric varices. In: Defranchis R. edr. Portal hypertension, proceeding of the third Baveno International Consensus Workshop on definitions, methodology and therapeutic strategies. Oxford: Blackwell Science 2001:576-9.
- [20] Sarwar S, Khan AA, Alam A. Non-endoscopic prediction of presence of esophageal varices in cirrhosis. *J Coll Physicians Surg Pak* 2005;15(9):528-31.
- [21] Dib N, Konate A, Obesti F, et al. Non-invasive diagnosis of portal hypertension in cirrhosis. Application to the primary prevention of varices. *Gastroenterol Clin Biol* 2005;29(10):975-87.
- [22] Sarangapani A, Shanmugam C, Kalyanasundaram M, et al. Noninvasive prediction of large esophageal varices in chronic liver disease patients. *Saudi J Gastroenterol* 2010;16(1):38-42.
- [23] Sotoudehmanesh R, Ainechi S, AliAsgari A, et al. Endoscopic ultrasonography in patients with chronic liver disease: a case control study. *Govaresh* 2012;16(4):270-4.
- [24] Schwarzenberger E, Meyer T, Golla V, et al. Utilization of platelet count /spleen diameter ratio in predicting the presence of esophageal varices in patients with cirrhosis. *J Clin Gastroenterol* 2010;44(2):146-50.
- [25] Zaman A, Becker T, Lapidus J, et al. Risk factors for the presence of varices in cirrhotic patients without a history of variceal hemorrhage. *Archives of Internal Medicine* 2001;161(21):2564-70.
- [26] The North Italian Endoscopic Club for the Study and Treatment of Esophageal Varices. Prediction of the first variceal hemorrhage in patients with cirrhosis of the liver and esophageal varices: a prospective multicenter study. *N Engl J Med* 1988;319(15):983-9.
- [27] Abbasi A, Butt N, Bhutto AR, et al. Correlation of thrombocytopenia with grading of esophageal varices in chronic liver disease patients. *Journal of the College of Physicians and Surgeons Pakistan* 2010;20(6):369-72.
- [28] Chalasani N, Imperiale TF, Ismail A, et al. Predictors of large esophageal varices in patients with cirrhosis. *Am J Gastroenterol* 1999;94(11):3285-91.
- [29] Mandal L, Mandal SK, Bandyopadhyay D, et al. Correlation of portal vein diameter and splenic size with gastro-oesophageal varices in cirrhosis of liver. *JIACM* 2011;12(4):266-70.
- [30] Sharma SK, Aggarwal R. Prediction of large esophageal varices in patients with cirrhosis of the liver using clinical, laboratory and imaging parameters. *J Gastroenterol Hepatol* 2007;22(11):1909-15.
- [31] Farooqi JI, Ahmed H, Ikramullah Q, et al. Predictors of esophageal varices in patients of liver cirrhosis. *JPMI* 2007;21(01):60-4.