STRESS DISTRIBUTION IN THE NASOMAXILLARY COMPLEX FOLLOWING APPLICATION OF ORTHOPAEDIC HEADGEAR FORCES- A 3-DIMENSIONAL FEM STUDY

Suruchi Seth¹, Shivalinga B. M², Sunil Seth³, Gurinderjit Singh Nagi⁴

¹Associate Professor, Department of Dentistry, PIMS, Jalandhar.

²Former Professor and HOD, Department of Orthodontics, JSS Dental College and Hospital, Mysore.

³Assistant Professor, Department of Radiodiagnosis, PIMS, Jalandhar.

⁴Associate Professor, Department of Surgery, PIMS, Jalandhar.

ABSTRACT

BACKGROUND

Extraoral force has been in use in orthodontics since Cellier introduced it in early 1800s. Like many other treatment modalities, interest in headgears and the understanding of its probable effect has varied over the years. From only dental changes to skeletal changes affecting the cranial base structures as well, all have been attributed to headgear use at various times. At present, it is well established that headgears have a retarding effect on the growth and development of maxilla. However, the exact biomechanical effect of different force directions on the sutural response is still under consideration.

MATERIALS AND METHODS

The present study was undertaken to examine the effect of posteriorly directed headgear forces on the various nasomaxillary sutures. FEM model was created having 236685 elements and 56052 nodes and 1 Kgf force was applied at -30°, 0°, 30°, 52.4° and 60° to the functional occlusal plane.

RESULTS

1. In loading with forces in 52.4° and 60° superior directions, compressive stresses were similarly generated in most anatomic areas and both the normal and shear stresses reduced and exhibited a convergence to a certain level.
2. As the force direction approached that of the CRe, mean principal stresses approached a uniform level of compressive stress.
3. Maximum strain or displacement of all the sutures was observed at a force direction of -30°. Strain approached to a minimum level at a force direction of 30° and 52.4°.

CONCLUSION

These findings show that the stress distribution in the sutures varies in relation to the direction of force. Force applied in the direction closest to that of the CRe may produce the most effective sutural modification for controlling maxillary growth. This is an observational study.

KEYWORDS

FEM, Nasomaxillary Complex, Sutures, CRe, Stress.

HOW TO CITE THIS ARTICLE: Seth S, Shivalinga BM, Seth S, et al. Stress distribution in the nasomaxillary complex following application of orthopaedic headgear forces- a 3-dimensional fem study. J. Evolution Med. Dent. Sci. 2017;6(91):6510-6514, DOI: 10.14260/jemds/2017/1413

BACKGROUND

The use of extraoral force began in the early 1800s when Cellier developed cervical and occipital traction to prevent luxation of mandible.¹ However, the widespread acceptance of orthopaedic therapy was brought about by Silas Kloehn's interesting study in 1947, where he showed headgear therapy to be an effective treatment for patients with maxillary excess.²

Since then, various types of headgears have been developed for clinical use in orthodontics.³ The response of the craniofacial structures to extraoral appliances continues to be a major concern for orthodontists. An understanding of orthopaedic effects of extraoral force depends on achieving

an understanding of the biological response of the maxillofacial sutures to clearly defined force systems.⁴ Animal studies have shown that headgear use results in compression or stretching of maxillary sutures resulting in remodelling changes of large magnitude and the force applied was also transmitted to adjacent bones with increased sutural activity.⁵

However, the interpretation of results in treatment proves extremely complicated. Variations in growth pattern of the craniofacial complex, timing of maturation, variation in tooth movement and variation in magnitude, direction, point of attachment of applied forces form a myriad of inter-related factors which greatly hinder precise analysis of treatment effects.

Due to these drawbacks, various animal studies^{4,6,7,8,9,10} and cephalometric studies^{1,11,12,13,14,15,16,17,18,19,20,21,22} were carried out. Though these studies gave a lot of useful information, they could not quantify the stress and strain across the sutures. This led to the use of models to access the levels of stress and strain across the sutures. Strain gauges⁵, holography,^{23,24,25,26} laser metrology¹¹ and laser speckle interferometry²⁷ are all highly sensitive techniques, able to measure extremely small initial deformations of bone on macerated skull. Unfortunately, most of these techniques are

'Financial or Other Competing Interest': None.

Submission 16-09-2017, Peer Review 11-11-2017,

Acceptance 17-11-2017, Published 27-11-2017.

Corresponding Author:

Dr. Suruchi Seth,

Department of Dentistry,

Punjab Institute of Medical Sciences,

Jalandhar.

E-mail: suruchikalra@yahoo.com

DOI: 10.14260/jemds/2017/1413

limited in their ability to measure internal displacements and stress-strain levels in a complicated structure such as the craniofacial skeleton.²⁸

To overcome the drawbacks of the aforementioned methods, FEA study was taken up to study the stress distribution in various sutures of the nasomaxillary complex on application of various headgear forces.

Aims and Objectives

The objectives of the Present Study are

- To study the stress distribution in the various nasomaxillary sutures during application of headgear forces.
- To study the displacement of various nasomaxillary sutures during application of headgear forces.
- To compare stress distribution and displacement of nasomaxillary sutures on application of headgear forces in various directions.

MATERIALS AND METHODS

A three-dimensional mathematical finite element model was generated for analysis from spiral CT scan images of 3 mm sections of an intact skull, with full complement of teeth. The sections obtained in DICOM format and the data were fed into the computer. Using the MIMICS software, these cross sections were converted into a three-dimensional mathematical model. Thus, a virtual geometric model of the skull was obtained. The virtual geometric model is divided into several finite elements. The mesh chosen for the study was HyperMesh 7.0, using a four-noded tetrahedral element. ANSYS software was used to create the finite element model. The finite element model thus built comprised of 236685 elements and 56052 nodes. All the nasomaxillary sutures like Frontomaxillary, Sphenomaxillary, Frontozygomatic, Sphenozygomatic, Temporozygomatic and Lamina Cribrosa were generated in the finite element model.

Mechanical properties such as Young's modulus and Poisson's ratio of bone, tooth and sutures were fed to the finite element model.^{28,29,30,31} All materials were assumed to be isotropic and to follow elastic behaviour. Because no direct data were available for the material properties of sutures, two values of stiffness were introduced. Firstly, the modulus of elasticity of suture was considered as half of the stiffness of cortical bone and secondly, stiffness was reduced to a minimum of $E = 1 \text{ N/mm}^2$.²⁸ Due to the initial character of force application and because no changes in time under force load of the apparent Young's modulus occurred, it was not necessary to take viscoelasticity into consideration.

Material	Modulus of Elasticity	Poisson's Ratio
Bone	12,200 MPa	0.3
Tooth	20,000 MPa	0.3
Sutures	6,850 MPa	0.3

Table 1. Mechanical Properties of Various Materials

The study is observational in nature.

The model was constrained at the Foramen Magnum to prevent sliding movement of the model, simulating clinical application of orthopaedic extraoral forces to patients. The generated model was then strained to a load of 1.0 Kgf applied at the first molar region in the 5 prescribed directions: -30°, parallel, 30° superior, 52.4° superior and 60°

superior to the occlusal plane to simulate the various clinical situations. A 52.4° superior direction was demonstrated in advance to connect the centre of resistance of the nasomaxillary complex and the first molar.³² The load was applied and the mean principal stress, shear stress and strain were studied using the ANSYS solver.

Following Statistical Methods were applied in the Present Study-

1. One-Way ANOVA
2. Scheffe's Post Hoc test

All the statistical operations were done through SPSS for Windows (Version 15 evaluation Version, 2006), SPSS Inc. New York.

RESULTS

Graph 1. Mean Principal Stresses in the Anatomic Areas Resisting Upward Displacement of the Complex

Graph 2. Mean Principal Stresses in the Anatomic Areas Resisting Backward Displacement of the Complex

Graph 3. Total Shear Stresses in the Anatomic Areas Resisting Upward Displacement of the Complex

Graph 4. Total Shear Stresses in the Anatomic Areas Resisting Backward Displacement of the Complex

Graph 5. Strain in the Anatomic areas Resisting Upward Displacement of the Complex

Graph 6. Strain in the Anatomic Areas Resisting Backward Displacement of the Complex

Graph 7. Association between Changes in Mean Principal Stresses at Five Different Points and Force Directions to the Functional Occlusal Plane Determined by the Maxillary Dentition

DISCUSSION

FEM is a powerful contemporary research tool that has been used to study the stress distribution of non-living as well as natural and restored craniofacial structures affected by three-dimensional stress fields, which are difficult to assess otherwise.

The three-dimensional FEM used in the present study provides the freedom to simulate orthodontic force systems applied clinically and allows analysis of the response of the craniofacial skeleton to the orthodontic loads in three-dimensional space.

The directions of loading chosen in the present study were taken so as to simulate clinical conditions. 30° inferior to the occlusal plane represents cervical traction; 0° represents a straight pull headgear; 30° superior to the occlusal plane represents an occipital pull headgear and 60° superior to the occlusal plane represents the high pull or parietal pull headgear. The orthopaedic loading of 52.4° to the occlusal plane was based on previous studies.

The posterior and anterior parts of the frontozygomatic suture showed almost uniform tensile stresses, which turned to compressive, when the direction of loading changed to 52.4° and 60°. However, the middle part of the suture showed tensile stresses which were much higher than observed in the anterior and posterior regions.

The lamina cribrosa showed almost uniformly compressive forces which decreased in magnitude as the force direction became more vertical. Throughout the loading range, the posterior part of the suture showed much higher stresses than the anterior.

The posterior part of the frontomaxillary suture showed tensile stresses, which decreased as the direction of pull became more upwards and showed slight compressive stress when the direction of pull changed to 60°. The anterior part of the suture showed compressive stresses, which again decreased in magnitude as the direction of pull became more vertical.

The superior part of the sphenozygomatic suture showed tensile stresses, which became compressive at 52.4° and 60°. However, the middle and inferior parts of the suture were uniformly compressive, irrespective of the force direction, showing the least stresses at the direction of 52.4°.

In the inferior regions of the temporozygomatic suture, slight tensile stresses were induced in loading with forces applied in different directions, whereas stresses in the superior region were mainly compressive.

Stresses in the superior, middle and inferior regions of the sphenozygomatic suture were mainly compressive in nature, with a substantial decrease in magnitude, in loading with more superiorly directed forces.

The total amount of shear stress at seven anatomic points resisting upward displacement of the nasomaxillary complex for each of five directions was calculated. When a downward force of 30° degrees was applied, total shear stress was highest, indicating a substantial shear deformation of the sutural structure. These stresses decreased gradually when the direction was varied from 30° inferior to 52.4° or 60° superior. This indicates that the loading directions of 52.4° and 60° produce least shear deformation in the anatomic areas resisting superior displacement of the complex.

The total amount of shear stresses at eight anatomic points in the three sutures resisting backward displacement of the nasomaxillary complex for each of five directions was calculated. In loading with a posteriorly directed horizontal force, total shear stress was greatest, indicating the sliding of two adjacent bones and the subsequent substantial shear deformation of the sutural structure associated with counter clockwise rotation of the complex.

Strain or displacement for all the six sutures was calculated for all the five loading conditions. Maximum strain was seen when load was applied at 30° inferior to the occlusal plane. Minimum strain was seen at 30° superior to the occlusal plane.

At headgear loading directions of 52.4° and 60°, the sphenozygomatic suture showed negative strain in its superior and middle regions and high positive strain in its inferior regions. This indicates a bending of the sphenozygomatic suture, with the inferior regions showing a widening of the suture, compared to compression in the superior and middle regions of the suture.

The stresses exhibited a gradual increase or decrease approaching a uniform level of compressive stress as the direction was varied from -30° to 60° to the functional occlusal plane. Thus, variation in these stresses decreased and the nature of stresses became more uniform when directing the line of force closer to the CRe of the nasomaxillary complex.

CONCLUSION

Stress distribution in the nasomaxillary complex was investigated by means of 3-Dimensional Finite Element Analysis. A finite element model of the craniofacial complex was created, that consisted of 56052 nodes and 236685 elements. An extraoral force of 1.0 Kgf was applied to the maxillary first molar in a posterior direction, 30° inferior, parallel and 30°, 52.4° and 60° superior to the occlusal plane. Mean Principal stresses, Shear stresses and Strain were evaluated at six anatomic areas resisting posterior and superior displacement of the complex. The following results were obtained:

1. When 30° inferior, parallel and 30° superior forces were applied, considerable variation in normal stresses at the sutural interfaces was observed in association with substantial shear stresses.
2. In loading with forces in 52.4° and 60° superior directions, compressive stresses were similarly generated in most anatomic areas and both the normal and shear stresses reduced and exhibited a convergence to a certain level.
3. As the force direction approached that of the CRe, mean principal stresses approached a uniform level of compressive stress.
4. Maximum strain or displacement of all the sutures was observed at a force direction of -30°. Strain approached to a minimum level at a force direction of 30° and 52.4°.

These findings show that the stress distribution in the sutures varies in relation to the direction of force. Force applied in the direction closest to that of the CRe may produce the most effective sutural modification for controlling maxillary growth.

REFERENCES

- [1] Barton JJ. High pull headgear versus cervical traction: a cephalometric comparison. *Am J Orthod* 1972;62(5):517-29.
- [2] Kloehn SJ. Guiding alveolar growth and eruption of teeth to reduce treatment time and produce a more balanced denture and face. *The Angle Orthodontist* 1947;17(1):10-33.
- [3] Proffit WR. Contemporary orthodontics. Mosby: St. Louis, 2001:494-508.
- [4] Nanda R, Hickory W. Zygomaticomaxillary suture adaptations incident to anteriorly-directed forces in rhesus monkeys. *Angle Orthod* 1984;54(3):199-210.
- [5] Wieslander L. The effect of force on craniofacial development. *Am J Orthod* 1974;65(5):531-8.
- [6] Droschl H. The effect of heavy orthopedic forces on the maxilla in the growing Saimiri sciureus (squirrel monkey). *Am J Orthod* 1973;63(5):449-61.
- [7] Elder JR, Rodger HT. Cephalometric and histologic changes produced by extraoral high-pull traction to the maxilla in Macaca mulatta. *Am J Orthod* 1974;66:599-617.
- [8] Meldrum RJ. Alterations in the upper facial growth of Macaca mulatta resulting from high-pull headgear. *Am J Orthod* 1975;67(4):393-411.
- [9] Triftshauser R, Walters RD. Cervical retraction of the maxillae in the Macaca mulatta monkey using heavy orthopedic force. *Angle Orthod* 1976;46(1):37-46.
- [10] Katsaros C, Zissis A, Bresin A, et al. Functional influence of sutural bone apposition in the growing rat. *Am J Orthod Dentofac Orthop* 2006;129(3):352-7.
- [11] Kragt G, Duterloo HS, Bosch JJ. The initial reaction of a macerated human skull caused by orthodontic cervical traction determined by laser metrology. *Am J Orthod* 1982;81(1):49-56.
- [12] Poulton DR. The influence of extraoral traction. *Am J Orthod* 1967;53(1):8-18.
- [13] Armstrong MM. Controlling the magnitude, direction and duration of extraoral force. *Am J Orthod* 1971;59(3):217-43.
- [14] Watson WG. A computerize appraisal of the high-pull face-bow. *Am J Orthod* 1972;62(6):561-79.
- [15] Baumrind S, Korn EL, Molthen R, et al. Changes in facial dimensions associated with the use of forces to retract the maxilla. *Am J Orthod* 1981;80(1):17-30.
- [16] Baumrind S, Korn EL. Patterns of change in mandibular and facial shape associated with the use of forces to retract the maxilla. *Am J Orthod* 1981;80(1):31-47.
- [17] Pfeiffer JP, Grobety D. A philosophy of combined orthopedic-orthodontic treatment. *Am J Orthod* 1982;81(3):185-201.
- [18] Baumrind S, Korn EL, Isaacson RJ, et al. Quantitative analysis of the orthodontic and orthopedic effects of maxillary traction. *Am J Orthod* 1983;84(5):384-98.
- [19] Boecler PR, Riolo ML, Keeling SD, et al. Skeletal changes associated with extraoral appliance therapy: an evaluation of 200 consecutively treated cases. *Angle Orthod* 1989;59(4):263-70.

- [20] Lagerstrom LO, Nielsen IL, Lee R, et al. Dental and skeletal contributions to occlusal correction in patients treated with the high-pull headgear-activator combination. *Am J Orthod Dentofac Orthop* 1990;97(6):495-504.
- [21] Ucem TT, Yuksel S. Effects of different vectors of force applied by combined headgear. *Am J Orthod Dentofacial Orthop* 1998;113(3):316-23.
- [22] Lima Filho RM, Lima AL, de Oliveira Ruellas AC. Longitudinal study of anteroposterior and vertical maxillary changes in skeletal class II patients treated with Kloehn cervical headgear. *Angle Orthod* 2003;73(2):187-93.
- [23] Kragt G, Duterloo HS. The initial effects of orthopedic forces: a study of alterations in the craniofacial complex of a macerated human skull owing to high-pull headgear traction. *Am J Orthod* 1982;81(1):57-64.
- [24] Kragt G, Duterloo HS, Algra AM. Initial displacement and variation of eight human skulls owing to high pull traction determined with laser holography. *Am J Orthod* 1986;89(5):399-406.
- [25] Zentner A, Sergl HG, Filippidis G. A holographic study of variations in bone deformations resulting from different headgear forces in a macerated human skull. *Angle Orthod* 1996;66(6):463-72.
- [26] Billiet T, de Pauw G, Dermaut L. Location of the centre of resistance of the upper dentition and the nasomaxillary complex. An experimental study. *Eur J Orthod* 2001;23(3):263-73.
- [27] Dermaut LR, Kleutghen JP, De Clerck HJ. Experimental determination of the center of resistance of the upper first molar in a macerated, dry human skull submitted to horizontal headgear traction. *Am J Orthod Dentofacial Orthop* 1986;90(1):29-36.
- [28] Verrue V, Dermaut L, Verhegghe B. Three-dimensional finite element modelling of a dog skull for the simulation of initial orthopaedic displacements. *Eur J Orthod* 2001;23(5):517-27.
- [29] Jafari A, Shetty KS, Kumar M. Study of stress distribution and displacement of various craniofacial structures following application of transverse orthopedic forces-a three-dimensional FEM study. *Angle Orthod* 2003;73(1):12-20.
- [30] Cattaneo PM, Dalstra M, Melsen B. The finite element method: a tool to study orthodontic tooth movement. *J Dent Res* 2005;84(5):428-33.
- [31] Tanne K, Matsubara S. Association between the direction of orthopedic headgear force and sutural responses in the nasomaxillary complex. *Angle Orthod* 1996;66(2):125-30.
- [32] Mantysaari R, Kantomaa T, Pirttiniemi P, et al. The effects of early headgear treatment on dental arches and craniofacial morphology: a report of a two year randomized study. *Eur J Orthod* 2004;26(1):59-64.