STUDY OF THE ASSOCIATION OF SERUM 25-HYDROXYVITAMIN D LEVELS WITH THE GLYCAEMIC CONTROL IN TYPE 2 DIABETES MELLITUS AT A TERTIARY CARE HOSPITAL IN GOA

Chitralekha Anilkumar Nayak¹, Anar Viraj Khandeparkar²

¹Senior Resident, Department of Medicine, Goa Medical College.

²Associate Professor, Department of Medicine, Goa Medical College.

ABSTRACT

BACKGROUND

Vitamin D deficiency is widely prevalent across all ages, races, geographical regions and socioeconomic strata. There is ample evidence indicating role of vitamin D in insulin secretion and insulin resistance in patients with type 2 diabetes mellitus. Evidence shows that Vitamin D deficiency can affect the glycaemic control in type 2 diabetes mellitus.

Objective- To study the association between serum 25-hydroxyvitamin D levels with glycaemic control in patients of type 2 diabetes mellitus.

MATERIALS AND METHODS

This case series study was performed on 50 consecutive patients of type 2 diabetes mellitus of 30 to 60 years admitted in General Medicine Ward at Goa Medical College. Serum 25-hydroxyvitamin D levels were assessed using Fully Automated Chemiluminescent Immunoassay (CLIA). Patients with renal failure, type 1 diabetes, chronic illness, pregnancy, thyroid disorders and previous intake of calcium and vitamin D were excluded from the study. The relationship between serum 25-hydroxyvitamin D levels and HbA1c, fasting blood glucose and postprandial blood glucose levels was assessed.

RESULTS

72% of the study population had vitamin D deficiency and 26% of the study group had insufficiency of the vitamin and the remaining 2% had normal 25-hydroxyvitamin D levels. The type 2 diabetic individuals with vitamin D deficiency showed a poor glycaemic control as evidenced by inverse relationship of serum vitamin D levels with FBSL and PPBSL values. There was inverse association between the low serum 25-hydroxyvitamin D levels and elevated HbA1c in the study population.

CONCLUSION

Vitamin D deficiency may be associated with impairment of glycaemic control in type 2 diabetes mellitus.

KEYWORDS

Vitamin D, 25-Hydroxyvitamin D, 25-(OH) D, Diabetes Mellitus, Glycaemic Control.

HOW TO CITE THIS ARTICLE: Nayak CA, Khandeparkar AV. Study of the association of serum 25-hydroxyvitamin D levels with the glycaemic control in type 2 diabetes mellitus at a tertiary care hospital in Goa. J. Evolution Med. Dent. Sci. 2017;6(38):3087-3090, DOI: 10.14260/jemds/2017/666

BACKGROUND

As per WHO estimates, prevalence of diabetes may reach 299 million by year 2025 and it is projected to be the leading cause of death by 2030.¹ Vitamin D deficiency is widely prevalent and it is known to contribute to many conditions like osteomalacia, osteoporosis, falls and fractures. Epidemiological studies have associated low vitamin D status with an increased risk of non-musculoskeletal conditions such as cancer, multiple sclerosis, diabetes mellitus and cardiovascular diseases. Evidence from studies suggests that vitamin D supplementation may help prevent type 2 diabetes mellitus in population at risk of diabetes and also to reduce incidence of complications in known diabetes individuals. The present study was undertaken as limited numbers of studies have attempted to determine the association of serum

25-hydroxyvitamin D levels with glycaemic control in type 2 diabetes mellitus.

Objectives

To study the association between serum 25-hydroxyvitamin D levels and impairment of glycaemic control in patients of type 2 diabetes mellitus.

MATERIALS AND METHODS

The present case series study was performed on 50 consecutive patients of type 2 diabetes mellitus of 30 to 60 years admitted in the Department of Medicine at Goa Medical College from March 2013 to July 2014 upon approval by the Institutional Ethics Committee of the Goa Medical College and Hospital. Informed consent was obtained from all study participants. Serum 25-hydroxyvitamin D levels were measured and its relationship with glycaemic control was assessed among all the study participants.

Inclusion Criteria

- Age between 30 to 60 years.
- Type 2 diabetes mellitus.
- Indian national.

Financial or Other, Competing Interest: None.

Submission 17-03-2017, Peer Review 30-04-2017,

Acceptance 06-05-2017, Published 11-05-2017.

Corresponding Author:

Dr. Chitralekha Anilkumar Nayak,

Rukmini, La Marcela Colony,

Marcela-403107, Goa.

E-mail: nayakchitralekha@gmail.com

DOI: 10.14260/jemds/2017/666

Exclusion Criteria

- Serum creatinine > 1 mg/dL.
- Preexisting chronic cardiac, hepatic, renal and oncologic disease.
- Subjects with type 1 diabetes mellitus.
- Pregnancy and lactation.
- Preexisting thyroid and parathyroid disorders.
- Use of drugs affecting the lipid profile or calcium and bone metabolism.
- History of intake of vitamin D as dietary supplements.
- Preexisting rickets, osteomalacia.
- Patients having disorders that change the metabolism of vitamin D.
- Patients who have been on anticonvulsants, hormone replacement therapy and steroids.

After overnight fasting, 10 mL of peripheral blood was withdrawn. The following biochemical investigations were performed: fasting blood glucose, 2-hour postprandial blood sugar, HbA1c, serum 25-hydroxyvitamin D, calcium, phosphorus, blood urea, serum creatinine, liver function test and lipid profile.

Total 25-hydroxyvitamin D levels were measured in Thyrocare Laboratory, Panaji and analysed on Siemens ADVIA Centaur, standardised against ID-LC/MS/MS, as per Vitamin D Standardisation Program (VDSP). The technology used was Fully Automated Chemiluminescent Immunoassay (CLIA).

Serum 25-Hydroxyvitamin D Levels were interpreted as follows-

1. Less than 20 ng/mL - vitamin D deficiency.
2. 20 to 30 ng/mL - vitamin D insufficiency.
3. 30 - 100 ng/mL - normal levels.
4. > 100 ng/mL - vitamin D toxicity.

All other biochemical investigations were performed in biochemistry laboratory of Goa Medical College using Automated Analyser. Detailed history of the patients was taken regarding occupation, sunlight exposure and intake of foods rich in vitamin D, duration of diabetes, treatment and family history of diabetes. Patients were also evaluated for neuropathy and retinopathy. The data was analysed on SPSS software and correlation coefficients were calculated.

The Study Group was categorised into Groups Based on Duration of Diabetes-

1. Less than 1 year, which also included newly detected subjects.
2. 1 to 5 years.
3. 5 to 10 years.
4. $\geq$ 0 years.

RESULTS

A total of 50 patients were included in this study, 32 being males and 18 females. Mean age was 51.5 ± 7.18 years. Mean age was 52.7 ± 7.08 years in males and 49.2 ± 6.94 years in females; 72% of these diabetic patients were working indoors and 28% were working outdoors. Most of the people who were working indoors were either involved in clerical jobs or were housewives; 60% of the study population had diabetes of duration 1 - 5 years, while 14% were diabetic since 5 - 10

years. In this study, 72% of the subjects were severely deficient in vitamin D and 26% of the subjects were insufficient in vitamin D. The remaining 2% had normal vitamin D levels. This study also showed that 75% of male study groups were vitamin D deficient, while 21.8% had vitamin D insufficiency; 3% of males had normal vitamin D levels; 66% of female subjects were vitamin D deficient and 33.33% were insufficient in vitamin D; 66.66% of severely vitamin D deficient patients were working indoors and 33.34% were working outdoors; 76.92% of vitamin D insufficient were working indoors and 23.08% were working outdoors; 94.44% of the patients with 25-OH-D deficiency and 100% of the patients with insufficiency spent less than 6 hours under sunlight outdoors per day.

Serum Vitamin D Levels (ng/mL)	Males	Females	Total
< 20	24	12	36
20 - 30	7	6	13
30 - 100	1	0	1
> 100	0	0	0

Figure 1. 25-Hydroxyvitamin D Level Distribution among Study Group

Mean Variable	Serum Vitamin D Levels (ng/mL)		
	< 20 ng/mL (Deficiency)	20-30 ng/mL (Insufficiency)	Total (49 Patients)
Age (years)	49.4 ± 4.24	50.24 ± 3.15	51.5 ± 7.18
Duration of Diabetes (yrs.)	4.36 ± 1.28	3.8 ± 1.43	3.88 ± 1.75
FBSL (mg/dL)*	213.86 ± 85.6	200.69 ± 80.78	208.48 ± 84.66
PPBSL (mg/dL)†	321.17 ± 105.1	289.15 ± 100.23	308 ± 112.56
HbA1C (%) §	10.30 ± 2.01	9.3 ± 1.95	9.824 ± 2.36
Serum Calcium (mg/dL)	8.625 ± 1.33	8.669 ± 1.78	8.654 ± 1.86

Figure 2. Characteristics of Participants by Vitamin D Status

*Fasting plasma glucose levels † 2 hours post prandial plasma glucose levels § glycosylated haemoglobin.

The study showed a negative correlation of serum vitamin D levels with FBSL values and glycosylated haemoglobin, but was statistically insignificant. It also showed inverse significant association between vitamin D and post prandial blood glucose levels.

Variable	Statistical Values	
	Correlation Coefficient	P value
FBSL (mg/dL)*	-0.188	0.1905
PPBSL(mg/dL)†	-0.3968	0.00433
HbA1C (%) §	-0.1299	0.435
Serum Calcium (mg/dL)	0.094	0.258

Figure 3. Correlation Coefficients of Variables with 25-Hydroxyvitamin D Levels

*Fasting plasma glucose levels † 2 hours post prandial plasma glucose levels § glycosylated haemoglobin.

DISCUSSION

The present study indicated that 72% of the study population was severely deficient in vitamin D i.e. 25-(OH) D levels < 20 ng/mL, while 26% of the study group was insufficient in vitamin D i.e. 25-(OH) D levels 20 - 30 ng/mL and the remaining 2% had normal (25-OH) D levels. None were found to be in toxic range group, i.e. 25-(OH) D levels > 100 ng/mL. Correlating with the result of the present study was the study by Lalitha A et al² in Vishakhapatnam who observed 38% of subjects being severely deficient (< 10 ng/mL), 42% moderately deficient and 20% being insufficient in the vitamin implying that 80% of the study population was deficient. In contrast to this, severe vitamin D deficiency group comprised of 58.34% and insufficiency of 41.66% in a study done by Saedisomeolia et al³ in Iran. Results of another study by Dalgard et al⁴ on 158 type 2 diabetic patients indicate that more than 50% of the study group was vitamin D deficient, which is actually lower than present study results.

Thus, it is observed that the incidence of vitamin D deficiency is at least 10% higher in the present study and the study by Lalitha A et al² which can be explained by skin colour, lifestyle factors and ethnic differences. An interesting fact was observed that 27% of the vitamin D deficient patients had levels below 10 ng/mL correlating with the previous study by Doddamani et al⁵ where 20% of the patients had such levels.

Data depicted that mean age of vitamin D deficiency patients was 49.4 ± 4.24 years and that of insufficiency being 50.24 ± 3.15 years. Study by Saedisomeolia et al³ also found similar mean age in deficient patients i.e. 47.66 ± 12.38 years and 47.63 ± 11.8 years in insufficient subjects. P value in both studies was statistically insignificant (present study: 0.1073, Saedisomeolia et al 0.74), thereby ignoring age as a factor for vitamin D deficiency. In our study, age had an inverse relationship with serum vitamin D levels but was statistically insignificant probably due to small sample size.

The data showed that 75% of male study group were vitamin D deficient, while 21.8% had vitamin D insufficiency; 3% of males had normal vitamin D levels; 66% of female subjects were vitamin D deficient and 33.33% were insufficient in vitamin D. The difference in number of vitamin D deficient patients between two genders (66% males, 33% females) can be attributed to unequal size of sample population. This finding is in contrast with the results of Saedisomeolia et al study,³ where 42.9% males and 57% females were deficient in vitamin. Of patients with vitamin D insufficiency 53.8% were males and 46.15% were females similar to Saedisomeolia et al³ where 62.2% were males and 37.8% were females. This can be attributed to the equal sample size for both genders in their study. The statistically reverse results in our study may be due to unequal sample population, the type of clothing, hours of sunlight exposure.

25-hydroxyvitamin D levels were suboptimal in diabetic patients pursuing indoor occupation. Data explains 66.66% of severely deficient patients were working indoors and 33.34% were working outdoors; 76.92% of vitamin D insufficient group was working indoors and 23.08% was working outdoors. Similar study in diabetic population is lacking in literature for comparison, but a study by Sari et al⁶ in general

population depicted 72% of vitamin D deficient patients working indoors and 27% working outdoors, results being consistent with the present study.

Another factor which may influence Vitamin D levels is duration of type 2 diabetes mellitus. In this study, mean duration of type 2 diabetes in patients with vitamin D deficiency (< 20 ng/L) was 4.36 years and 3.8 years in those with insufficient vitamin levels respectively. Contrary to this, a study by Christel Joergensen et al⁷ showed mean diabetes duration of 11 years in vitamin deficient subjects and 6 years in patients with insufficient vitamin D levels of > 20 ng/mL. Another study by Ahmadih et al⁸ showed no difference in mean diabetes duration between the two vitamin D groups (Mean duration of diabetes- 8 years). The present study observed a trend of inverse association between vitamin D levels and duration of diabetes, but it was not statistically significant (p value: 0.26). This study had small sample size and excluded nephropathy patients, while in other studies large population was recruited and subjects with nephropathy were also included.

Considering the glycaemic control of diabetes mellitus, the current study showed an inverse but insignificant relationship of serum vitamin D levels and FBSL values ($r = -0.188$, $p = 0.1905$). This was consistent with the results of study done by Balasubramanian et al,⁹ which also showed negative correlation of serum vitamin D levels with fasting blood sugar values and did not show statistical significance ($r = -0.093$, $p = 0.534$). This may be attributed to smaller sample size in our study. Studies by Saedisomeolia et al³ and Doddamani GB et al⁵ showed that mean levels of FBSL were significantly higher in diabetic patients with vitamin D deficiency than those with insufficiency. (Mean FBSL 145 ± 75.66 in deficiency and 129 ± 53.38 insufficiency). The present study revealed similar results. The mean FBSL in deficiency group and insufficiency was 213.86 and 200.69 mg/dL respectively. Contrary to the above, Orwol et al 1994¹⁰ proved no correlation of vitamin D with fasting glucose levels. Many studies conducted on normal glucose tolerant subjects, e.g. studies by Scragg et al 2004,¹¹ Chonchol and Scragg et al 2007,¹² Lu et al 2009,¹³ Need et al 2005,¹⁴ Gamnag Yaered et al 2009, Ford et al 2005¹⁵ also found an inverse relationship of serum vitamin concentration with fasting glucose levels.

The results of the present study revealed a trend towards an inverse vitamin D and post prandial blood glucose levels association, which was statistically significant ($r = -0.3968$, $p = 0.00433$). Similar results were obtained by Balasubramanian et al,⁹ but a statistically insignificant p value was obtained ($r = -0.095$, $p = 0.511$). A study by Chiu et al 2004¹⁶ also showed negative correlation 25 (OH) D levels with PPBSL values. On the contrary, Baynes et al 1997¹⁷ study did not show any association between the two variables.

The low serum vitamin D levels were negatively correlated with glycosylated haemoglobin value, which was not statistically significant probably due to smaller study group. Studies by Balasubramanian et al ($r = -0.173$, $p = 0.229$)⁹ and Dalgard et al⁴ found similar inverse association between the two variables. Contrary to this, Saedisomeolia et al⁶ study and Christel Joergensen et al⁶ study did not show any association between HbA1c and vitamin D levels. In our study, mean HbA1c values were higher in vitamin D deficient

individuals (mean: 10.30%) compared to patients with vitamin insufficiency (mean: 9.3%). Consistent with these results were those of studies by Hypponen and Power et al,¹⁸ Doddamani et al,⁵ Saedisomeolia et al³ and Christel Joergensen et al.⁶

25-Hydroxyvitamin D Levels (ng/L)	HbA1c (Glycosylated Haemoglobin)			
	< 6%	6 - 8%	> 8%	Total
< 20	0	12	24	36
20 - 30	0	5	8	13
30 - 100	0	0	1	1
> 100	0	0	0	0

Figure 4. HbA1c Distribution among 25-Hydroxyvitamin D Level Groups

CONCLUSION

72% of the study population had vitamin D deficiency and 26% of the study group had insufficiency of the vitamin; 27% of the vitamin D deficient patients had very low levels below 10 ng/mL. The present study found no effect of age on serum vitamin D concentration. 25-OH vitamin levels were suboptimal in diabetic patients pursuing indoor occupation compared to those with outdoor occupation. A trend of inverse insignificant association between vitamin D levels and duration of diabetes was observed. There was a positive correlation between serum vitamin D levels and sunlight exposure score and number of hours spent outdoors. The trend towards an association between serum 25-OHD levels and time spent outdoors suggest that production of vitamin D in the skin is the principal determinant of 25-OHD levels. The present study observed that the type 2 diabetic individuals with vitamin D deficiency showed a poor glycaemic control as evidenced by inverse relationship of serum vitamin D levels with FBSL and PPBSL values. The inverse association between the low serum 25(OH) D levels and elevated HbA1c in the study population may be inscribed into a wider context, portraying vitamin D deficiency as a poor prognostic factor, which may play a vital role in impairing the glycaemic control.

REFERENCES

- [1] WHO. Global status report on non-communicable diseases 2010. Geneva: WHO 2011.
- [2] Lalitha A, Tripathi MK. Impact of treating vitamin D deficiency on blood sugar levels in patients with type 2 diabetes and prediabetes. Paper presented at: APICON, 67th Annual Conference of the Association of Physicians of India 2012, Jan 12-15, Kolkata, India.
- [3] Saedisomeolia A, Taheri E, Djalali M, et al. Association between serum level of vitamin D and lipid profiles in type 2 diabetic patients in Iran. J Diabetes Metab Disord 2014;13:7.

- [4] Dalgård C, Petersen MS, Weihe P, et al. Vitamin D status in relation to glucose metabolism and type 2 diabetes in septuagenarians. Diabetes Care 2011;34(6):1284-8.
- [5] Doddamani GB, Umakant Boke, Kora S, et al. Serum vitamin D levels in newly detected type 2 diabetes mellitus. Sch J App Med Sci 2013;1(6):786-8.
- [6] Bajaj S, Singh RP, Swivedi NC, et al. Vitamin D levels and microvascular complications in type 2 diabetes. Indian J Endocrinol Metab 2014;18(4):537-41.
- [7] Joergensen C, Hovind P, Schmedes A, et al. Vitamin D levels, microvascular complications, and mortality in type 1 diabetes. Diabetes Care 2011;34(5):1081-5.
- [8] Ahmadi H, Azar ST, Lakkis N, et al. Hypovitaminosis D in patients with type 2 diabetes mellitus: a relation to disease control and complication. ISRN Endocrinol 2013;2013:641098.
- [9] Shanthi B, Revathy C, Devi AJM, et al. Serum 25 OH D and diabetes mellitus. Journal of Clinical and Diagnostic Research 2012;6(5):774-6.
- [10] Orwoll E, Riddle M, Prince M. Effects of vitamin D on insulin and glucagon secretion in non-insulin-dependent diabetes mellitus. Am J Clin Nutr 1994;59(5):1083-7.
- [11] Scragg R, Sowers M, Bell C. Serum 25-hydroxyvitamin D, diabetes and ethnicity in the Third National Health and Nutrition Examination Survey. Diabetes Care 2004;27(12):2813-8.
- [12] Chonchol M, Scragg R. 25-hydroxyvitamin D, insulin resistance, and kidney function in the Third National Health and Nutrition Examination Survey. Kidney International 2007;71(2):134-9.
- [13] Lu L, Yu Z, Pan A, et al. Plasma 25-hydroxyvitamin D concentration and metabolic syndrome among middle-aged and elderly Chinese individuals. Diabetes Care 2009;32(7):1278-83.
- [14] Tai K, Need AG, Horowitz M, et al. Glucose tolerance and vitamin D: effects of treating vitamin D deficiency. Nutrition 2008;24(10):950-6.
- [15] Ford ES, Ajani UA, McGuire LC, et al. Concentrations of serum vitamin D and the metabolic syndrome among US adults. Diabetes Care 2005;28(5):1228-30.
- [16] Chiu KC, Chu A, Go VL, et al. Hypovitaminosis D is associated with insulin resistance and beta cell dysfunction. Am J Clin Nutr 2004;79(5):820-5.
- [17] Baynes KC, Boucher BJ, Feskens EJM, et al. Vitamin D, glucose tolerance and insulinaemia in elderly men. Diabetologia 1997;40(3):344-7.
- [18] Hypponen E, Power C. Vitamin D status and glucose homeostasis in the 1958 British birth cohort: the role of obesity. Diabetes Care 2006;29(10):2244-6.