COMPARISON OF INTRAVENOUS BOLUS PHENYLEPHRINE, EPHEDRINE AND MEPHENTERMINE FOR MAINTENANCE OF ARTERIAL PRESSURE DURING SPINAL ANAESTHESIA IN CAESAREAN SECTION

M. Srinivasa Rao¹, Satyanarayana², Ramesh³, Saisuraj⁴

¹Associate Professor, Department of Anaesthesiology, SVRR Government General Hospital, Tirupathi, Andhra Pradesh.

²Assistant Professor, Department of Anaesthesiology, S.V. Medical College, Tirupathi, Andhra Pradesh.

³Assistant Professor, Department of Anaesthesiology, S.V. Medical College, Tirupathi, Andhra Pradesh.

⁴Senior Resident, Department of Anaesthesiology, S.V. Medical College, Tirupathi, Andhra Pradesh.

ABSTRACT

BACKGROUND

Caesarean section under spinal anaesthesia is commonly associated with hypotension, which can be detrimental to mother and fetus. It is the responsibility of the anaesthesiologist to ensure stable arterial blood pressure throughout surgery to avoid any decrease in maternal organ blood flow and placental insufficiency. Due to the absence of definitive evidence showing absolute clinical benefit of one over the other, especially in emergency and high-risk caesarean sections, we did examine the effect of intravenous bolus doses of vasopressor agents for maintenance of arterial pressure during spinal anaesthesia in caesarean section today.

MATERIALS AND METHODS

Vasopressors are the cornerstone in treatment of hypotension during spinal anaesthesia. Phenylephrine, α_1 agonist, ephedrine and mephentermine, a direct and indirect acting sympathomimetic can be used to increase the arterial blood pressure. The present study aims to compare the effect of phenylephrine, ephedrine and mephentermine for maintenance of arterial blood pressure.

RESULTS

The study shows that three vasopressors can be used for the indication of hypotension. The clinical outcomes like neonatal Apgar score, incidence of nausea and vomiting and other adverse events were comparable with both the vasopressors. Phenylephrine however has the advantage of decreased heart rate and requirements of lesser amount and more efficacy at maintaining the arterial blood pressure. The systolic blood pressure was significantly higher with phenylephrine 6 minutes after administration compared to mephentermine.

CONCLUSION

Mephentermine should be avoided in patients in whom increased heart rate may be undesirable. Phenylephrine seems to be a better choice for the treatment of hypotension during spinal anaesthesia.

KEYWORDS

Vasopressor Agents, Caesarean Section, Hypotension, Spinal Anaesthesia.

HOW TO CITE THIS ARTICLE: Rao MS, Satyanarayana, Ramesh, et al. Comparison of intravenous bolus phenylephrine, ephedrine and mephentermine for maintenance of arterial pressure during spinal anaesthesia in caesarean section. J. Evolution Med. Dent. Sci. 2018;7(15):1905-1911, DOI: 10.14260/jemds/2018/429

BACKGROUND

Spinal anaesthesia for caesarean section has several advantages over general anaesthesia like decreased risk of failed intubation, decreased risk of pulmonary aspiration of gastric contents, avoidance of the depressant effects of general anaesthetics on neonates, etc.¹

Single shot spinal is most commonly confirmed because it is simple, quicker, has faster onset with superior block and infrequent failure, lesser risk of systemic toxicity due to local anaesthetic agent and lesser transfer to fetus as lower doses are used and its cost effectiveness. However, single shot spinal anaesthesia has its own bag of adverse effects.

'Financial or Other Competing Interest': None.

Submission 10-02-2018, Peer Review 22-03-2018,

Acceptance 29-03-2018, Published 09-04-2018.

Corresponding Author:

Dr. M. Srinivasa Rao,

Associate Professor,

Department of Anaesthesiology,

SVRR Government General Hospital,

Tirupathi,

Andhra Pradesh.

E-mail: manduri60@gmail.com

DOI: 10.14260/jemds/2018/429

The most common adverse effect is hypotension, primarily because of sympathectomy associated with the lumbosacral block. The incidence of hypotension during spinal anaesthesia is as high as 75% - 85%, placental perfusion may be reduced in supine parturient even when mean arterial blood pressure is measured normal. Hypotension during spinal anaesthesia for caesarean delivery may thus further reduce it and may result in foetal acidosis, hypoxia and neurological injury besides maternal nausea and vomiting, dizziness and severe hypotension may result in loss of consciousness and sudden cardiac arrest.² Vasopressors like (Ephedrine, Phenylephrine, Mephentermine) are used for management of hypotension apart from preload of IV fluids.

Recent studies favour phenylephrine, an alpha-1 agonist which elevates arterial blood pressure by increasing systemic vascular resistance secondary to vasoconstriction. Since the primary mode of hypotension during spinal anaesthesia is vasodilation, it seems physiologic to use the vasoconstrictor. However, it causes bradycardia and serial dilution for IV administration is the source of error. It may cause uterine arteriolar constriction and thus diminishing uterine blood flow. Mephentermine has direct and indirect sympathomimetic action and probably increase in arterial

blood pressure is chiefly by increased cardiac output. This may be favourable for placental circulation. Ephedrine increases arterial blood pressure in humans both by peripheral vasoconstriction and by cardiac stimulation. Heart rate, cardiac output is increased. Cardiovascular response to ephedrine may, however, be rather variable. In some cases, arterial blood pressure is not elevated and the peripheral resistance has been reported as increased, decreased or unchanged. Placental transfer of ephedrine occurs and foetal blood levels of approximately 70% of those of maternal blood have been reported. Ephedrine is excreted in breast milk at concentrations that can affect infant.

Aetiopathogenesis of Hypotension

In healthy patients, hypotension during anaesthesia may be defined as a mean arterial pressure of less than 60 mmHg. A systolic blood pressure of 25% less than the patient's preoperative value also indicates hypotension, especially in patients with a pre-existing hypertension. Hypotension is clinically significant if vital organ perfusion is compromised (e.g. myocardial ischaemia or oliguria). As left ventricular coronary blood flow occurs predominantly in diastole, the value of diastolic blood pressure is particularly important in patients with coronary artery disease.

Pathophysiology of Hypotension due to Central Neuraxial Blockade

The principal site of action for neuraxial blockade is the nerve root. Local anaesthetic injected into CSF or the epidural space bathes the nerve root in the subarachnoid space or epidural space respectively. Blockade of neural transmission in the posterior nerve root fibres interrupts somatic and visceral sensations, while blockade of anterior nerve root fibres prevents efferent, motor and autonomic outflow.

Autonomic Blockade

Interruption of efferent autonomic transmission at the spinal nerve roots can produce sympathetic and some parasympathetic blockade. Sympathetic outflow from the spinal cord may be described as thoracolumbar, while parasympathetic outflow is craniosacral. Sympathetic preganglionic nerve fibres exit from the spinal cord with the spinal nerves from T1 to L2 level and may course many levels up or down the sympathetic chain before synapsing with a post-ganglionic cell in a sympathetic ganglion. In contrast, parasympathetic pre-ganglionic fibres exit from the spinal cord with the cranial and sacral nerves. Neuraxial anaesthesia does not block the vagal nerve. Neuraxial blocks therefore primarily result in varying degrees of sympathetic blockade and physiologic responses, resulting from decreased sympathetic and/ or unopposed parasympathetic tone.³

The preganglionic sympathetic fibres arise in the spinal cord between T1 and L2 and travel in the corresponding anterior roots across the subarachnoid and epidural spaces. Sympathetic blockade leads to dilatation of both resistance and capacitance vessels in the region supplied by the blockade nerves. If this is extensive, a fall in peripheral vascular resistance will result together with a posture-dependent fall in cardiac output. The fall in blood pressure accompanying spinal and epidural anaesthesia depends upon height of the blockade, therefore a substantial number of thoracic segments. Sympathetic block extends somewhat

higher than sensory block because B fibres are more sensitive than A and C fibres to local anaesthetics.⁴ There is no difference between spinal and epidural block in this respect, though with epidurals hypotension is usually of slower onset and can be minimised by preloading and by use of small volumes of higher concentrations of local anaesthetic in preference to bigger volumes of lower concentration.

In the presence of sympathetic blockade of the lower part of the body, compensatory vasoconstriction in the upper part can help to maintain the total peripheral resistance, provided that this reflex is not inhibited for example by general anaesthesia. If blockade extends above the level of T5, it becomes progressively more difficult to compensate for the haemodynamic changes and blood pressure will be markedly reduced.⁵ The baroreceptors in the carotid sinus and aortic arch normally respond to a fall in blood pressure by producing a compensatory tachycardia (Marey's law) through vagal afferent and efferent pathways.

Most patients under spinal or epidural anaesthesia, however, exhibit bradycardia and it would appear that the Bainbridge reflex predominates. Thus, in spinal or epidural anaesthesia, venous pooling in the periphery reduces stimulation of the volume receptors thereby diminishing activity of the cardiac sympathetic nerves. The result is vagal preponderance and a slowing of the heart rate. Block of the cardiac sympathetic nerves by either high spinal or epidural block may have beneficial effects. The blood flow to the legs and splanchnic bed are increased, which may confer a considerable advantage compared to general anaesthesia. Thus, the incidence of deep vein thrombosis after hip surgery is significantly reduced by spinal anaesthesia.

Vasodilatation is a predictable and occasionally a desirable feature of blockade involving nerve roots above L2. Serious hypotension is commonly provoked by a high block of rapid onset, because spread of effect may outstrip adequate cardiovascular compensation and fluid preloading. Hypotension therefore may appear as a more prominent feature of spinal than epidural blockade and of blockade with an agent of rapid onset such as chloroprocaine, not because of any differential neural effects by different techniques or drugs.^{6,7,8} Conversely, its incidence can be reduced when an extensive epidural is required by giving the full dose in multiple small increments. Hypotension is minimised by adequate fluid loading which for spinal anaesthesia and for a rapid acting local anaesthetic must be given early. A litre or two of Hartmann's solution is commonly recommended, though colloid solution such as 5% albumin and polygeline (Haemaccel) are effective in reducing hypotension at lower volumes than is crystalloid solution. However, because colloids are more expensive and carry a slightly increased risk of allergic reaction, crystalloid is generally preferred for routine prophylaxis against hypotension.

Factors Influencing the Level of Block

1. Baricity of anaesthetic solution.
2. Position of the patient.
3. Drug dosage.
4. Site of injection.
5. Age.
6. Cerebrospinal fluid volume.
7. Intra-abdominal pressure and curvature of the spine.
8. Patient's height.

Hyperbaric bupivacaine solution most commonly used drug for spinal anaesthesia in caesarean section. Its duration of action is 1.5 to 2 hours. The drug is available in 0.5%, 0.75% and 1%.

Phenylephrine

Is a synthetic non-catecholamine derivative, relatively pure agonist. It is not inactivated by Catechol-O-methyltransferase and has a much longer duration of action than catecholamines. Phenylephrine provides vasoconstriction that lasts longer than that of epinephrine and ephedrine. Responses are more sustained than those to epinephrine, lasting 20 minutes after intravenous and as long as 50 minutes after subcutaneous injection. Its action on the heart contracts sharply with that of epinephrine and ephedrine, in that it slows the heart rate and increases the stroke output producing no disturbance in the rhythm of the pulse.

Phenylephrine is a powerful post-synaptic alpha-receptor stimulant with little effect on the beta receptors of the heart. The predominant actions of phenylephrine are on the cardiovascular system. The drug is a powerful vasoconstrictor with properties very similar to those of epinephrine, but almost completely lacking the chronotropic and inotropic actions on the heart. Cardiac irregularities are seen only very rarely even with large doses.⁹⁻¹⁰

Small intravenous boluses of 50 - 100 micrograms of phenylephrine rapidly reverse blood pressure reductions caused by peripheral vasodilatation. A continuous infusion (100 µg/mL at a rate of 0.25 - 1 µg/kg/min) will maintain arterial blood pressure, but at the expense of renal blood flow. The addition of 2 mg to 5 mg of phenylephrine HCL to the anaesthetic solution increases the duration of motor blockade by as much as approximately 50% without any increase in the incidence of complications such as nausea, vomiting or blood pressure disturbance.

Ephedrine

Is a non-catechol phenylisopropylamine, occurs in various plants. Ephedrine increases arterial blood pressure in humans, both by peripheral vasoconstriction and by cardiac stimulation. Heart rate and cardiac output are increased. Hypoglycaemia has been reported following Ephedrine administration.

In general, the effects of Ephedrine are similar to those observed with Epinephrine but are slower in onset, less potent and longer in duration. Ephedrine is rapidly and completely absorbed after oral administration. Maximum plasma concentration of 45 to 140 mg/L are obtained after a single dose of 22 mg Ephedrine hydrochloride. The mean plasma half-life is about 6 hours. Clearance is 13.6 - 44.3 L/h. Up to 95% of an oral dose may be excreted in the urine in 24 hours. 55% to 75% as unchanged drug and the rest as metabolites. The urinary excretion of Ephedrine is pH dependent and is increased in acidic urine. In alkaline urine, excretion is reduced to 20% - 35% of the dose.

Placental transfer of ephedrine occurs and foetal blood levels of approximately 70% of those of maternal blood have been reported. Ephedrine is excreted in breast milk at concentration that can affect the infant. In adults, ephedrine as an IV bolus of 2.5 - 10 mg. Subsequent doses are increased due to offset of the development of tachyphylaxis, which is probably due to depletion of norepinephrine stores.

Ephedrine is available in 1 mL ampoules containing 25 or 50 mg of the agent.

Mephentermine

Acts by release of noradrenaline at the adrenergic synapses. Cardiac output and systolic and diastolic blood pressures are increased and there is peripheral vasoconstriction. It acts almost immediately after intravenous injection with action of up to about 30 minutes. It is excreted as unchanged drug and metabolites in the urine. IV- 5 to 10 mg bolus, repeat if necessary.

The ideal vasopressor would be one which is reliable and easy to use, has rapid onset, short duration of action, easy, titratable, can potentially be used prophylactically and lack any adverse maternal and foetal impact.

Objectives

To compare the effects of phenylephrine, ephedrine and mephentermine in the management of hypotension during spinal anaesthesia for caesarean section based on the-

1. Efficacy of vasopressor in treating hypotension.
2. Incidence of undesirable side effects.
3. Effect on neonatal outcome.

MATERIALS AND METHODS

After approval by the Institute Ethical Committee, we enrolled singleton full-term patients, aged 20 to 35 yrs. of ASA grade I and II, scheduled for elective caesarean delivery under spinal anaesthesia, consenting to participate in the study at Govt. Maternity Hospital attached to SVMC, Tirupati. Patients having resting BP > 140/100 mmHg, H/O hypertension and having co-existing neurological or cardiovascular disorders, renal, metabolic or psychiatric disorders were excluded. This is a non-randomised clinical study.¹¹

The sample size was taken for convenience and patients were non-randomly allocated as minimum 20 patients per group were necessary to achieve 80% power to detect a 20% difference between the groups. A recruitment target of 60 (20 in each group) was calculated to detect a difference of 6 mmHg in systolic blood pressure (SBP) with 80% power of study and 5% probability of type 1 error. This calculation assumed a standard deviation of 10 mmHg in this parameter within group.^{12,13} The participant was allocated to receive either IV bolus of phenylephrine 100 µg, Ephedrine 5 mg or mephentermine 6 mg to treat hypotension. Hypotension was defined as a fall in systolic blood pressure to a value less than 80% of base value. In order to maintain blinding, the vasopressor solutions were prepared in identical syringe by an anaesthetist who was not involved in the study. The intraoperative monitoring and postoperative observation is done by the same anaesthesiologist who will be administering the drug, but unaware of the contents of the syringe.

Study period was 3 hours after administration of spinal anaesthesia.

Conduct of the Study

The following data was recorded from the selected participant on the morning of surgery, i.e. body weight, 3 readings of the systolic, diastolic blood pressure and HR were obtained at 2 minutes interval with patient at supine with a

15-degree wedge under right hip after the preloading. The lowest reading of BP and highest reading of HR were taken as baseline values. Highest nausea and vomiting score value was taken as baseline.

The participant received bolus of either Phenylephrine 100 µg or Ephedrine 5 mg or mephentermine 6 mg upon developing hypotension. The volume of each dose was equalled by adding 0.9% NaCl solution making the concentration of mephentermine 6 mg/mL and that of phenylephrine 100 µg/ mL. The same, previously calibrated multiple parameter monitor was used for all the participants and for all the readings including the baseline values. Foetal HR was monitored using stethoscope till the dressing and draping of the participant.

Systolic and diastolic blood pressure was noted every 2 mins. after administration of SA (1.8 mL of 0.5% bupivacaine with dextrose 8% solution) till next 10 minutes and every 5 mins thereafter till the completion of surgery or at least 45 mins and every 30 mins for rest of study period. HR was monitored continuously using lead II. Nausea and vomiting and other maternal undesired effects are noted.¹⁴

The time from intrathecal administration of bupivacaine to development of hypotension (SAB-Hypo) from SA to delivery of baby (cord clamping) (SAB-Del) and duration of surgery (Dur-S) was noted. The time of 1st repeat dose of the vasopressor and number of doses was noted. APGAR score of the neonate was noted at 1 and 5 mins of delivery. The data collected were grouped at the end of the whole trial depending on the type of vasopressor. Group P (phenylephrine), Group E (ephedrine) and Group M (mephentermine). The data of failed SA or those who did not develop hypotension were excluded. In appropriate or severe bradycardia was defined as heart rate less than 60 beats/minute if SBP was < 80% of base value, HR less than 50 beats/min when SBP was above the value or heart rate less than 45 beats/min, whatever the blood pressure.

The present study intends to focus on the phenylephrine, ephedrine and mephentermine for maintenance of the arterial blood pressure in women undergoing caesarean delivery under spinal anaesthesia. An examination of the impact of the effect of drugs on arterial blood pressure subjects is also examined in the current study. This chapter includes various methodological aspects which are discussed under the following headings: research design, selection of the sample, sampling technique, inclusion criteria, exclusion criteria, used in the study, procedure of data collection and statistical used in data analysis.¹⁵

This study is used to describe the effect of drugs on arterial blood pressure among caesarean delivery under spinal anaesthesia and also to examine the relationship between those variables. Based on the inclusion and exclusion criteria, 60 patients are taken into the study. They are divided into 3 groups non-randomly using a computer-generated set of numbers before the start of the study, each group containing 20, 20 and 20 respectively. The therapeutic efficacy and safety are assessed by the following assessment scales. A comparison of all the 3 groups is done by using the appropriate statistical procedures mentioned in this protocol, so that results can be obtained.

Statistical analysis was done using SPSS software (version 21.0). The analysis included descriptive statistics and inferential statistics. Descriptive statistics including frequencies, means and standard deviations were used to present the description of the subjects regarding patient's data. Group comparisons between type of vasopressor used and various quantitative variables were conducted by One-Way ANOVA. Results were considered statistically significant when $p < 0.05$.

RESULTS

The groups were comparable in physical characteristic. All the three groups were similar in sensory block level, time to develop hypotension and mean time to delivery and uterine incision to delivery interval.

	Groups		
	Group P	Group E	Group M
Age	23.20 ± 2.608	22.85 ± 2.455	23.60 ± 2.349
Weight	63.50 ± 3.086	63.20 ± 3.019	64.15 ± 3.249
Height	154.35 ± 4.557	155.70 ± 5.592	156.70 ± 5.222
SABHP	4.10 ± .788	4.40 ± .821	4.40 ± .821
SAB-DEL Interval	9.50 ± .688	9.40 ± .598	9.30 ± .571
UI-DEL Interval	1.05 ± .224	1.00 ± .000	1.05 ± .224
Foetal Heart Rate			
Apgar Score of Baby	8 10	8 10	8 10

Table 1. Patient's Characteristic and Relevant Data

Intervals	Systolic Blood Pressure (mmHg)			P-value	Sig. (2-Tailed)
	Group P	Group E	Group M		
HP (VP Given)	82.8 ± 8.02	83.2 ± 7.85	82.6 ± 7.55	p>0.05	Not Significant
2 mins after VP	109.2 ± 8.37	92.4 ± 6.52	92.2 ± 3.55	p<0.001	Significant at 0.001 level
4 mins after VP	112.9 ± 7.24	98.4 ± 3.72	104.2 ± 7.55	p<0.001	Significant at 0.001 level
6 mins after VP	115.1 ± 4.09	106.8 ± 3.37	106 ± 4.85	p<0.001	Significant at 0.001 level
8 mins after VP	116.2 ± 4.60	110.5 ± 3.50	107.8 ± 7.03	p<0.001	Significant at 0.001 level
10 mins after VP	116.5 ± 4.72	111.3 ± 7.75	111 ± 4.52	p<0.05	Significant at 0.05 level
15 mins after VP	114.3 ± 8.90	112.8 ± 7.35	110.8 ± 8.62	p>0.05	Not Significant
At 30 mins after VP	114.5 ± 7.80	109.1 ± 12.45	105.6 ± 14.75	p>0.05	Not Significant
At the end of the surgery	117.1 ± 3.43	117.4 ± 3.65	117.4 ± 3.8	p>0.05	Not Significant

Table 2. Changes in Systolic BP (mean ± SD)

Figure 1. Systolic BP

VP- Vasopressor agents.

On intergroup comparison rise of systolic blood pressure at 2, 6, 8 and 10 minutes post study drug were less in ephedrine and mephentermine groups as compared to phenylephrine group ($p < 0.001$).

Intervals	Diastolic Blood Pressure (mmHg)			P-value	Sig. (2-Tailed)
	Group P	Group E	Group M		
HP (VP given)	61.8 ± 7.72	61.2 ± 7.59	58.2 ± 5.90	$p > 0.05$	Not Significant
2 mins after VP	72.5 ± 4.04	66.4 ± 5.59	65.40 ± 4.90	$p < 0.001$	Significant at 0.001 level
4 mins after VP	74.62 ± 3.03	69.5 ± 4.88	69.32 ± 3.75	$p < 0.001$	Significant at 0.001 level
6 mins after VP	76.12 ± 1.78	72.6 ± 2.05	70.8 ± 2.82	$p < 0.001$	Significant at 0.001 level
8 mins after VP	77.25 ± 2.32	75.5 ± 2.05	72.0 ± 4.60	$p < 0.001$	Significant at 0.001 level
10 mins after VP	76.42 ± 2.71	76.2 ± 3.08	74 ± 3.85	$p > 0.05$	Not Significant
15 mins after VP	76.00 ± 8.90	77.5 ± 3.42	74.4 ± 5.69	$p > 0.05$	Not Significant
At 30 mins after VP	76.7 ± 7.70	74.12 ± 6.50	72.1 ± 8.9	$p > 0.05$	Not Significant
At the end of the surgery	79.10 ± 3.33	78.5 ± 1.53	78.8 ± 3.42	$p > 0.05$	Not Significant

Table 3. Changes in Diastolic BP (mean ± SD)

Figure 2. Diastolic BP

Intervals	Heart Rate Changes			P-value	Sig. (2-Tailed)
	Group P	Group E	Group M		
HP (VP given)	101.85 ± 9.40	98.2 ± 7.85	96.1 ± 5.81	$p > 0.05$	Not Significant
2 mins after VP	96.60 ± 7.25	100.40 ± 9.20	96.50 ± 8.21	$p > 0.05$	Not Significant
4 mins after VP	92.40 ± 6.85	103.32 ± 10.45	95.85 ± 8.10	$p < 0.001$	Significant at 0.001 level
6 mins after VP	90.70 ± 7.10	104.5 ± 13.06	97.20 ± 12.80	$p < 0.001$	Significant at 0.001 level
8 mins after VP	89.60 ± 8.65	101.45 ± 8.50	97.70 ± 12.80	$p < 0.05$	Significant at 0.05 level
10 mins after VP	86.35 ± 10.54	96.6 ± 13.85	96.4 ± 9.65	$p < 0.05$	Significant at 0.05 level
15 mins after VP	82.85 ± 6.40	96.0 ± 6.50	94.50 ± 12.56	$p < 0.001$	Significant at 0.001 level
At 30 mins after VP	84.10 ± 6.50	94.25 ± 5.25	96.3 ± 9.67	$p < 0.001$	Significant at 0.001 level
At the end of the surgery	80.20 ± 3.20	81.36 ± 4.02	89.85 ± 5.72	$p < 0.001$	Significant at 0.001 level

Table 4. Changes in Heart Rate (mean ± SD) mmHg

Figure 3. Heart Rate

In Group P, post study drug values of HR were decreased significantly from the values at onset of hypotension till the end of the surgery. Whereas in Group E values remained high upto 10 mins and then decreased (< 0.05). In Group M, rates were higher after study drugs and remain statistically significant with values at onset of the hypotension till end of the surgery.

In Group P 80% patients require single bolus dose, while 15% two and 5% three to maintain SBP within 20% limit of base value. In Group E 45% required single, 45% required two and 10% three bolus doses. Whereas in Group M 55% required single, 40% two and 5% three doses. One patient (5%) in each group developed bradycardia. In Group P and E, 10% developed nausea and vomiting. Apgar score did not reveal any untoward effect on foetal status, since all new born of three groups had Apgar score greater than 7.

Sab-Delivery Time Sab-Hypo Time

Group P	4.1 ± 0.79
Group E	4.4 ± 0.82
Group M	4.4 ± 0.82
P-value	$p > 0.05$

Group P	9.5 ±0.68
Group E	9.4 ±0.59
Group M	9.3 ±0.57
P-value	p>0.05

DISCUSSION

The current study was conducted to compare the three vasopressors for their efficacy at maintaining the arterial blood pressure. The method of non-randomisation was allocation by sealed envelope technique. Data was collected and the results were summarised and analysed statistically by appropriate method.

The demographic data, baseline parameters were comparable in all groups, the haemodynamic parameters at the development of hypotension were also comparable between the groups. The mean time for development of hypotension after SA in Group P was 4.1 ± 0.79 (Mean $\pm$ SD), 4.4 ± 0.82 (Mean $\pm$ SD) minutes in Group M and 4.4 ± 0.82 in Group E. The values were comparable with $p > 0.05$. After administration of the vasopressors, the SBP was comparable between the groups at all time points, except at 10 minutes when the mean SBP was significantly higher in Group P (116 ± 4.72 ; 113 ± 7.75 ; 114 ± 4.52 ; $p < 0.05$).

Sahu D and colleagues, however, observed mean SBP to be statistically higher after 2 minutes when phenylephrine was the vasopressor in comparison to mephentermine 116.5 ± 14 vs 106.9 ± 11 (mean $\pm$ SD), $p < 0.05$. The mean DBP was significantly higher in Group P in comparison to Group M at all time points after administration of vasopressors in the current study, where it was observed to be comparable by Sahu D and colleagues.

The mean HR decreased significantly after administration of phenylephrine as compared to mean HR at the onset of hypotension, where it remained high in Group E and was statistically significant at 0.001 level. Within the group analysis of SBP shows peak effect of phenylephrine to occur after 6 to 8 minutes, while that of mephentermine after 8 minutes. The incidence of nausea and vomiting and other effects was comparable between the three groups. The Apgar score at 1 and 5 minutes was also statistically not significant between the groups.

CONCLUSION

The primary mechanism of elevation of arterial BP by phenylephrine is by vasoconstriction due to predominately α_1 agonist activity, whereas mephentermine increases MAP by increasing HR and myocardial contractility due to its α and β agonist activity. We conclude from the study that phenylephrine, ephedrine and mephentermine maintain SBP above hypotensive range, though phenylephrine might be better because number of doses needed is less. Phenylephrine increases DBP more than mephentermine and hence MAP is increased. Thus, it can probably enhance organ blood flow more than mephentermine. Mephentermine increases HR and thus may be avoided in population, where the effect may be detrimental.

REFERENCES

- [1] Riley ET, Cohen SE, Rubenstein AJ, et al. Prevention of hypotension after spinal anesthesia for cesarean section: six percent hetastarch versus lactated Ringer's solution. *Anesth Analg* 1995;81(4):838-42.
- [2] Lee A, Kee NWD, Gin T. A quantitative, systematic review of randomized controlled trials of ephedrine versus phenylephrine for the management of hypotension during spinal anesthesia for cesarean delivery. *Anesth Analg* 2002;94(4):920-6.
- [3] Ganeshanavar A, Uday AS, Adarsh SE, et al. Comparison of bolus phenylephrine, ephedrine and mephentermine for maintenance of arterial pressure during spinal anaesthesia in caesarean section. *Journal of Clinical and Diagnostic Research* 2011;5(5):948-52.
- [4] Kee NWD, Khaw KS, Lee BB, et al. A dose-response study of prophylactic intravenous ephedrine for the prevention of hypotension during spinal anesthesia for cesarean delivery. *Anesth Analg* 2000;90(6):1390-5.
- [5] Corke BC, Datta S, Ostheimer GW, et al. Spinal anaesthesia for Caesarean section. The influence of hypotension on neonatal outcome. *Anaesthesia* 1982;37(6):658-62.
- [6] Ramanathan S, Grant GJ. Vasopressor therapy for hypotension due to epidural anaesthesia for caesarean section. *Acta Anaesthesiol Scand* 1988;32(7):559-65.
- [7] Thomas DG, Robson SC, Redfern N, et al. Randomized trial of bolus phenylephrine or ephedrine for maintenance of arterial pressure during spinal anesthesia for Caesarean section. *Br J Anaesth* 1996;76(1):61-5.
- [8] Simin A, Zahra F, Pouya H, et al. Comparison the effect of ephedrine and phenylephrine in treatment of hypotension after spinal anesthesia during cesarean section. *Open Journal of Obstetrics and Gynaecology* 2012;2(3):192-6.
- [9] Alday ME, Palacio AF, De Diego PR, et al. Ephedrine vs. phenylephrine by intravenous bolus and continuous infusion to prevent hypotension secondary to spinal anesthesia during cesarean section: a randomized comparative trial. *Rev Esp Anestesiología Reanim* 2011;58(7):412-6.
- [10] Nazir I, Bhat MA, Qazi S, et al. Comparison between phenylephrine and ephedrine in preventing hypotension during spinal anesthesia for cesarean section. *Journal of Obstetric Anaesthesia and Critical Care* 2012;2(2):92-7.
- [11] Kansal A, Mohta M, Sethi AK, et al. Randomised trial of intravenous infusion of ephedrine or mephentermine for management of hypotension during spinal anaesthesia for Caesarean section. *Anaesthesia* 2005;60(1):28-34.
- [12] Bhattarai B, Bhat SY, Upadya M. Comparison of bolus phenylephrine, ephedrine and mephentermine for maintenance of arterial pressure during spinal anesthesia in cesarean section. *JNMA J Nepal Med Assoc* 2010;49(177):23-8.
- [13] Bhardwaj N, Jain K, Arora S, et al. A comparison of three vasopressors for tight control of maternal blood pressure during cesarean section under spinal anesthesia: effect on maternal and fetal outcome. *Journal of Anaesthesiology Clinical Pharmacology* 2013;29(1):26-31.

- [14] Lin FQ, Qiu MT, Ding XX, et al. Ephedrine versus phenylephrine for the management of hypotension during spinal anesthesia for cesarean section: an updated meta-analysis. *CNS Neurosci Ther* 2012;18(7):591-7.
- [15] Hall PA, Bennett A, Wilkes MP, et al. Spinal anaesthesia for caesarean section: comparison of infusions of phenylephrine and ephedrine. *Br J Anesth* 1994;73(4):471-4.