BACTERIOLOGICAL PROFILE AND THE ANTIBIOTIC SUSCEPTIBILITY PATTERN OF ENDOTRACHEAL SECRETIONS IN ICU OF A TERTIARY CARE HOSPITAL

Pooja Gupta¹, Shilpa Gupta², Jung Bahadur Singh³

¹Consultant Microbiologist, Department of Microbiology, Max Super Speciality Hospital, Bathinda, Punjab.

²Consultant, Department of Internal Medicine, Max Super Speciality Hospital, Bathinda, Punjab.

³Consultant, Department of Internal Medicine, Max Super Speciality Hospital, Bathinda, Punjab.

ABSTRACT

BACKGROUND

Patients on mechanical ventilation are at higher risk of acquiring hospital acquired infection due to interplay of compromised host defense, virulent organism and presence of invasive device. Knowledge of local bacteriological trend of the antibiotic susceptibility pattern is a must to initiate a judicious antibiotic treatment.

MATERIALS AND METHODS

The study was a retrospective descriptive study analysis of endotracheal (ET) samples of patients on mechanical ventilation done over a period of 6 months from July 2017 to Dec 2017. This was done to analyse bacteriological profile and their antibiotic susceptibility.

RESULTS

44 out of 82 samples showed growth accounting for total of 53% (positive cultures). Gram negative bacteria were the most common isolates with *Klebsiella spp.* being the most common bacteria followed by *Pseudomonas aeruginosa*. *Acinetobacter baumannii*, though the most resistant organism was a rare isolate. *Klebsiella* was the most common as well as an MDR (multidrug resistant) isolate.

CONCLUSION

Gram negative bacteria are the major isolates from ET secretions of patients on mechanical ventilation in our ICU setup, *Klebsiella* being the most common and resistant too. An antibiogram specific for healthcare setting is a must that shall be based on bacteriological profile and organism's sensitivity pattern.

KEYWORDS

Endotracheal Aspirate, MDR, *Klebsiella Pneumoniae*, Mechanical Ventilation, Gram Negative Bacteria.

HOW TO CITE THIS ARTICLE: Gupta P, Gupta S, Singh JB. Bacteriological profile and the antibiotic susceptibility pattern of endotracheal secretions in ICU of a tertiary care hospital. J. Evolution Med. Dent. Sci. 2018;7(18):2210-2213, DOI: 10.14260/jemds/2018/497

BACKGROUND

Invasive devices continue to be essential for the management of critically ill patients, despite breath-taking advances in modern day medicine.¹ These ubiquitous devices unfortunately are a major cause of nosocomial infections, especially in the ICU.² As a result following these invasive methods, the incidence of nosocomial infections in ICU has raised.³

These infections are associated with significant rise in morbidity, mortality and healthcare cost. An additional problem is the emergence and rise in multi-drug resistant (MDR) pathogens. Several factors like new mutations, selection of resistant strains and suboptimal infection control along with the use high level antibiotics influences the rapid spread of MDR bugs in the ICU.⁴ In patients on mechanical ventilation with intubation, frequent colonisation occurs either through exogenous or endogenous origin.⁵

'Financial or Other Competing Interest': None.
Submission 17-03-2018, Peer Review 12-04-2018,
Acceptance 18-04-2018, Published 30-04-2018.

Corresponding Author:

Dr. Pooja Gupta,

Consultant Microbiologist,

Department of Microbiology,

Max Super Speciality Hospital,

Bathinda-151001, Punjab.

E-mail: Pooja.Gupta@maxhealthcare.com

DOI: 10.14260/jemds/2018/497

Ventilator-associated pneumonia (VAP) is the second-most common hospital-acquired infection (HAI), accounting for 15% of HAIs and has the highest morbidity and mortality.⁶ A large number of bacteria are found to be causative factors for colonisation that may later proceed to infection.⁷ Gram negative bacteria are predominant ones. Most common gram-negative bacteria are *Pseudomonas aeruginosa*, *Acinetobacter spp.* and *Klebsiella pneumoniae*.^{8,9} Prolonged hospital stay and colonisation with *Pseudomonas* species itself is an independent risk factor for gram negative bacterial infection.

The longer the hospital stay is as in case of head trauma, the more are the chances of change of bacterial flora causing infection, thereby complicating therapy because of alteration of susceptibility pattern.^{10,11}

It is very essential for the clinicians to be aware of local bacteriological flora and their susceptibility pattern to encourage rational use of antibiotics. The aim of our study was to present a data that represents the local bacterial trend in respiratory secretions of the ventilated patients and their antibiotic susceptibility patterns.

MATERIALS AND METHODS

The study was a retrospective descriptive study conducted in 2 ICUs; medical (MICU) and surgical (SICU) of a tertiary care hospital over a period of six months from July 2017 to Dec 2017. The analysis of the reports was done at the Microbiology Department.

Samples were endotracheal secretions collected from patients, who were on mechanical ventilation for more than 2 days for various reasons. Patients were selected regardless of age and sex. Samples were aseptically inoculated on blood agar and MacConkey agar. The plates were further incubated aerobically at 37°C overnight. Significant growth was considered when $> 10^5$ colonies were obtained from tracheal secretions.¹² The antibiotics were used as per the strips suitable for the kind of growth as per standard protocol.

A total of 82 samples were collected during the study period. A total of 44 samples showed significant growth when cultured using standard microbiological techniques.¹³ Significant growth was considered when $> 10^5$ colonies were obtained.¹² Organisms were identified by automated system; ATB reader and their antibiotic susceptibility was done as per latest CLSI guidelines 2017.¹⁴ Exclusion criteria were the isolates identified as contaminant or same isolates from repeat culture of patient. Our gram negative bacterial isolates were defined as multi-drug resistant (MDR) when they were resistant to at least 3 different types of antimicrobials.¹⁵

RESULTS

The percentage of positive cultures was approximately 53% (44 positive cultures out of 82 samples). The most common bacteria isolated were *Klebsiella pneumoniae* (50%), *Pseudomonas aeruginosa* (25%), *Escherichia coli* (9%), *Enterococcus faecalis* (9%), *Staphylococcus aureus* (4.5%) and *Acinetobacter baumannii* (2.5%). Thus, gram negative bacteria (GNB) contributed to major number of isolates (86.5%); the remaining 13.5% was as a result of gram positive infections. Moreover, amongst GNB, Enterobacteriaceae (*K. pneumoniae* and *E. coli*) accounted for 59%, while non-fermenters (*P. aeruginosa* and *A. baumannii*) comprised the rest (Fig. 1).

ICU wise, the common isolates in MICU were *Pseudomonas* and *E. coli* followed by *Klebsiella*, while in

NSICU *Klebsiella pneumoniae* was the most common isolate. The only *Acinetobacter* isolate was isolated from NSICU.

Figure 1. Bacteriological Profile of ET Secretions in MICU and NSICU

Antibiotic Susceptibility Profile

Amongst gram negative bacteria, *E. coli* was the least resistant organism being sensitive to aminoglycosides and cotrimoxazole followed by *P. aeruginosa*. On the other hand, *Klebsiella* isolates and *Acinetobacter* spp. were resistant to lower antibiotics namely cephalosporins, aminoglycosides and fluoroquinolones. While some *K. pneumoniae* isolates showed susceptibility to Tigecycline and few to carbapenems, *A. baumannii* was only sensitive to polymyxins.

Amongst gram positive bacteria, 100% susceptibility was shown to higher antibiotics viz. teicoplanin, linezolid and Vancomycin, while resistance was observed towards beta lactam antibiotics. The resistance was high towards fluoroquinolones for both gram positive and gram-negative isolates (Table 1).

None of the patients who were mechanically ventilated in our study period developed VAP.

	Pathogens	No. of Isolates	% of Isolates	Sensitive	Resistant
1.	<i>Klebsiella</i>	22	50	PB, Col (100%) Tgc- (70%), I, Mr (55%)	All cephalosporins, aminoglycosides, FQ
2.	<i>Pseudomonas</i>	11	25	PB, Col (100%), I, Mr (100%), CfS (100%), Ak (50%), PTz (50%)	Cephalosporins (100%), FQ (100%), CoT (100%)
3.	<i>Enterococcus faecalis</i>	4	9	Va, Lz, Tei (100%), PTz (50%), Tetra (70%),	Beta Lactams (100%), FQ (100%)
4.	<i>E. coli</i>	4	9	Col (100%), I/Mr (100%), PTz (75%), CfS (75%), Ak (50%),	Amc, Cip, CoTri
5.	<i>S. aureus</i>	2	4.5	Va, Lz, Tei (100%), PTz (50%), Tetra (70%), I (100%), CD-60%	Cz, E, Cip, Penicillin
6.	<i>A. baumannii</i>	1	2.5	PB, Col (100%)	Carbapenems, FQ, Aminoglycosides, cephalosporins
7.	No growth	38			

Table 1. Antibiotic Susceptibility Profile of Isolates in ET Secretions

AK= Amikacin, GEN= Gentamycin, CIP= Ciprofloxacin, IMI= Imipenem, AMC= Amox-co=Clavulanate, COT= Cotrimoxazole, PB= Polymyxin B, Col= Colistin, Mr= Meropenem, FQ= Fluoroquinolones (Ciprofloxacin, Ofloxacin), PTz (Piperacillin=tazobactam), Va= vancomycin, Lz (linezolid), Tei= teicoplanin, tetra= tetracycline, Cz= (cefazolin).

DISCUSSION

Endotracheal intubation and mechanical ventilation are life-saving procedures done to prevent or combat respiratory failure.¹⁶ On the other hand, it may even lead to life-threatening lung infections due to various reasons that bypass the host's immune response, whereby organisms enter the airways by exogenous or endogenous route causing ventilator associated pneumonia (VAP). In our study, we chose samples from the patients who were on mechanical ventilation for more than 2 days. This would help us identify VAP.

During our study period, the VAP count was 0 as per the CDC definition. Joseph et al study in Pondicherry showed a high incidence rate of 22.4 per 1000 ventilator days.¹⁷

There have been many studies done in the Indian subcontinent that have identified the aetiological agents of VAP as well as the susceptibility pattern which has been showing increasing resistance. Most common VAP pathogens are *P. aeruginosa*, *Acinetobacter spp.*, *E. coli*, *K. pneumoniae* and *S. aureus*.¹⁵ In our study, though *Klebsiella spp.* was the most common organism. Nishat et al study had *Pseudomonas*, *Acinetobacter spp.* and *Klebsiella pneumoniae* as most common isolates (26% each) followed by *E. coli* and *S. aureus*⁽¹⁵⁾ and 86.3% isolates were MDR. In our study *Klebsiella* and *Acinetobacter spp.* were MDR, while rest of the isolates were quite sensitive except to the beta-lactam antibiotics and fluoroquinolones. In Anusha et al study, *Klebsiella* was the most common bacteria followed by *Pseudomonas*, *Acinetobacter* and *E. coli*, just like our findings.¹⁶ But their isolates were mostly susceptible to aminoglycosides and carbapenem class. In our study, *Klebsiella* isolates were resistant to aminoglycosides.

Summaiya M et al, while assessing biofilm formation by VAP aetiological agents found that the most common isolates are *Pseudomonas aeruginosa* and *Acinetobacter spp.*¹⁸ While Trilok Patil study at Aurangabad found *Pseudomonas aeruginosa* to be the most common organism (37.4%) followed by *Klebsiella pneumoniae* (28.5%).¹⁹ George P et al study had *Acinetobacter* as the most common isolate (37.5%) followed by *Pseudomonas* (21.8%) and *Klebsiella* (5%).²⁰ These studies differed from our observation of *Klebsiella* being the most common organism in endotracheal secretions. As per Deepti et al study, Gram negative enteric aerobic bacteria were isolated from most of the patients, most common being *Klebsiella* species (32.35%) followed by *Acinetobacter* and *Pseudomonas*.²¹ In their study Gram positive cocci were mostly sensitive to Penicillin derivatives, Vancomycin and Clindamycin, while our GPC were resistant to Penicillin derivatives.²¹ These variations in the aetiological bacteria can be due to varied population in ICU, their prior antibiotic therapy and the commensal carriage.

Gram negative bacteria as the most common isolates in ET secretion was similar in various studies, but the multi-drug resistant property of *Klebsiella* isolates and resistance of gram positive bacteria to penicillins in our study is alarming.

The limitation of the study is that the study was of short duration and a retrospective one. The results are not applicable to generalised healthcare settings, since the findings are from a single tertiary care hospital. The BAL sample was not included in samples and the outcome of patients could not be monitored.

CONCLUSION

Gram negative bacteria form the predominant isolates in our ICU setup; the worrisome being *Klebsiella* isolates which were resistant to even carbapenem group of antibiotics. Clinicians shall be aware of local bacteriological profiles and their antibiotic sensitivity pattern since that helps them in initiation of an appropriate antibiotic therapy. Thus, an updated antibiogram for a hospital and timely ET secretion culture are the keys to reduce emergence of resistant microbes in patients on mechanical ventilation.

REFERENCES

- [1] Safdar N, Crnich CJ, Maki DG. Nosocomial infections in the intensive care unit associated with invasive medical devices. *Curr Infect Dis Rep* 2001;3(6):487-95.
- [2] Anonymous, National Nosocomial Infections Surveillance (NNIS) system report, data summary from Jan 1992-April 2000. *Am J Infect Control* 2000;28:429-48.
- [3] Mehta A, Rosenthal VD, Mehta Y, et al. Device associated nosocomial infection rates in intensive care units of seven Indian cities. Findings of the international nosocomial infection control consortium (INICC). *J Hosp Infect* 2007;67(2):168-74.
- [4] Brusselaers N, Vogelaers D, Blot S. The rising problems of antimicrobial resistance in the intensive care unit. *Ann Intensive Care* 2011;1:47.
- [5] Vadivoo NS, Santharam P, Sudha K, et al. Dynamic bacterial profile of endotracheal aspirates and its sensitivity pattern-a cause of concern. *Int J Curr Res Rev* 2014;6(10):112-9.
- [6] Fiel S. Guidelines and critical pathways for severe hospital-acquired pneumonia. *Chest* 2001;119(2):412S-8.
- [7] Mukhopadhyay C, Bhargava A, Ayyagari A. Role of mechanical ventilation and development of multidrug resistant organisms in hospital acquired pneumonia. *Indian J Med Res* 2003;118:229-35.
- [8] Bonten MJ, Kollef MH, Hall JB. Risk factors for ventilator associated pneumonia: from epidemiology to patient management. *Clin infect Dis* 2004;38(8):1141-9.
- [9] Tullu MS, Deshmukh CT, Baveja SM. Bacterial profile and antibiotic susceptibility pattern in catheter related nosocomial infections. *JPGM* 1998;44(1):7-13.
- [10] Gotsman MS, Whitby JL. Respiratory infections following tracheostomy. *Thorax* 1964;19(1):89-96.
- [11] Kaul S, Brahmadathan KN, Jagannati M, et al. One year trends in the gram-negative bacterial antibiotic susceptibility pattern in a medical intensive care unit in South India. *Indian J Med Microbiol* 2007;25(3):230-5.
- [12] Tablan OC, Anderson LJ, Besser R, et al. Guidelines for preventing health-care-associated pneumonia, 2003: recommendations of CDC and the healthcare infection control practices advisory committee. *MMWR Recomm Rep* 2004;53(RR-3):1-36.

- [13] Collee JG, Marr W. Culture of bacteria. In: Marmion BP, Fraser AG, Simmons A. (eds). Mackie and McCartney's practical medical microbiology. 14th edn. New York: Churchill Livingstone, 1996:113–30.
- [14] Clinical and laboratory standards institute (CLSI). Performance standards for antimicrobial susceptibility testing; twenty seventh informational supplement. CLSI document 2017:M100-S26.
- [15] Ahmed NH, Hussain T, Biswal I. Antimicrobial resistance of bacterial isolates from respiratory secretions of ventilated patients in a multi-specialty hospital. *Avicenna J Med* 2015;5(3):74–8.
- [16] Anusha N, Madhu KP, Arun BJ, et al. Microbiological profile and sensitivity pattern of endotracheal secretions in mechanically ventilated patients in ICU. *Journal of Evidence Based Medicine And Healthcare* 2014;1(9):1177-84.
- [17] Joseph NM, Sistla S, Dutta TK, et al. Ventilator-associated pneumonia in a tertiary care hospital in India: incidence and risk factors. *J Infect Dev Ctries* 2009;3(10):771–7.
- [18] Summaiya MA, Urmi JN. Assessment of biofilm formation by the causative organisms of ventilator associated pneumonia at intensive care unit of a tertiary care hospital. *Natl J Med Res* 2012;2(1):15-8.
- [19] Patil T. The study of the organisms colonizing trachea in mechanically ventilated patients admitted in the intensive care unit (ICU). *International Journal of Medical Science and Education* 2014;1(1).
- [20] George P, Sequiera A. Antimicrobial sensitivity pattern among organisms which were isolated from the endotracheal aspirates of patients with ventilator associated pneumonia. *J Clin Diag Res* 2010;4(6):3400-404.
- [21] Chandra D, Laghawe A, Sadawarte K, et al. Microbiological profile and antimicrobial sensitivity pattern of endotracheal tube aspirates of patients in ICU of a tertiary care hospital in Bhopal, India. *Int J Curr Microbiol App Sci* 2017;6(3):891-5.