AN OBSERVATIONAL STUDY OF CLINICAL PROFILE AND MANAGEMENT OF NON-TRAUMATIC SMALL BOWEL PERFORATION AT TERTIARY CARE CENTRE

Kiran Somani¹, Rishikant Vashistha², Sanjay Datey³, Abhishek Gurjar⁴, Akhilesh Patel⁵

¹Professor, Department of General Surgery, Sri Aurobindo Medical College and Postgraduate Institute, Indore, Madhya Pradesh, India.

²Associate Professor, Department of General Surgery, Sri Aurobindo Medical College and Postgraduate Institute, Indore, Madhya Pradesh, India.

³Professor, Department of General Surgery, Sri Aurobindo Medical College and Postgraduate Institute, Indore, Madhya Pradesh, India.

⁴Senior Resident, Department of General Surgery, Sri Aurobindo Medical College and Postgraduate Institute, Indore, Madhya Pradesh, India.

⁵Senior Resident, Department of General Surgery, Sri Aurobindo Medical College and Postgraduate Institute, Indore, Madhya Pradesh, India.

ABSTRACT

BACKGROUND

Perforation peritonitis is a common surgical emergency faced by general surgeon in India. It is associated with significant morbidity and high mortality rate. This work has been carried out to study various aetiology, clinical feature, diagnostic dilemmas and management of non-traumatic small bowel perforation at a tertiary care centre. Traumatic small bowel perforations were not included in the study.

MATERIALS AND METHODS

This cross-sectional observational study was performed at a tertiary care centre. All patients undergoing exploratory laparotomy for non-traumatic small bowel perforation were included in the study from 1st January 2015 to 30th June 2016. The patient's demographic data, clinical profile, diagnostic studies, intraoperative findings during laparotomy, surgical interventions performed, postoperative complications and final outcome were recorded and evaluated.

RESULTS

A total of 80 patients, 21 females and 59 males, between age ranges of 18 years and 90 years underwent exploratory laparotomy for non-traumatic small bowel perforation. Symptoms were abdominal pain, obstipation, abdominal distension, fever and vomiting. The cause of perforation was typhoid in 66 patients followed by ischaemic bowel disease in 6 and tuberculosis in 5 patients, and malignancy, intussusceptions, worm infestation in 1 patient each. 67 patients had ileal perforation, while 13 patients had jejunal perforation. 67 patients had single perforation, while 13 patients had multiple perforations. Various surgical procedures according to intraoperative findings were performed, 90% patients were discharged.

CONCLUSION

Typhoid fever is the most common cause of non-traumatic small bowel perforation followed by ischaemic bowel disease and tuberculosis in our study. Early surgical intervention under the cover of broad-spectrum antibiotics preceded by aggressive resuscitation and correction of electrolyte imbalance is imperative for good outcomes minimising morbidity and mortality.

KEY WORDS

Small Bowel Perforation, Typhoid, Exploratory Laparotomy.

HOW TO CITE THIS ARTICLE: Somani K, Vashistha R, Datey S, et al. An observational study of clinical profile and management of non-traumatic small bowel perforation at tertiary care centre. J. Evolution Med. Dent. Sci. 2018;7(32):3581-3584, DOI: 10.14260/jemds/2018/804

BACKGROUND

Non-traumatic small bowel perforation is the most common form of gastrointestinal perforation in developing and underdeveloped nations.¹ Perforation peritonitis is the most common emergency faced by general surgeon in India.² Small bowel perforation is an acute surgical emergency, which may lead to multiorgan failure, sepsis and death if appropriate aggressive intervention is not done in a limited time.

Financial or Other Competing Interest: None.
Submission 12-07-2018, Peer Review 28-07-2018,
Acceptance 30-07-2018, Published 06-08-2018.

Corresponding Author:

Dr. Rishikant Vashistha,

Associate Professor,

2-BB, Slice No. 5, Scheme No. 78,

Vijay Nagar, Indore, Madhya Pradesh, India.

E-mail: vashistha.rishikant@gmail.com

DOI: 10.14260/jemds/2018/804

Perforation of small bowel leads to chemical and bacterial contamination, which may result in diffuse acute peritonitis. Patients with perforation are most of them usually presents with complaint of pain in abdomen, vomiting, fever and obstipation. In developing countries, perforation of small bowel due to typhoid and tuberculosis is common.^{3,4} Perforation of ileum due to typhoid is one of the commonest causes in developing countries, most probably due to low socio-economic condition, poor sanitation and poor personal hygiene. It is one of the most common causes of ileal perforation, which occur in 2nd and 3rd week of illness. The most lethal complications of typhoid fever are ileal perforation and intestinal bleeding both arising from necrosis of Peyer's patches in the terminal ileum. Typhoid ulcers can occur anywhere from stomach to rectum, but terminal ileum is more commonly involved due to higher density of Peyer's patches in the terminal ileum.^{5,6}

Despite aggressive management and best of critical care facilities available, morbidity and mortality rates remain high. With the availability of new better antibiotics and advances in anaesthesia technique, laparotomy remains the first line of treatment for perforated bowel. The present mortality due to diffuse suppurative peritonitis ranges from 10% to 20%, which continues to be unacceptably high.⁷ This study was done in our hospital to find out age and sex incidence, aetiological factors, clinical features and various surgical procedures for small bowel perforation and its complication and outcomes in our set-up.

MATERIALS AND METHODS

This cross-sectional observational study was conducted in the Department of General Surgery of Sri Aurobindo Medical College and Postgraduate Institute, Indore, over a period of 18 months between 1st January 2015 and 30th June 2016. The work was started after the review and approval of protocol of study by Institutional Ethics and Research Committees.

Study sample- 80 consecutive patients with diagnosis of non-traumatic small bowel perforation were included. Patients aged > 18 years were included in the study. Proper consent from the patient was taken for his/ her inclusion in this study. A detailed clinical history was obtained for all the patients with the emphasis on the presenting complaints. A thorough physical examination was done for all patients. Vital signs were recorded. Information was documented for demographical data (age, gender), clinical features (Abdominal pain, distension of abdomen, vomiting, fever, cold peripheries, decreased urine output, constipation, dyspnoea, duration of symptoms, clinical examination findings (pulse, BP, temperature and abdominal examination to see for tenderness, localised guarding, rigidity). All patients underwent exploratory laparotomy after routine haematological and radiological investigations and record was made about the intraoperative findings and type of procedure done, any postoperative complications were noted and final outcome of the patients was evaluated.

Statistical Analysis

The mean and percentage value were calculated by using simple statistical methods. The final data was presented in the form of graphs and tables.

RESULTS

The table shows the distribution of patients according to age group. There were 3 (3.8%) patients in the age group < 20 years, 20 (25%) were in the age group of 21 - 30 years, 19 (23.8%) were in the age group of 31 - 40 years, 9 (11.3%) were in the age group of 41 - 50 years and 17 (21.3%) were in the age group of 51 - 60 years and 12 (15%) were in the age group more than 60 years.

Sr. No.	Age Group	Number	Percentage
1	<= 20 years	3	3.8
2	21-30 years	20	25.0
3	31-40 years	19	23.8
4	41-50 years	9	11.3
5	51-60 years	17	21.3
6	> 60 years	12	15.0

Table 1. Distribution according to Age Group

In the present study, 73.7% of the patients were males and 26.3% were females. The male-to-female ratio was 2.8: 1

Sr. No.	Gender	Number	Percentage
1	Female	21	26.3
2	Male	59	73.7

Table 2. Distribution according to Gender

According to clinical presentation in 80 patients, abdominal pain in 80 (100%) patients, abdominal distension in 57 (71.3%) was present and obstipation in 70 (87.5%) were the commonest presenting symptoms.

Sr. No.	Clinical Presentation	Number	Percentage
1	Abdominal pain	80	100.0
2	Fever	51	63.7
3	Abdominal distension	57	71.3
4	Obstipation	70	87.5
5	Vomiting	44	55.0

Table 3. Distribution according to Clinical Presentation (Symptom)

Abdominal tenderness, guarding and rigidity with absent bowel sounds were the commonest presenting signs.

Sr. No.	Clinical Presentation	Number	Percentage (%)
1	Tachycardia	80	100
2	Tenderness	28	35
3	Guarding/ Rigidity	51	63.7
4	Absent bowel sound	70	87.5

Table 4. Distribution according to Clinical Presentation (Sign)

In our study typhoid in 66 (82.5%), ischaemic bowel disease in 6 (7.5%) and tuberculosis in 5 (6.25%) patients are the most common causes of perforation.

Sr. No.	Causes of Perforation	Number	Percentage
1	Typhoid	66	82.5
2	Tuberculosis	5	6.25
3	Ischaemic bowel disease	6	7.5
4	Malignancy	1	1.25
5	Intussusception	1	1.25
6	Worm infestation	1	1.25

Table 5. Causes of Perforation

In our study, ileal perforation 67 (83.8%) are more common than jejunal perforation 13 (16.3%).

Sr. No.	Site	Number	Percentage (%)
1	Ileum	67	83.8
2	Jejunum	13	16.3

Table 6. Following Anatomical Site of Perforation

There were single perforations, 67 (83.8%) were more common than multiple perforations 13 (16.3%).

Sr. No.	Number of Perforations	Number	Percentage
1	Single	67	83.8
2	Multiple	13	16.3

Table 7. Distribution according to Number of Perforations in each Patient

Distribution of patients according to the Procedures Performed

51 (63.75%) cases had undergone primary repair, followed by 11 (13.75%) resections and primary anastomosis, 6 (7.5%) resections and exteriorisation of bowel without anastomosis (N= 80).

Procedure	Number	Percentage
Primary repair	51	63.75%
Resection and primary anastomosis	11	13.75%
Resection and exteriorisation of bowel	6	7.5%
Resection and anastomosis with stoma	8	10.0%
Resection and exteriorisation of bowel with pre-op stenting of thrombosed artery	2	2.5%
Palliative drainage	2	2.5%
Total	80	100.0

Table 8. Distribution according to Types of Surgical Procedure Performed

In our study in 38 (47.5%) surgical site infection was seen, in 13 (16.3%) sepsis was seen and in 12(15%) patient's chest infection were the most common complication (N= 80).

Complications	Number	Percentage
Surgical site infection	38	47.5
Wound dehiscence	6	7.5
Sepsis	13	16.3
Chest infection	12	15
Faecal fistula	3	7.5
Death	8	10
Uneventful	15	18.8

Table 9. Postoperative Complication and Course

There were 8 (10%) deaths in our patients, while 72 (90%) patients were discharged. Majority of the patients in our study were discharged (N= 80).

Outcome	Number	Percentage
Death	8	10.0
Discharged	72	90.0
Total	80	100.0

Table 10. Distribution of Patients according to Outcome

DISCUSSION

Infection is the commonest cause of such perforations in developing countries. Small bowel perforation is one of the commonest surgical emergencies. Present study was carried out on 80 patients. The age range observed in the study was from 20 years - 90 years. The mean age of the study population was 43.36 ± 16.9 years (Ratio of males-to-female participants was 2.8: 1). Most of the patients presented in the age group of 21 - 30 years. This might be due to the fact that typhoidal perforation are seen in young age group 20 - 30 years. Similar observation was found in other studies of Rodolfo et al (2002),⁷ Dani et al (2015),⁸ Memon et al (2008).⁹

The most common complaint noticed in the study was abdominal pain followed by fever, abdominal distension, vomiting and constipation and the same being reported by Abdullah et al (2011).¹⁰ Ambikavathy et al (2013)¹¹ and

Shaikh et al (2009).¹² In our study, the main cause of perforation was due to typhoid. Other causes such as ischaemic bowel disease, tuberculosis, malignancy, intussusception and worm infestation were found in few cases. Typhoid fever was the predominant cause of non-traumatic small bowel perforation. It is the most common cause in developing countries due to lack of proper sanitation facilities. This was a similar finding in other studies where large bowel and malignant perforations were least common.^{11,13}

Majority of the patients had purulent type of peritoneal fluid. This was the similar finding in other studies.⁸ But in contrary percentage of patients having faecal peritoneal fluid was significantly more, since in our study most common site of perforation was found to be in ileum which was attributed to typhoid infection. It is one of the most common infectious causes of small bowel perforation in developing countries like ours with poor basic sanitation and drinking water facilities. Majority of the patients had diffuse type of peritonitis. Our hospital being a tertiary centre most of the patients comes late, referred from periphery. Majority of the patients had single perforation, which was similar finding in comparison to other studies.^{10,11}

CONCLUSION

There are innumerable causes, which may lead to small bowel perforation. Thus, there always persists a diagnostic dilemma in determining the aetiology leading to the small bowel perforation. Typhoid fever is the most common cause of ileal perforation followed by ischaemic bowel disease and tuberculosis.

Most of the patients presenting at tertiary centre are referred from peripheries with more diagnosis-surgery interval. Preoperative management of such cases are challenges and tests the skill techniques and knowledge of the surgeon.

No particular surgery is gold standard for the management of perforation peritonitis. Thus, every surgery is tailored in accordance with scenario on laparotomy, considering various parameters like age of patient, site, number and size of perforation, anaesthesia duration, type of peritoneal fluid, level of contamination, comfort of operating surgeon and protocol followed in particular institute.

REFERENCES

- [1] Abantanga FA, Nimako B, Amoah M. Perforation of the Gut in children as a result of enteric fever: a 5 years single institute review. *Annals of Pediatric Surgery* 2009;5(1):1-10.
- [2] Jhobta RS, Attri AK, Kaushik R, et al. Spectrum of perforation peritonitis in India-review of 504 consecutive cases. *World Journal of Emergency Surgery* 2006;1:26.
- [3] Wani RA, Parray FQ, Bhat NA, et al. Nontraumatic terminal ileal perforation. *World J Emerg Surg* 2006;1:7.
- [4] Atamanalp SS, Aydinli B, Ozturk G, et al. Typhoid intestinal perforations: twenty-six year experience. *World J Surg* 2007;31(9):1883-8.
- [5] Beniwal US, Jindal D, Sharma J, et al. Comparative study of operative procedures in typhoid perforation. *Indian J Surg* 2003;65:172-7.

- [6] Na'aya HU, Eni UE, Chama CM. Typhoid perforation in Maiduguri, Nigeria. *Ann Afr Med* 2004;3(2):69-72.
- [7] Bracho-Riquelme RL, Melero-Vela A, Torres-Ramirez A. Mannheim Peritonitis Index validation study at the hospital general de Durango (Mexico). *Cir Ciruj* 2002;70:217-25.
- [8] Dani T, Ramchandra L, Nair R, et al. Evaluation of prognosis in patients' with perforation peritonitis using Mannheim's peritonitis index. *International Journal of Scientific and Research Publications* 2015;5(5)1-35.
- [9] Memon AA, Bhutto AA, Shaikh SG, et al. Spectrum of diseases in patients with non-traumatic acute abdomen. *J Liaquat Univ Med Health Sci* 2008;7:180-3.
- [10] Abdullah MS, Rassam RE, Almarzooq TJ. A study of 82 patients of non-traumatic terminal ileal perforation in Al-Kindy teaching hospital. *J Fac Med Baghdad* 2011;2:147-51.
- [11] Ambikavathy M, Bhaskaran A, Kumar S, et al. Non-traumatic ileal perforation: Surgical experience in rural population in Indian scenario. *IJBAR* 2013;04(01):47-55.
- [12] Shaikh GS, Soomro Q, Bhutto AA, et al. Typhoid ileal perforation experience of 62 cases at Chandka Medical College Hospital, Larkana. *J Med Channel* 2009;15:187-90.
- [13] Malhotra MK, Singal R, Chowdhary K, et al. Spectrum of perforation peritonitis in a rural medical college. *Bangladesh Journal of Medical Science* 2016;15(1):70-3.