CASE REPORT

GOSSYPIBOMA: A RARE PRESENTATION

Veena Rahatgaonkar¹, U.T.Bhosale², Ankita Gawade³.

1. Associate Professor, Department of Gynaecology & Obstetrics, Bharati Vidyapeeth Medical College, Sangli.
2. Professor & Head of the Department, Department of Gynaecology & Obstetrics, Bharati Vidyapeeth Medical College, Sangli.
3. Junior Resident, Department of Gynaecology & Obstetrics, Bharati Vidyapeeth Medical College, Sangli.

CORRESPONDING AUTHOR

Dr.Veena Rahatgaonkar,
Swami Swarupananda Nursing Home,
Behind Wardhaman Plaza,
Kolhapur Road, Sangli, 416416,
Maharashtra, India.
Email: ven_rah@yahoo.com

HOW TO CITE THIS ARTICLE:

Veena Rahatgaonkar, U.T.Bhosale, Ankita Gawade. "Gossypiboma: A Rare presentation". Journal of Evolution of Medical and Dental Sciences 2013; Vol2, Issue 23, June 10; Page: 4094-4096.

ABSTRACT: Inadvertent retention of foreign materials in the body after surgical procedures continues to feature in the literature. Retained foreign material like sponge or instruments is a major contributing factor for increased morbidity. Gossypiboma (retained surgical sponge) is rare but preventable occurrence.

In this case it presented as chronic abdominal mass simulating ovarian tumour. Due to adhesions formation and encapsulation it was a diagnostic challenge for us.

KEY WORDS: Ovary, cotton, retained.

INTRODUCTION: Inadvertent retention of foreign materials in the body after surgical procedures continues to feature in the literature. Gossypiboma (retained surgical sponge) is rare but preventable occurrence. It is estimated that Gossypiboma may occur in 1 out of 1,000 - 1,500 intra-abdominal operations. [1]

In this case it presented as chronic abdominal mass simulating ovarian tumour in a lady who had undergone abdominal hysterectomy with right salpingo - oophorectomy.

CASE REPORT: A 55 years multiparous patient who had undergone abdominal hysterectomy 9 years back for abnormal uterine bleeding admitted in Bharati Vidyapeeth Medical College, Sangli. She presented with mass and pain in lower abdomen since 1 year prior to admission.

On examination, vital parameters were normal. A lump of size approximately 15 x 15 cm. with variable consistency and restricted mobility was present in lower abdomen. There was no free fluid in abdomen. Vaginal vault was healthy and same mass was felt on Per vaginal examination.

On Investigations, Ca125 was less than 4 units/ml.

On Ultrasonography of Pelvis, Uterus was not seen, mixed echogenic lesion of 113 x 101 mm arising from pelvis having solid and cystic component, suggestive of Dermoid was seen.

CASE REPORT

On CT examination of Pelvis, there was 12x12 cm cystic lesion in Right side of pelvis with well defined enhancing walls and internal nodules, ovarian in origin with bowel loops adherent to it. 5x3 cm cystic lesion was seen in left ovary.

Decision of exploratory laparotomy was taken. On opening abdomen large mass approximately 15x15 cm plastered with omentum was noted. Omental adhesions were separated. While delivering the tumor, it ruptured accidentally and thick yellow fluid of around 300cc with centrally located large mop adherent to inside of formed abscess cavity wall was seen. (Fig.1 and Fig.2)

The mop was removed along with the cyst wall. Left sided cystic ovary was removed. Right sided tube and ovary were not seen. Abdomen was closed in layers after achieving complete homeostasis. Instruments and mop count checked & found clear.

The post operative period was uneventful.

Histopathological examination revealed evidence of foreign body inflammation on right side and simple ovarian cyst on left Side.

DISCUSSION: Gossypiboma is retention of cotton material, gossypium (cotton) derived from Latin and boma (place of concealment) from Swahili [2]. Usually gauze or abdominal sponge is a frequent cause of Gossypiboma which causes morbidity and mortality. Patient may require another surgery to recover the material.

Acute presentation generally follows a septic course with abscess formation or a pyogenic foreign body granuloma. Foreign body may remain in abscess cavity and creates multiple fistulae and intestinal obstruction. In such case the patient who was recovering well after surgery complains of vague abdominal symptoms.

Delayed presentation may follow months or years after original surgical intervention . Due to adhesions formation and encapsulation it may present as intra-abdominal tumor which really is a diagnostic challenge for treating clinicians. The presentation may be like ovarian tumor as in present case or simulate primary small bowel tumor. Non specific clinical symptoms and inconclusive imaging findings may preclude an accurate diagnosis.

Risk of retention of foreign body is higher in emergency surgery, unexpected change in surgical procedure, obese patient and when mop count is omitted [3]. Change in assistant, multiple surgical teams, staff fatigue, prolongation of surgery is contributing factors for retention of sponge.

Routine Radiographic screening of patients before they leave the operation theatre is suggested in difficult and prolonged surgeries. Use of RFID (Radiofrequency Identification) tag on the gauzes is cost effective solution to avoid the complications related to retained surgical material [4, 5].

CONCLUSION: Medico legal consequences of Gossypiboma are significant. Considering mortality and morbidity associated with this condition, surgeons should make deliberate efforts to prevent it. This emphasizes importance of guidelines for operative theatre record keeping as recently set down by Royal college of Surgeons of England. All surgical textile material and instruments should be counted once at the start and twice at the end of surgery.

REFERENCES:

CASE REPORT

---

1. Lincourt AE, Harrel A, Cristiano J, Schrist C, Kercher K, Heniford BT. Retained foreign bodies after surgery. J surgery Res 2007; 138(2):170-174
2. Rajgopal A, Martin J. Gossypiboma – “a surgeon’s legacy”: report of a case and review of literature. Dis Colon Rectum 2002; 45(1): 119-120.
3. Gawande AA, Studdert DM, Orav EJ, Brennan TA, Zinner MJ. Risk factors for retention of instruments and sponges after surgery. N Engl J Med 2003; 348(3):229-35.
4. Rogers A, Jones E, Oleynikov D. Radio frequency identification (RFID) applied to surgical sponges. Surg Endosc 2007; 21(7):1235-7.
5. Radio-frequency surgical sponge detection: a new way to lower the odds of leaving sponges (and similar items) in patients. Health Devices 2008; 37(7):193-202.

FIGURE 1: Gossypiboma

FIGURE 2: Abdominal mass with sponge and thick yellow fluid coming out