ISSN 1313 - 8820
Volume 8, Number 1
March 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Tsanko Yablanski
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Co-Editor-in-Chief

Dimitar Panaiotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Atanas Atanasov (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Bojin Bojinov (Bulgaria)
Stoicho Metodiev (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Zervas Georgios (Greece)
Ivan Varlyakov (Bulgaria)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Georgi Petkov (Bulgaria)
Ramesh Kanwar (USA)
Martin Banov (Bulgaria)

Product Quality and Safety

Marin Kabakchiev (Bulgaria)
Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to editorial review and the editors reserve the right to improve style and return the paper

for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone.: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone.: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820

Volume 8, Number 1
March 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Partial factor productivity of nitrogen fertilizer on grain and grain protein yield of durum wheat cultivars

G. Panayotova^{1*}, S. Kostadinova²

¹Department of Plant Production, Faculty of Agriculture, Trakia University, 6000 Stara Zagora, Bulgaria

²Department of Agrochemistry and Soil Science, Faculty of Agronomy, Agricultural University, 12 Mendelev, 4000 Plovdiv, Bulgaria

(Manuscript received 19 January 2016; accepted for publication 2 February 2016)

Abstract. This study was conducted at the Field Crops Institute in Chirpan, Bulgaria, on Pellic vertisols soil type under non-irrigated conditions in the period of 2011–2013. Tests were made on the effect of nitrogen fertilization at rates of 0; 60; 120 and 180 kg N/ha on seven durum wheat cultivars. The cultivar response was studied regarding the yield produced out of a unit of applied nutrient. It was found that the partial nitrogen productivity of grain and protein decreased by increasing the applied amounts of nitrogen fertilizer into the soil. The rate of N₆₀ yielded the highest average partial productivity of nitrogen (69.8 kg/kg). The Elbrus and Deana cultivars had the highest partial nitrogen productivity of grain (47–50 kg N/kg). The Elbrus cultivar manifested a tendency for the highest efficiency in forming the grain protein yield (7.2 kg N/kg) out of a unit of applied nitrogen fertilizer. The partial nitrogen productivity increased in favourable temperature and moisture weather conditions during wheat vegetation.

Keywords: durum wheat, partial productivity, nitrogen, genotype

Abbreviations: PFP-N - Partial factor productivity of nitrogen

Introduction

Over the last years Bulgaria has increased its production of grain from durum wheat (*Triticum Durum* Desf.) as well as the consumption of products made of durum wheat. The issues of variety and applied farming practices, including levels of nitrogen fertilization are pressing for obtaining stable grain yields. The biology of durum wheat and the need to enhance the productivity require the application of optimal fertilizer rates which are in conformity with the area and field specificity. Most studies on wheat fertilization refer mainly to common wheat (*Triticum aestivum*) and only a limited number to durum wheat.

The modern varieties of durum wheat have higher productive potential but they realize it after greater nutrient input. The breeding process rarely considers the characteristics of the source forms regarding the mineral nutrition and does not envisage the possible results. Thus, over the last years the agrochemical evaluation of cultivars and hybrids was established as a compulsory part of modern breeding for high productivity (Sylvester-Bradley and Kindred, 2009; Johnson, 2004).

Human demands for food, fiber and biofuel production are rising with the world's population. It is estimated that almost half of the world's population is currently fed as a result of increased crop productivity due to the use of inorganic nitrogen fertilizers (Erismann et al., 2008). Use of mineral fertilizers sustains the world's growing population and sparing millions of hectares of natural and ecologically-sensitive systems that otherwise would have been converted to agriculture (Cassman, 1999). Today, economic and environmental challenges are driving increased interest in nitrogen use efficiency. Higher prices for both crops and fertilizers have focused interest in efficiency-improving technologies and practices

that also improve productivity (Ladha et al., 2005). Risks of increased environmental N losses via leaching, runoff, volatilization and denitrification, which may be associated with increased global N use and that harm air and water quality can be reduced by improving use efficiencies of nutrients (Kuzmanov and Tomov, 2000; Voicu and Soare, 2012). Since fertilizers are made from non-renewable resources, pressure to increase their use efficiencies will continue and at the same time, efforts should increase to enhance fertilizer use effectiveness for improved productivity and profitability of cropping systems (Horhota, 2012; Koteva and Marcheva, 2012b). Nitrogen fertilization is expected to increase to satisfy these growing human needs. Future fertilizer N management decisions must be increasingly based on economic, social, and environmental goals. On-farm N use efficiency and effectiveness can be improved, through better management of N sources, rates, timing, and placement (Koteva and Marcheva, 2012a). A goal of improving N use efficiency by 25% from current levels is considered achievable in the United States and may also be within the reach of many developing countries (Dobermann and Cassman, 2002).

Sustainable agricultural systems performance required to take account the effects in a long term period (Mosier et al., 2004). According to the best management practices (BMPs) the fertilizers efficiency is maximal when suitable sources of nutrients are applied at the exact rate, time and place – the 4R principle, which increases the yield, decreases the loss of nutrients and improves the soil fertility (Bruulsema et al., 2004; Fixen, 2009; Roberts, 2008). For certain soil types and cultivation conditions, the efficiency of nutrients use in most of the cases decreases with increase of the amount of applied element (Dibb, 2000). It is of particular importance for the cereals which are consumers of the greater part of the industrial fertilizers produced worldwide, but the efficiency principles and indicators are similar for all farm crops (Craswell and Godwin,

* e-mail: galia_panayotova@abv.bg

1984; Dawson et al., 2008; Deloqua et al., 1998; Dobermann, 2007; Moll et al., 1982; Weih et al., 2011; Witt et al., 2005).

Partial factor productivity (PFP) of nutrients due to factor fertilization answers the question what the productivity of a system considering the nutrient inputs is. This index is the simplest form of yield efficiency and is calculated per units of crop yields per unit nutrient element. It is used as long-term indicator of trends (Fixen, 2009). Higher levels indicate a higher amount of nutrient input, while lower indicate productivity limiting deficit. Typical values of the partial productivity of nitrogen are about 40 – 80 kg.kg⁻¹. The rates higher than 60 kg.kg⁻¹ are used in very efficiently managed systems, at low nitrogen rates or low soil nitrogen supply. PFP-N (often called nitrogen use efficiency or NUE) is changed from a high value at low N fertilization to lower values by increasing the nitrogen levels. PFP-N for cereals worldwide declined from 245 kg grain per kg N imported in 1961 – 1965 to 52 kg.kg⁻¹ in 1981 – 1985 and in recent years was an average of about 44 kg per kg N (Snyder and Bruulsema, 2007). The increase in yield with the increase of nitrogen fertilization was the reason for the decline. In many developed countries the yield of cereals continued to exceed the past 20 years without significant changes in fertilization. Sustainable PFP-N has increased in Western Europe, North America, Japan since the mid 1980s (Dobermann and Cassman, 2005).

Today, the average cereals yields in these regions are 60 to 100% higher than those worldwide at nitrogen rates only higher with 30 to 60% than the ones worldwide (Dobermann and Cassman, 2005). High yield and high PFP-N rates were the result of fertile soil, favorable weather conditions and excellent technologies. Giller et al. (2004) also reported that significant increases in nitrogen use efficiency are often achieved through reductions in N fertilizer use by 10 to 30%, while increases in yield tended to be small.

The cropping systems are a complex of soil, soil microorganisms, plants and crop rotation (Snyder, 2009). Improving the efficiency of one component may or may not be efficient for improving the efficiency of the cropping systems. A higher efficiency in a short-term period can lead to more expenses compared to the longer period. Lower fertilization rates in short term improve the efficiency of the nutrients even when the yields decrease. However, in long term the lower yields leave smaller amounts of plant residue and thus increase the risk of erosion, reduce organic matter in the soil and deteriorate the soil productivity (Snyder and Bruulsema, 2007).

According to Dobermann and Cassman (2005) 66% of fertilizer N was used to fertilize cereal crops, mainly corn and wheat. By raising N use efficiency and effectiveness through better cropping system and fertilizer N management, societal food fiber and biofuel demands may be met while also protecting air, water, and soil resources for current and future generations. A goal of improving N use efficiency by 25% from current levels is considered achievable in the United States and may also be within the reach of many developing countries (Dobermann and Cassman, 2002).

Since 1960, improved nitrogen efficiency has led to a sharp decline in the PFP-N on average 1 – 2% per year in developing countries (Dobermann and Cassman, 2005; Kitchen and Goulding, 2001). Too high values of this parameter in Africa (122 kg.kg⁻¹ imported N), Eastern Europe and Central Asia (84 kg.kg⁻¹) are indicators of unsustainable use of soil due to the low nitrogen requirements.

The partial productivity of the imported element is the most important indicator for grain farmers because it integrates the use efficiency of soil elements and the efficiency of applied fertilizers

(Doberman, 2007; Hawkesford, 2012; Snyder and Bruulsema, 2007). The partial productivity of nutrient is easily calculated for each farm that observes the holdings and income. Bulgaria doesn't have scientific data on indices of nitrogen uptake by newly established varieties of durum wheat.

The objective of this study was to examine the response of modern Bulgarian genotypes of durum wheat to nitrogen fertilization in relation with the formed production through one of the main indicators for nitrogen effectiveness – partial factor productivity per unit of applied nitrogen.

Material and methods

This study was conducted at the experimental field of the Field Crops Institute, Chirpan, with two field crops rotation (cotton-durum wheat) under rainy conditions in the period of 2011 – 2013. The experimental design consisted of split-plots design with four replications. The harvested size of the plots was 10 m². The treatments were as follows: 0; 60; 120 and 180 kg N.ha⁻¹. Nitrogen fertilization in the form of NH₄NO₃ was applied before sowing (1/3 of the rate) and in early spring at the beginning of vegetation (2/3 of the rate). The cotton predecessor was fertilized with P₈₀ in the form of triple superphosphate.

Seven Bulgarian cultivars of durum wheat were studied: Progress, Vazhod, Victoria, Predel, Deana, Zvezdica and Elbrus. The field trial was set by the common agrotechnology adopted for durum wheat cultivation in Bulgaria.

The main nitrogen use efficiency indicator of PFP was used to assess the durum wheat response to nitrogen (Dobermann, 2007). PFP was calculated on dry weight basis using the following formulas:

$$PFP = Y/F \text{ (kg.kg}^{-1}\text{)},$$

where Y is grain yields (grain protein) from fertilized treatments, and F is amount of N fertilizer applied (kg.ha⁻¹).

The soil type of the experimental field was *Pellic vertisols* (FAO) and generally refers to the so called Mediterranean chernozems. The soil type is one of the most generous, widely spread and significant in Bulgaria. It is suitable for growing most of the field crops and has a potential for high yield. The main parent materials are pliozen clay deposits. It has a high-powered humus horizon (70 – 100 cm), with a compact zone of the profile (united horizon). By humus content it belongs to the mean humus soils. It is characterized with high humidity capacity, caused by the high percentage of clay minerals, with clay soil texture, small water-permeability, bulk density of the arable soil layer 1.2-1.3 g.cm⁻³, with specific gravity 2.4 – 2.6 and low total porosity, weakly acid to neutral soil reaction and high cation exchange capacity (CEC) – 35-46 meq per 100 g soil, with high degree of bases saturation (93.4 – 100.0%), with total N in the arable layer 0.095 – 0.14% and low content of total phosphorus (0.06 – 0.11%), poor to medium supplied with hydrolyzed nitrogen, poorly supplied with available phosphorus and well-supplied with available potassium.

The agrochemical analyses from the field trial showed that the soil had neutral reaction and the content of mineral nitrogen was an average of 41.20 mg/kg soil for the layer of 0 – 20 cm (within 40.04 – 43.12 mg/kg soil), and under the plough layer it was 30.03 mg/kg soil, which varied from 29.26 to 30.80 mg/kg soil. The content of available phosphorus was low – an average of 3.9 mg/100 g soil in the 0 – 20 cm layer and from 2.6 to 3.0 mg/100 g soil for the 20 – 40

Table 1. Agrochemical characteristics of the soil, Chirpan, 2011

Depth, cm	pH _(KCl)	Mineral nitrogen, mg/kg soil			P ₂ O ₅ , mg/100 g	K ₂ O, mg/100 g
		NH ₄ -N	NO ₃ -N	N min		
Average 0 – 20	6.3	25.80	15.40	41.20	3.9	19.8
Average 20 – 40	6.5	20.02	10.01	30.03	2.8	17.8

cm soil layer. The soil was well-supplied with available potassium – an average of 19.8 mg/100 g soil (19.5 – 20.1 mg/100 g soil) in the 0–20 cm layer and 17.4 – 18.0 mg/100 g soil for the 20 – 40 cm layer (Table 1).

The experimental field of the Field Crops Institute in Chirpan is situated on the lowlands of the Upper Thracian Plain – 42°12'58" northern latitude, 25°17'00" eastern longitude and at 175 m altitude. According to the climatic zoning the region is transitional continental. As for precipitation and air temperature, the region is particularly suitable for durum wheat cultivation. Typical for this region is the low winter temperature and warm summer. The average annual temperature sum is 4317 °C. The coldest month is January with an average temperature of 0.3 °C, and the warmest month is July with an average temperature of 23.2°C. The annual average precipitation sum is 567 L/m², with its maximum in June – an average of 70.0 L/m². The annual minimum precipitation is in September.

Regarding the grain yield, crop year 2011 was the most favorable for the development, yield and quality of durum wheat, and 2012 was the most unfavorable. In the spring-summer period (March – June) of 2011 the temperature sum (1780.1 °C) exceeded the average for a long-term period by 122.7 °C. Unfavorable influence was exerted by the high temperatures in April – June and in the three years, as well as by the abundant precipitation in May and June of crop years 2012 and 2013. January and February of crop year 2012 reached the record low temperatures of -27°C, but it was a favorable condition for the wheat that those temperatures coincided with thick 25–30 cm snow cover. The abundant precipitation in January (97.0 L/m²) and February (61.3 L/m²), which significantly exceeded the norm, was in the form of snow and led to delay in plant development (Figures 1 and 2).

Figure 1. Average air temperature (°C) during the vegetation period of durum wheat, 2011 – 2013

During the vegetation period of the durum wheat, 2013 was characterized as warm and medium dry regarding the temperature and precipitation supply, with insufficient precipitation during the autumn months and during the period of ear formation – flowering. The temperature sum exceeded the average sum with 443.6 °C. The precipitation sum during the vegetation period was close to the average value but the values were lower during the mass tillering and ear formation-flowering (March and May), when the yield elements were set.

The data was statistically analyzed with the ANOVA procedure within the SPSS statistical program and Duncan's multiple range test ($P = 0.05$) to find significant differences among the average values.

Results and discussion

The differences in the grain yield of durum wheat as affected by the rates of nitrogen fertilization were significant (Table 2). Without fertilization the cultivars formed an average yield of 3.77 t.ha⁻¹. The nitrogen fertilization increased the yield up to N₁₂₀ – 4.80 t.ha⁻¹, which was 27% above unfertilized. The high N₁₈₀ rate reduced the yield compared to N₁₂₀.

Average for the period, the cultivars did not distinguish significantly by grain yield. Standard Progress was characterized with the lowest yield – 3.99 t.ha⁻¹, and the new cultivars Elbrus and Deana formed the highest yield, 4.72 and 4.47 t.ha⁻¹, respectively, which was 18% and 12% above standard. The Predel yield was the lowest. The highest average grain yield characterized the favorable 2011, 35% and 12% above 2012 and 2013.

In the interaction of nitrogen rate and cultivar, average for the

Figure 2. Monthly precipitation sum during the vegetation period of durum wheat, 2011 – 2013

Table 2. Grain yield (t.ha⁻¹) depending on the nitrogen rate and cultivar, 2011 – 2013

Main factors	Year			Average	
	2011	2012	2013	t.ha ⁻¹	%
A. N rate					
N ₀	4.12 ^d	3.09 ^b	4.09 ^b	3.77 ^c	100
N ₆₀	4.71 ^c	3.36 ^b	4.50 ^{ab}	4.19 ^b	111
N ₁₂₀	5.63 ^a	3.84 ^a	4.93 ^a	4.80 ^a	127
N ₁₈₀	5.20 ^b	4.27 ^a	4.40 ^{ab}	4.62 ^a	123
B. Cultivar					
Progress	4.76 ^{ns}	3.38 ^{ns}	3.84 ^b	3.99 ^{ns}	100
Vazhod	5.02	3.36	4.27 ^{ab}	4.22	106
Victoria	5.09	3.41	4.40 ^{ab}	4.30	108
Predel	4.90	3.28	4.26 ^{ab}	4.15	104
Deana	4.74	4.14	4.52 ^{ab}	4.47	112
Zvezdica	4.64	3.78	4.30 ^{ab}	4.24	106
Elbrus	5.25	4.11	4.79 ^a	4.72	118
C. Year - t.ha⁻¹					
	4.91 ^a	3.64 ^c	4.48 ^b	4.34	
- %	135	100	123		

^{a, b, c} – Means within a column having a different superscript letter differ significantly (p < 0.05)

period, the unfertilized Progress cultivar had the lowest yield – 3.26 t.ha⁻¹. The highest average values of grain yield were observed for Elbrus, fertilized at N₁₂₀ – 5.21 t.ha⁻¹ and with N₁₈₀ – 4.99 t.ha⁻¹, which was 60% and 53% above the unfertilized Progress and significantly exceeded all other interactions (Table 3). Average for the study, the highest yield was from Victoria x N₁₂₀ and Elbrus x N₆₀₋₁₂₀ in 2013 – 88 – 96% above standard, and the lowest was in 2012 from unfertilized Vazhod and Predel.

Increasing the nitrogen rate up to N₁₈₀ also increased the grain protein yield, reaching the highest average value of 718 kg.ha⁻¹, where the differences with N₁₂₀ were insignificant (Table 4). The medium and high nitrogen rates greatly affected the protein yield and were the most efficient for this parameter. This showed the strong influence of nitrogen fertilization on the yield of grain protein. The differences between N₀ and N₆₀ were insignificant. In 2011 and 2013, the yields of grain protein increased by increasing the imported nitrogen up to rate N₁₂₀, and in 2012 – up to N₁₈₀. There was no significant difference between the cultivars regarding the mean yield

of grain protein. The standard Progress was characterized with low protein yield – 534 kg.ha⁻¹. The cultivars Elbrus and Zvezdica distinguished with high values, reaching 675 and 630 kg.ha⁻¹, respectively, which was 26% and 18% above Progress. Elbrus had the highest protein yield in the three studied years.

The N rate x Cultivar interaction had very well proven differences related to the protein yield, where the highest average values were observed for Elbrus, with applied fertilizer at N₁₂₀ and N₁₈₀ – 823 and 785 kg.ha⁻¹, respectively, exceeding the unfertilized Progress with 92% and 83% (Table 5). For the grain protein yield the greatest effect was manifested by N – 45.7%, followed by year – 32.5%, cultivar – 7.4%, and the smallest insignificant effect was the N rate x Cultivar x Year interaction – 1.9% (Figure 3).

PFP is the most important index of total economic outputs for the cereal farmers relative to the use of all N sources (soil N and applied fertilizer) (Doberman, 2007; Hawkesford, 2012; Snyder and Bruulsema, 2007). The obtained values in the study on partial nitrogen productivity for grain and grain protein, average for the period, decreased by increasing the applied nitrogen amount

Table 3. Grain yield (t.ha⁻¹) at Nitrogen rate x Cultivar interaction, 2011 – 2013

Cultivar	N rate	N ₀	N ₆₀	N ₁₂₀	N ₁₈₀	Average for cultivar	
						t/ha	%
Progress		3.26	3.61	4.36	4.15	3.99 ^{ns}	100
Vazhod		3.55	4.15	4.69	4.70	4.22	106
Victoria		3.81	4.17	4.88	4.76	4.30	108
Predel		3.71	4.05	4.75	4.55	4.15	104
Deana		3.96	4.45	4.88	4.80	4.47	112
Zvezdica		3.82	4.15	4.83	4.40	4.24	106
Elbrus		4.26	4.72	5.21	4.99	4.72	118
Average for N - t/ha		3.77	4.19	4.80	4.62	4.34	
- %		100	111	127	123		

Table 4. Effect of nitrogen rate and cultivar on the yield of grain protein, kg.ha⁻¹

Main factors	Year			Average	
	2011	2012	2013	kg.ha ⁻¹	%
A. N rate					
N ₀	536 ^c	364 ^c	510 ^b	470 ^b	100
N ₆₀	613 ^b	414 ^c	576 ^b	534 ^b	114
N ₁₂₀	862 ^a	542 ^b	723 ^a	709 ^a	151
N ₁₈₀	833 ^a	636 ^a	684 ^a	718 ^a	153
B. Cultivar					
Progress	684 ^{ns}	451 ^{ns}	468 ^b	534 ^{ns}	100
Vazhod	678	429	577 ^b	561	105
Victoria	741	459	659 ^a	619	116
Predel	757	458	671 ^a	628	118
Deana	653	534	629 ^a	606	113
Zvezdica	695	535	661 ^a	630	118
Elbrus	768	558	699 ^a	675	126
C. Year - t.ha⁻¹					
- %	145	100	127		

^{a, b, c} – Means within a column having a different superscript letter differ significantly (p < 0.05)

Table 5. Effect of interaction of Nitrogen rate x Cultivar on grain protein yield (kg.ha⁻¹)

Cultivar	N rate	N ₀	N ₆₀	N ₁₂₀	N ₁₈₀	Average for cultivar	
						kg/ha ⁻¹	%
Progress		429	481	624	603	534	100
Vazhod		399	460	607	779	561	105
Victoria		478	544	744	710	619	116
Predel		495	554	732	732	62.8	118
Deana		469	526	703	723	606	113
Zvezdica		525	576	730	691	630	118
Elbrus		496	597	823	785	675	126
Average for N - t/ha		470	534	709	718		
- %		100	114	151	153		

Figure 3. Individual effect and interaction of factors N, genotype and year (%) on the grain and protein yield of durum wheat

Table 6. Partial factor productivity of nitrogen for grain yield (kg/kg N) at Nitrogen rate x Cultivar interaction

Main factors	PFP-N for grain				
	2011	2012	2013	Average	
				kg/kg N	%
A. N rate					
N ₆₀	78.5 ^a	56.0 ^a	74.9 ^a	69.8 ^a	174.5
N ₁₂₀	46.9 ^b	32.0 ^b	41.1 ^b	40.0 ^b	100.0
N ₁₈₀	28.9 ^c	23.7 ^c	24.4 ^c	25.7 ^c	64.2
B. Cultivar					
Progress	49.8 ^{ns}	34.2 ^{ns}	35.5 ^{ns}	39.8 ^{ns}	100.0
Vazhod	53.4	34.4	46.7	44.8	112.6
Victoria	53.5	34.4	48.8	45.5	114.3
Predel	50.8	33.6	48.0	44.1	110.8
Deana	49.6	43.1	48.9	47.2	118.6
Zvezdica	47.9	39.2	46.9	44.6	112.1
Elbrus	55.0	41.8	53.0	49.9	125.4

a, b, c – Means within a column having a different superscript letter differ significantly ($p < 0.05$)

(Tables 6 and 7). The reduction of values for the PFP-N index for the higher nitrogen fertilization rates was due to the fact that a unit of imported nutrient formed lower production compared to the lower rates. Respectively, the highest values, average for the period, were observed for the low nitrogen fertilization rate of 60 kg N/ha – 69.8 kg/kg for grain and 9.2 kg/kg for grain protein, and the plants used most efficiently the nutrient from the fertilizer and the soil and were proven to exceed the values at N₁₂₀ and N₁₈₀, by 74.5% and 171.5% for grain, and 58.6% and 135.8% for grain protein, respectively. Similar effect of the nitrogen fertilization was observed in each of the experimental years, where the PFP-N for grain and grain protein was proven to decrease with the increase of imported nitrogen.

In 2011 the values ranged within 28.9 – 78.5 kg/kg for grain and 4.4–11.0 kg/kg for grain protein, in 2012 – 23.7 – 56.0 kg/kg for grain and 3.5 – 6.9 kg/kg for grain protein, and in 2013 – 24.4 – 74.9 kg/kg

for grain and 3.8–9.6 kg/kg for grain protein. No cultivar specific was established towards the PFP-N for grain and grain protein for both the average for the period and the individual years, and the differences in values were insignificant and unproven. The standard Progress had the lowest average values for grain and grain protein, 39.8 and 5.5 kg/kg, respectively, whereas the new cultivar Elbrus had the highest index values – 49.9 and 7.2 kg/kg (Tables 6 and 7).

In the interaction of factors nitrogen fertilization and cultivar, the PFP-N for grain and grain protein average for the studied period was the highest for Elbrus, fertilized at the low rate of 60 kg N/ha – 78.7 kg/kg grain and 10.5 kg/kg grain protein, and the lowest for the standard Progress fertilized at the high rate of 180 kg N/ha – 23 kg/kg grain and 3.4 kg/kg grain protein (Tables 8 and 9). In 2011 and 2013 the values of PFP-N for grain and grain protein were the highest for N₆₀ x Elbrus, for grain 85.8 and 90.6 kg/kg, and for grain protein 13.0

Table 7. Partial factor productivity of nitrogen for grain protein (kg/kg N) at Nitrogen rate x Cultivar interaction

Main factors	PFP-N for grain protein				
	2011	2012	2013	Average	
				kg/kg N	% to St
A. N rate					
N ₆₀	11.0 ^a	6.9 ^a	9.6 ^a	9.2 ^a	158.6
N ₁₂₀	6.8 ^b	4.5 ^b	6.0 ^b	5.8 ^b	100.0
N ₁₈₀	4.4 ^c	3.5 ^c	3.8 ^c	3.9 ^c	67.2
B. Cultivar	6.9 ^{ns}	4.5 ^{ns}	4.9 ^{ns}		
Progress	7.4	4.3	5.9	5.5 ^{ns}	100.0
Vazhod	7.9	4.6	6.8	5.9	107.3
Victoria	7.1	4.7	7.0	6.4	116.4
Predel	7.3	5.5	6.5	6.2	112.7
Deana	6.9	5.5	6.9	6.4	116.4
Zvezdica	8.4	5.7	7.4	6.4	116.4
Elbrus				7.2	130.9

a, b, c – Means within a column having a different superscript letter differ significantly ($p < 0.05$)

Table 8. Partial factor productivity of nitrogen for grain (kg/kg N) at Nitrogen rate x Cultivar interaction, 2011 – 2013 (kg.kg⁻¹)

Factors	PFP-N for grain				
	2011	2012	2013	Average	
				kg/kg N	% to St
N ₆₀ x Progress	76.9	50.3	53.0	60.1	165.6
N ₆₀ x Vazhod	82.7	49.7	75.3	69.2	190.6
N ₆₀ x Victoria	80.5	51.1	77.0	69.5	191.5
N ₆₀ x Predel	76.2	50.5	76.0	67.6	186.2
N ₆₀ x Deana	76.3	67.8	78.6	74.2	204.4
N ₆₀ x Zvezdica	70.9	62.7	74.2	69.2	190.6
N ₆₀ x Elbrus	85.8	59.6	90.6	78.7	216.8
N ₁₂₀ x Progress - St	45.2	29.6	34.1	36.3	100.0
N ₁₂₀ x Vazhod	47.1	29.9	40.2	39.1	107.7
N ₁₂₀ x Victoria	49.3	29.3	43.5	40.7	112.1
N ₁₂₀ x Predel	46.5	28.8	43.4	39.6	109.1
N ₁₂₀ x Deana	44.7	36.0	41.4	40.7	112.1
N ₁₂₀ x Zvezdica	46.8	32.5	41.4	40.2	110.7
N ₁₂₀ x Elbrus	48.7	37.9	43.6	43.4	119.6
N ₁₈₀ x Progress	27.2	22.6	19.4	23.0	63.4
N ₁₈₀ x Vazhod	30.3	23.5	24.6	26.1	71.9
N ₁₈₀ x Victoria	30.6	22.8	25.9	26.4	72.7
N ₁₈₀ x Predel	29.9	21.4	24.5	25.3	69.7
N ₁₈₀ x Deana	27.7	25.5	26.7	26.6	73.3
N ₁₈₀ x Zvezdica	25.9	22.3	25.1	24.5	67.5
N ₁₈₀ x Elbrus	30.5	27.8	24.8	27.7	76.3
Year	51.4 ^a	37.2 ^b	46.8 ^{ab}	45.14	-

Table 9. PFP-N for grain protein (kg N/kg) in factor correlation with nitrogen fertilization x cultivar. 2011–2013

Factors	Year			Average	
	2011	2012	2013	kg/kg N	% to St
N ₆₀ x Progress	10.0	6.3	7.1	7.8	150.0
N ₆₀ x Vazhod	11.3	5.6	8.6	8.5	163.5
N ₆₀ x Victoria	11.5	6.4	9.8	9.2	176.9
N ₆₀ x Predel	10.4	6.6	10.3	9.1	175.0
N ₆₀ x Deana	11.0	7.8	9.5	9.5	182.7
N ₆₀ x Zvezdica	9.7	8.5	10.4	9.5	182.7
N ₆₀ x Elbrus	13.0	7.1	11.4	10.5	201.9
N ₁₂₀ x Progress - St	6.7	4.1	4.8	5.2	100.0
N ₁₂₀ x Vazhod	6.6	3.7	5.2	5.2	100.0
N ₁₂₀ x Victoria	7.4	4.2	6.5	6.0	115.4
N ₁₂₀ x Predel	6.5	4.2	6.7	5.8	111.5
N ₁₂₀ x Deana	6.6	5.0	5.9	5.8	111.5
N ₁₂₀ x Zvezdica	6.7	4.8	6.3	5.9	113.5
N ₁₂₀ x Elbrus	7.4	5.7	6.9	6.7	128.8
N ₁₈₀ x Progress	4.1	3.2	2.8	3.4	65.4
N ₁₈₀ x Vazhod	4.3	3.6	4.0	4.0	76.9
N ₁₈₀ x Victoria	4.8	3.3	4.0	4.0	76.9
N ₁₈₀ x Predel	4.3	3.3	3.9	3.8	73.1
N ₁₈₀ x Deana	4.2	3.7	4.0	4.0	76.9
N ₁₈₀ x Zvezdica	4.2	3.4	3.9	3.8	73.1
N ₁₈₀ x Elbrus	4.8	4.3	4.0	4.3	82.7
Year	7.4 ^a	5.0 ^b	6.5 ^{ab}	6.3	-

and 11.4 kg/kg, respectively, whereas in 2012 were the highest for grain for N_{60} x Deana – 67.8 kg/kg and for grain protein for N_{60} x Zvezdica – 8.5 kg/kg. The index values for grain protein in the three studied years were the lowest for the correlation of the standard Progress fertilized with 180 kg N/ha, 4.1; 3.2 and 2.8 kg/kg, respectively, and for grain in 2013 for the same interaction – 19.4 kg/kg, and in 2011 for N_{180} x Zvezdica – 25.9 kg/kg and in 2012 for N_{120} x Predel – 21.4 kg/kg.

Among the studied years, average for all cultivars and fertilization rates, the highest values of partial factor productivity of nitrogen were observed in the favorable 2011 – 51.4 kg/kg grain and 7.4 kg/kg grain protein, which were proven to exceed the lowest values reported in 2012 by 38% (for grain) and by 48% (for grain protein), whereas the second position was taken by 2013 with 10% (for grain) and 14% (for grain protein) less than 2011 but there was no significant difference. The higher values of the PFP-N in the first year of study resulted from the higher yields of grain and protein under the influence of the favorable combination of temperature and precipitation. The sudden drought and high temperatures in the grain filling period (June) caused the low productivity in 2012 and as a result the values of PFP-N were the lowest (Tables 3 and 4).

Conclusions

Average for the period, the grain yield at fertilization rates of N_{60} , N_{120} and N_{180} increased with 11, 27 and 23% compared to the unfertilized control. Fertilization rates of 120 and 180 kg N/ha increased the grain protein yield with 51 – 53%. To realize high yield the optimal level of fertilization out of the tested ones was 120 kg N/ha. The highest average grain yield was obtained in the climatically favorable 2011 – 4.91 t/ha. Regarding the grain yield there was efficient interaction between cultivar Elbrus and N_{120} , which exceeded the control with 60% and significantly exceeded all other interactions. The new cultivar Elbrus stood out as the most perspective in relation to grain and protein yield. The protein content increased with the increase of nitrogen fertilization. The strong effect on the protein yield was due to the medium and high nitrogen rate. The cultivar characteristics did not influence the grain protein yield significantly.

The agrochemical assessment of durum wheat showed that the partial factor productivity of nitrogen for grain and protein reduces with the increase of the imported quantity of nitrogen fertilizer in the soil. Rate N_{60} led to obtaining the highest average PFP-N at 69.8 kg/kg.

Regarding the genotype response of formed production per unit of imported nitrogen the cultivars Elbrus and Deana distinguished with the highest PFP-N for grain – 47.2 – 49.9 kg/kg N, where the plants used the nutrient most efficiently. The Elbrus cultivar manifested a trend of highest efficiency in forming grain protein yield (7.2 kg/kg N) per unit of imported nitrogen fertilizer. Favorable hydrothermal conditions during wheat vegetation assisted the higher PFP-N.

References

- Bruulsema TW, Fixen PE and Snyder CS, 2004. Fertilizer nutrient recovery in sustainable cropping systems. *Better Crops*, 88, 15-17.
- Cassman K, 1999. Ecological intensification of cereal production systems: Yield potential, soil quality, and precision agriculture. *Proceedings of the National Academy of Science of the United States of America*, 96, 5952-5959.
- Cassman KG, Dobermann A and Walters DT, 2004. Agroecosystems, nitrogen-use efficiency, and nitrogen management. *Ambio*, 31, 132-140.
- Craswell A and Godwin P, 1984. The efficiency of N fertilizers applied to cereals in different climates. *Advances in Plant Nutrition*, 1, 1-55.
- Dawson JC, Huggins DR and Jones SS, 2008. Characterizing nitrogen use efficiency in natural and agricultural ecosystems to improve the performance of cereal crops in low-input and organic agricultural systems. *Field Crops Research*, 107, 89-101.
- Delogu G, Cattivellia L, Pecchionia N and De Falcis D, 1998. Uptake and agronomic efficiency of nitrogen in winter barley and winter wheat. *European Journal of Agronomy*, 9, 1, 11-20.
- Dibb DW, 2000. The mysteries (myths) of nutrient use efficiency. *Better Crops*, 84, 3-5.
- Dobermann A, 2007. Nutrient use efficiency – measurement and management, 22 p. In: Krauss A, Isherwood K, Heffer P, editors. *Proc. of International Fertilizer Industry Association (IFA) Workshop on Fertilizer Best Management Practices*. Brussels, Belgium. March, 7-9, 2007, 28.
- Dobermann A and Cassman K, 2002. Plant nutrient management for enhanced productivity in intensive grain production systems of the United States and Asia. *Plant Soil*, 247, 153-175.
- Dobermann A and Cassman K, 2005. Cereal area, yield and nitrogen use efficiency are drivers for future nitrogen fertilizer consumption. *Science in China*, 48, 745-758.
- Erismann JW, Sutton MA, Galloway J, Klimont Z and Winiwarter W, 2008. How a Century of ammonia synthesis changed the world. *Nature Geoscience*, 1, 636-639.
- Fixen P, 2009. Nutrient use efficiency in the context of sustainable agriculture, In J. Espinosa and F. García (Eds.), *Nutrient Use Efficiency*, International Plant Nutrition Institute (IPNI), USA, 1-10.
- Giller K, Chalk P, Dobermann A, Hammond L, Heffer P, Ladha J, Nyamudeza P, Maene L, Ssali H and Frene J, 2004. Emerging technologies to increase the efficiency of use of fertilizer nitrogen. In: AR Mosier et al. (ed.) *Agriculture and the nitrogen cycle: assessing the impacts of fertilizer use on food production and the environment*. SCOPE 65. Island Press, Washington, p. 35-52.
- Hawkesford M, 2012. The diversity of nitrogen use efficiency for wheat varieties and the potential for crop improvement. *Better Crops*, 96, 10-15.
- Horhota L, 2012. Economic efficiency and environmental protection in agriculture. *Journal of Environmental Protection and Ecology*, 13, 811.
- Kitchen NR and Goulding KWT, 2001. On-farm technologies and practices to improve nitrogen use efficiency. In: *Nitrogen in the environment: sources, problems, and management*, eds. RF Follett and JL Hatfield, 335-370. The Netherlands, Amsterdam: Elsevier.
- Koteva V and Marcheua M, 2012a. Productivity of common wheat variety Mirjana grown with reduced mineral fertilizing. *Soil Science, Agrochemistry and Ecology*, 3, 55-62 (Bg).
- Koteva V and Marcheua M, 2012b. Productivity of barley variety Veslec grown with reduced mineral fertilizing. *Agricultural Sciences*, 4, 7-13 (Bg).
- Kuzmanov N and Tomov A, 2000. Seasonal monitoring of flowing and non-moving water sources in Parvomaj region in Bulgaria. *J. of Environm. Protection and Ecology*, 1, 324.
- Ladha JK, Pathak H, Krupnik TJ, Six J and Van Kessel C, 2005.

Efficiency of fertilizer nitrogen in cereal production: retrospects and prospects. *Advances in Agronomy*, 87, 85-156.

Moll R, Kamprath E and Jackson W, 1982. Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agronomy Journal*, 74, 562-564.

Mosier AR, Syers JK and Freney JR, 2004. Agriculture and the Nitrogen Cycle. Assessing the Impacts of Fertilizer Use on Food Production and the Environment. Scientific Committee on Problems of the Environment (SCOPE). Island Press. Washington, DC.

Roberts TL, 2008. Improving nutrient use efficiency. *Turk Journal of Agriculture and Forestry*, 32, 177-182.

Snyder C, 2009. Nutrient use efficiency: global challengers, trends and the future. *IPNI*, p. 10-18.

Snyder C and Bruulsema T, 2007. Nutrient use efficiency and effectiveness in North America: Indices of agronomic and

environmental benefit. International Plant Nutrition Institute, Norcross, Georgia, 3-15.

Sylvester-Bradley R and Kindred DR, 2009. Analysing nitrogen responses of cereals to prioritize routes to the improvement of nitrogen use efficiency. *Journal of Experimental Botany*, 60, 1939-1951.

Voicu Gh and Soare A, 2012. Protection of waters against pollution by nitrites and nitrates from agricultural sources. *Journal of Environmental Protection and Ecology*, 13, 69.

Weih M, Asplund L and Bergkvist G, 2011. Assessment of nutrient use in annual and perennial crops: A functional concept for analyzing nitrogen use efficiency. *Plant Soil*, 339, 513-520.

Witt C, Pasuquin JMCA, Muters R and Buresh RJ, 2005. New leaf color chart for effective nitrogen management in rice. *Better Crops*, 89, 36-39.

Review

- Classical and modern concepts of inbreeding and effects of inbreeding depression in animals** 3
S. Tanchev

Genetics and Breeding

- Genotype by environment interaction in mutant lines of winter barley for grain yield** 14
B. Dyulgerova, N. Dyulgerov
- Genotype-environment interaction and stability analysis for grain yield of winter barley in the conditions of North-East and South Bulgaria** 19
M. Dimitrova-Doneva, D. Valcheva, G. Mihova, B. Dyulgerova

Production Systems

- Effect of predecessors on the productivity and phytosanitary condition of hull-less oats in organic farming** 24
D. Atanasova, V. Maneva, T. Nedelcheva
- Partial factor productivity of nitrogen fertilizer on grain and grain protein yield of durum wheat cultivars** 28
G. Panayotova, S. Kostadinova
- Influence of the dimensions of lifting brushes on the losses at direct harvesting of standing vine dry bean** 37
I. Iliev, G. Milev
- Energy productivity, fertilization rate and profitability of wheat production after various predecessors** 41
II. Profitability of wheat production
Z. Uhr, E. Vasileva
- Selectivity and stability of new herbicides and herbicide combinations for the seed yields of some field crops I. Effect at coriander (*Coriandrum Sativum* L.)** 46
G. Delchev
- Determination of some macro and micro elements in grain of winter barley genotypes** 51
N. Markova Ruzdik, Lj. Mihajlov, V. Ilieva, S. Ivanovska, D. Valcheva, B. Balabanova, M. Ilievski

Agriculture and Environment

- Effects of irrigation and fertilization on soil microorganisms** 58
T. Dinev, I. Gospodinov, A. Stoyanova, G. Beev, D. Dermendzhieva, D. Pavlov

Investigation on some biotic factors in carp fish ponds	62
D. Terziyski, H. Kalcheva, A. Ivanova, R. Kalchev	
Investigation of some energy characteristics of pig farm	70
P. Kostov, K. Atanasov, I. Ivanov, K. Peychev, R. Georgiev	
Variability in the resistance to bacterial spot causal agents <i>Xanthomonas euvesicatoria</i> P and <i>Xanthomonas vesicatoria</i> PT2 among Bulgarian and introduced pepper varieties	75
T. Vancheva, S. Masheva, D. Ganeva, N. Bogatzevska	
Comparative analysis for macro and trace elements content in goji berries between varieties from China and R. Macedonia	79
B. Balabanova, I. Karov, S. Mitrev	
 Product Quality and Safety	
Extraction and characterization of anthocyanin colorants from plant sources	85
S. Dyankova, M. Doneva	
Heavy metal content in the meat of common carp (<i>Cyprinus carpio</i> L.) and rainbow trout (<i>Oncorhynchus mykiss</i> W.), cultivated under different technologies	90
St. Stoyanova, I. Sirakov, K. Velichkova, Y. Staykov	

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings:

Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 8, Number 1
March 2016

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz