ISSN 1313 - 8820
Volume 8, Number 3
September 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Editor-in-Chief

Georgi Petkov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Co-Editor-in-Chief

Dimitar Panayotov
Faculty of Agriculture
Trakia University, Stara Zagora
Bulgaria

Editors and Sections

Genetics and Breeding

Tsanko Yablanski (Bulgaria)
Atanas Atanasov (Bulgaria)
Nikolay Tsenov (Bulgaria)
Max Rothschild (USA)
Ihsan Soysal (Turkey)
Horia Grosu (Romania)
Bojin Bojinov (Bulgaria)
Stoicho Metodiev (Bulgaria)
Svetlana Georgieva (Bulgaria)

Nutrition and Physiology

Nikolai Todorov (Bulgaria)
Peter Surai (UK)
Zervas Georgios (Greece)
Ivan Varlyakov (Bulgaria)

Production Systems

Radoslav Slavov (Bulgaria)
Dimitar Pavlov (Bulgaria)
Bogdan Szostak (Poland)
Banko Banev (Bulgaria)
Georgy Zhelyazkov (Bulgaria)

Agriculture and Environment

Ramesh Kanwar (USA)
Martin Banov (Bulgaria)
Peter Cornish (Australia)

Product Quality and Safety

Marin Kabakchiev (Bulgaria)
Stefan Denev (Bulgaria)
Vasil Atanasov (Bulgaria)
Roumiana Tsenkova (Japan)

English Editor

Yanka Ivanova (Bulgaria)

Scope and policy of the journal

Agricultural Science and Technology /AST/ – an International Scientific Journal of Agricultural and Technology Sciences is published in English in one volume of 4 issues per year, as a printed journal and in electronic form. The policy of the journal is to publish original papers, reviews and short communications covering the aspects of agriculture related with life sciences and modern technologies. It will offer opportunities to address the global needs relating to food and environment, health, exploit the technology to provide innovative products and sustainable development. Papers will be considered in aspects of both fundamental and applied science in the areas of Genetics and Breeding, Nutrition and Physiology, Production Systems, Agriculture and Environment and Product Quality and Safety. Other categories closely related to the above topics could be considered by the editors. The detailed information of the journal is available at the website. Proceedings of scientific meetings and conference reports will be considered for special issues.

Submission of Manuscripts

All manuscripts written in English should be submitted as MS-Word file attachments via e-mail to editoffice@agriscitech.eu. Manuscripts must be prepared strictly in accordance with the detailed instructions for authors at the website

www.agriscitech.eu and the instructions on the last page of the journal. For each manuscript the signatures of all authors are needed confirming their consent to publish it and to nominate an author for correspondence.

They have to be presented by a submission letter signed by all authors. The form of the submission letter is available upon request from the Technical Assistance or could be downloaded from the website of the journal. Manuscripts submitted to this journal are considered if they have submitted only to it, they have not been published already, nor are they under consideration for publication in press elsewhere. All manuscripts are subject to editorial review and the editors reserve the right to improve style and return the paper

for rewriting to the authors, if necessary. The editorial board reserves rights to reject manuscripts based on priorities and space availability in the journal.

The journal is committed to respect high standards of ethics in the editing and reviewing process and malpractice statement. Commitments of authors related to authorship are also very important for a high standard of ethics and publishing. We follow closely the Committee on Publication Ethics (COPE), <http://publicationethics.org/resources/guidelines>

The articles appearing in this journal are indexed and abstracted in: DOI, EBSCO Publishing Inc. and AGRIS (FAO).

The journal is accepted to be indexed with the support of a project № BG051PO001-3.3.05-0001 "Science and business" financed by Operational Programme "Human Resources Development" of EU. The title has been suggested to be included in SCOPUS (Elsevier) and Electronic Journals Submission Form (Thomson Reuters).

Address of Editorial office:

Agricultural Science and Technology
Faculty of Agriculture, Trakia University
Student's campus, 6000 Stara Zagora
Bulgaria

Telephone.: +359 42 699330
+359 42 699446

www.agriscitech.eu

Technical Assistance:

Nely Tsvetanova
Telephone.: +359 42 699446
E-mail: editoffice@agriscitech.eu

ISSN 1313 - 8820

Volume 8, Number 3
September 2016

AGRICULTURAL SCIENCE AND TECHNOLOGY

2016

An International Journal Published by Faculty of Agriculture,
Trakia University, Stara Zagora, Bulgaria

Agriculture and Environment

Effect of some herbicides on weeds and vines in mother plantation of Cabernet sauvignon

N. Prodanova – Marinova*

Institute of Viticulture and Enology, 5800 Pleven, Bulgaria

(Manuscript received 1630 May 2016; accepted for publication 17 August 2016)

Abstract. To maintain soil surface clean of weeds in the parent vineyard for producing cuttings for scions, the efficiency and selectivity of Gardoprim plus Gold (312.5 g/l s-metolachlor + 187.5 g/l terbuthylazine), Wing P (pendimethalin 250 g/l + dimethenamid P 212.5 g/l) and Lumax 538 SC (375 g/l s-metolachlor + 125 g/l terbuthylazine + 337.5 g/l mesotrione) at doses of 0.4 and 0.6 l/da was studied. Lumax 538 SC, Wing P and Gardoprim Plus Gold controlling efficiently annual weeds established in parent vineyard, except *Xanthium strumarium* L.. Lumax 538 SC and Gardoprim plus Gold showed long-term (90 days) activity. Wing P activity decreases after the thirtieth day. The tested herbicides do not damage the buds and do not lead to a reduction of shoots developed from them. Lumax 538 SC and Gardoprim plus Gold at doses of 0.4 and 0.6 l/da and Wing P at dose of 0.4 l/da do not inhibit the growth of shoots. The largest number of cuttings for scions were obtained after treatment with Lumax 538 SC.

Keywords: parent vineyard, herbicides, shoots, cuttings for scions

Introduction

The production of quality cuttings is essential for the yield of grafted rooted vines. The precise choice of land and training systems for creating mother plantations, as well as the cultivation practices for vine growing are of great importance (Kandeva, 1986; Todorov, 2005). Maintaining the soil surface free of weeds eliminates the competition for the absorption of major nutrients and creates conditions for active vine growth. It has been found that bindweed species (*Convolvulus arvensis* L.), mustard (*Sinapis arvensis* L.), amaranth (*Amaranthus retroflexus* L.) and twitch (*Cynodon dactylon* Pers.) transpire much more intensively than vine. In addition to more water, weeds absorb from the soil greater quantities of nutrients (Boychev, 1980; Steenwerth et al., 2016). At an average density of weeds they extract from the soil 42 kg/ha nitrogen, phosphorus and potassium, while at high density - 186 kg/ha. Often a direct correlation has been observed between the degree of weed infestation and yield (Smirnov et al., 1987). The greater weed infestation usually results in inhibition of growth, less-developed shoots, lower cluster weight and lower yield per hectare (Moulis, 1992; Muganu et al., 2013).

The application of herbicides leads to a reduction or complete destruction of harmful plants in the vineyard and the proper selection of the active substances have a positive effect on vine growth. It has been shown that after prolonged use of herbicides (e.g., triazine group), vine growth is enhanced. The plants have better developed shoots and their annual growth is higher compared to the traditionally cultivated vineyards (Nikov et al., 1972; Magriso et al., 1979; Patil, 2006).

Hilling up and low-stem training of mother plantations in the region of Northern Bulgaria preclude the application of products based on glyphosate that have been registered for use in our country. An efficient way for weed control in such plantations is the

combination of soil herbicides imported immediately after the hilled up soil has been pulled away with appropriate herbicides for Johnson grass during vegetation. Thus the rotation of different active substances would contribute to more effective weed control, preventing to a great extent the negative consequences from flora evolution in the treated plantations (Boyer, 2004; Carsouille, 2004).

The objective of this study was to determine the efficiency and selectivity of some soil herbicides to weed species and vines in a mother vineyard; to determine the appropriate doses for treatment in plantations with ground trained vines.

Material and methods

A field trial was carried out in 2014 and 2015 for investigating the opportunities for application of chemical weed control in the mother vineyard with the following herbicides: Gardoprim plus Gold (312.5 g/l s-metolachlor + 187.5 g/l terbuthylazine), Wing P (pendimethalin 250 g/l + dimethenamid P 212.5 g/l) and Lumax 538 SC (375 g/l s-metolachlor + 125 g/l terbuthylazine + 337.5 g/l mesotrione). The treatment was performed immediately after the hilled up soil was pulled away before the weeds growth, without incorporation. The trial was conducted in the Experimental Base of the Institute of Viticulture and Enology - Pleven in plantation of Cabernet Sauvignon clone D-1. The soil type was slightly leached chernozem formed on clayish loess. The vine loading from all variants was preliminary made equal - 6 spurs per vine (12 winter eyes).

The variants were as follows:

V1 - Lumax 538 SC (4 l/ha);

V2 - Wing P (4 l/ha);

V3 - Wing P (6 l/ha);

V4 - Gardoprim plus Gold (4 l/ha);

V5 - Gardoprim plus Gold (6 l/ha);

* e-mail: neli_npm@abv.bg

C – Untreated cultivated control.

They were set in four repetitions with area of 12 m². The total area of each variant was 48 m², and the number of vines - 40.

The dynamics of the species composition and weed density (pc./m²) was monitored as the recordings were made in constantly marked in each repetition plots with area of 1m² on the 30th, 60th, 90th and 120th post-treatment day. For determining the effect of the tested herbicides on vines the following was recorded (5 vines per variant): number of germinated shoots, growth dynamics (cm) and number of cuttings for scions. The data were processed by analysis of variance (Dimova and Marinkov, 1999).

Results and discussion

The species diversity analysis showed a small number of weed species, which are typical for the region: field bindweed

(*Convolvulus arvensis* L.), whitetop (*Cardaria draba* L.), field sowthistle (*Sonchus arvensis* L.), Canada thistle (*Cirsium arvense* L.), field mustard (*Sinapis arvensis* L.), Black bindweed (*Falopia convolvulus* L.), chickweed (*Stellaria media* L.), Xanthium (*Xanthium strumarium* L.), White amaranth (*Amaranthus blitoides* L.), red-root amaranth (*Amaranthus retroflexus* L.), lamb's quarter (*Chenopodium album* L.), European heliotrope (*Heliotropium europaeum* L.), green foxtail (*Setaria viridis* L.), Johnson grass (*Sorghum halepense* L.).

The composition of the weed association has been changing over time and after the sixtieth day ephemera (chickweed) and early spring weeds (field mustard) gave way to late spring ones (Xanthium, amaranth species, orache, Setaria, etc.). Weed infestation with field bindweed and Johnson grass was constant throughout the growing season, as their density varied and it was insignificantly affected by the applied herbicides. The herbicides inhibited seed germination but did not kill the existing perennial

Table 1. Weed density per species in 2014 – 30th day and 60th day (pcs./m²)

Weed species	Density (pcs./m ²)											
	V1		V2		V3		V4		V5		C	
	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day
1 <i>Conv. arvensis</i>	56	74.5	22.5	40.5	48.0	67.0	49.5	63.5	41.5	45.0	160	185
2 <i>Cardaria draba</i>	2.5	4.5	4.0	10.0	6.0	10.5	6.0	7.5	8.5	13.5	4.5	7.0
3 <i>Sonchus arvensis</i>	-	-	-	1.0	-	0.5	-	-	-	-	1.0	3.0
4 <i>Cirsium arvense</i>	-	-	-	-	-	-	-	-	-	-	-	2.0
5 <i>Sinapis arvensis</i>	-	-	-	-	-	-	-	-	-	-	2.0	1.0
6 <i>Stellaria media</i>	-	-	-	-	-	-	-	-	-	-	1.0	-
7 <i>X. strumarium</i>	0.5	0.5	1.0	2.0	0.5	2.0	7.5	13.0	2.5	9.5	6.5	6.5
8 <i>A. blitoides</i>	-	-	-	-	-	-	-	-	-	-	23.5	25.0
9 <i>Ch. album</i>	-	-	-	3.0	-	1.0	-	-	-	-	13.5	15.0
10 <i>Setaria viridis</i>	-	-	-	0.5	-	0.5	-	-	-	-	8.5	35.5
11 <i>Sorghum halepense</i>	-	0.5	1.0	3.0	0.5	4.0	7.0	9.5	1.0	2.5	-	1.0
Σ	59.0	80.0	28.5	58.0	55.0	85.0	70.0	93.5	53.5	70.5	220.5	281.0

Table 2. Weed density per species in 2014 - 90th day and 120th day (pcs./m²)

Weed species	Density (pcs./m ²)											
	V1		V2		V3		V4		V5		C	
	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day
1 <i>Conv. arvensis</i>	81.0	83.5	46.0	54.5	70.5	83.0	73.5	79.5	55.0	67.5	83.5	51.0
2 <i>Cardaria draba</i>	5.0	5.0	11.0	12.0	9.5	9.5	8.0	9.5	15.0	16.0	3.0	1.0
3 <i>Sonchus arvensis</i>	-	-	1.5	1.5	0.5	0.5	-	1.0	-	-	0.5	3.0
4 <i>Cirsium arvense</i>	-	2.0	0.5	0.5	0.5	0.5	1.5	1.5	-	-	1.0	-
5 <i>X. strumarium</i>	1.0	1.0	2.0	1.0	5.0	5.0	18.5	14.0	9.5	10.5	2.5	0.5
6 <i>A. retroflexus</i>	-	-	-	-	-	-	-	-	-	-	1.0	1.5
7 <i>A. blitoides</i>	-	-	-	-	-	-	-	-	-	-	7.5	9.5
8 <i>Ch. album</i>	-	-	3.0	3.5	3.0	3.0	-	0.5	-	-	9.5	2.0
9 <i>H. europaeum</i>	-	-	3.0	3.0	1.0	1.0	-	0.5	-	0.5	3.0	2.5
10 <i>Setaria viridis</i>	-	3.0	3.0	3.0	1.0	3.0	-	3.0	0.5	0.5	0.5	8.5
11 <i>Sorghum halepense</i>	2.0	3.0	7.0	15.0	9.5	12.5	11.0	23.0	7.5	8.0	12.0	8.5
Σ	89.0	97.5	77.0	94.0	100.5	117.0	112.5	132.5	87.5	103.0	124	88.0

stems and rhizomes. The tested substances did not have any effect on *Xanthium* when applied to the soil, as in this case.

Wing P had remained active in the soil for 30 – 40 days and in 2014, on the sixtieth day in the plots treated with both doses plants of field sowthistle (0.5 pcs./m² for V2 and 1 pcs./m² for V3), orache (1 pcs./m² for V2 – 3 pcs./m² for V3) and *Setaria* (0.5 pcs./m²) were recorded – Table 1. Lumax 538 SC and Gardoprim plus Gold showed significantly greater persistency. On the ninetieth day the weed infestation in these variants was formed only by perennial weeds and *Xanthium* and the findings have shown that they continue to control the annual species afterwards, too (Table 2).

The high humidity in the spring of 2014 increased the density of weeds in the control variant – on the 30th day it was 220.5 pcs./m², while on the 60th day, in spite of the manual treatment, it reached 281 pcs./m². With the implementation of the monthly trenching round the

vines the undesired vegetation thereafter decreased and on the 125th day it was 88 pcs./m², as for the first time it was lower than in the treated variants.

In 2015, despite the different weather conditions, the tested herbicides showed the same permanence of action (Tables 3 and 4). For Wing P (V2), except the perennial weeds, it was reported 0.2 pcs./m² plants of European heliotrope (V3) and 0.8 pcs./m² black bindweed (V3). In both variants with this herbicide on the sixtieth day and after that higher density of the weeds was observed compared to the control. The most efficient control over the annual species throughout the vegetation season showed Lumax 538 SC at a dose of 4 l/ha (V1) and Gardoprim plus Gold at a dose of 6 l/ha. An increase in the density of *Xanthium* compared to 2014 was found in all variants, especially those treated with Wing P. On the 120th day *Xanthium* plants reached 30 pcs./m² (V2) and 31.8 pcs./m² (V3). In

Table 3. Weed density per species in 2015 - 30th day and 60th day (pcs./m²)

Weed species	Density (pcs./m ²)											
	V1		V2		V3		V4		V5		C	
	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day	30 th day	60 th day
1 <i>Conv. arvensis</i>	22.8	34.4	24.6	35.2	28.4	38.6	24.0	32.8	21.4	29.8	55	40
2 <i>Cardaria draba</i>	3.0	3.6	5.8	6.2	1.2	1.2	4.4	2.4	6.2	6.8	1.2	0.8
3 <i>Sonchus arvensis</i>	3.0	3.0	3.6	1.2	-	-	0.4	0.2	0.8	0.8	0.2	0.2
4 <i>Cirsium arvense</i>	0.4	3.0	-	-	0.2	0.4	-	-	-	-	0.4	-
5 <i>Sinapis arvensis</i>	-	-	-	-	0.2	-*	-	-	-	-	-	-
6 <i>Falopia convolvulus</i>	-	-	-	0.8	-	-	-	-	-	-	1.0	1.4
7 <i>X. strumarium</i>	17.4	19.2	25.6	29.0	28.4	30.0	7.0	7.4	22.6	23.6	12.4	18.4
8 <i>A. blitoides</i>	-	-	-	-	-	-	-	-	-	-	2.0	2.6
9 <i>H. europaeum</i>	-	-	-	0.2	-	-	-	-	-	-	2.0	0.4
10 <i>Ch. album</i>	-	-	-	-	-	-	-	-	-	-	0.4	0.4
11 <i>Setaria viridis</i>	-	-	-	-	-	-	-	-	-	-	2.8	5.8
12 <i>Sorghum halepense</i>	0.6	6.0	7.6	8.2	0.6	2.0	10.2	15.4			2.6	2.0
Σ	47.2	69.2	67.2	80.8	59.0	72.2	46.0	58.2	51.0	61	80.0	72.0

Table 4. Weed density per species in 2015 - 90th day and 120th day (pcs./m²)

Weed species	Density (pcs./m ²)											
	V1		V2		V3		V4		V5		C	
	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day	90 th day	120 th day
1 <i>Conv. arvensis</i>	35.8	39.2	37.4	40.6	38.4	41.8	32.8	37.6	29.8	31.8	31.6	30.4
2 <i>Cardaria draba</i>	3.6	3.6	6.2	6.2	1.2	1.2	2.4	4.8	6.8	7.0	5.6	1.2
3 <i>Sonchus arvensis</i>	3.0	3.0	1.2	1.2	-	0.2	0.2	0.4	0.8	0.8	-	0.6
4 <i>Cirsium arvense</i>	3.0	3.0	-	-	0.4	0.6	-	-	-	-	-	-
5 <i>X. strumarium</i>	20.0	22.2	29.0	30.0	30.4	31.8	7.4	8.2	23.6	24.2	13	10.0
6 <i>Falopia convolvulus</i>	-	-	0.8	0.8	-	-	-	-	-	-	-	-
7 <i>A. blitoides</i>	-	-	-	0.8	-	-	-	-	-	-	7.8	1.2
8 <i>Ch. album</i>	-	-	-	1.4	-	1.4					2	2.4
9 <i>H. europaeum</i>	-	-	0.2	0.4	-	0.2	-	-	-	-	1.6	3.4
10 <i>Setaria viridis</i>	-	2.2	-	2.4	-	1.2	-	2.4	-	2.0	2.4	9.2
11 <i>Sorghum halepense</i>	6.0	8.8	9.0	10.2	2.8	2.8	16.4	17.8	-	-	12.0	13.0
Σ	71.4	82.0	83.8	94.8	73.2	81.2	59.2	71.2	61.0	65.8	76.0	71.4

2015, there were no chickweeds both in the treated variants and in the control.

Besides on the soil surface during the treatment, the herbicide solution fell on vines. At this time (immediately after the hilled up soil had been pulled away) the buds have not yet broken open and the hulls covering them protect the future shoots from direct contact with the tested substances. The data for both years were meaningful. Twenty-five days post-treatment with herbicides the average number of germinated shoots varied from 24.3 pcs. (V2) to 26.7 pcs.

per vine (V4) - Figure 1. Minimum reduction compared to the control was recorded only for Wing P - 4 l/ha (V2) - 1.1 pc. and Wing P - 6 l/ha (V3) - 1 pc., as it was due to the smaller number of sprouts. The shoots from spurs ranged from 13.2 (V1) to 13.9 per vine (V3 and V4), while in control they were 13.3 pcs. That had definitely suggested that the tested herbicides introduced immediately after the hilled up soil had been pulled away did not damage the buds and did not lead to a reduction of the shoots that developed from the buds.

Figure 1. Number of shoots per vine, average for the period 2014 – 2015

Table 5. Growth dynamics

V	Shoot average length (cm)							
	35 th day		65 th day		95 th day		125 th day	
	2014	2015	2014	2015	2014	2015	2014	2015
C	35.2	100.8	135.8	220.7	202.8	333.1	253.4	348.6
V1	42.4	108.8	174.4	229.4	247.0	364.9	292.3	388.9
V2	32.8	105.5	150.0	240.5	247.7	339.8	298.1	367.6
V3	28.9	82.5	123.3	205.5	231.2	279.5	279.6	308.5
V4	36.7	104.0	162.9	241.7	261.6	350.6	296.1	387.5
V5	42.1	94.1	168.1	200.5	293.0	282.0	322.7	316.5

During the vegetation period of 2014 there was no evidence of negative effects of the tested active substances on the vine growth. Thirty-five days after treatment, the shoots of all variants (except V3) had more active growth compared to the control (Table 5). The vines of V3 (Wing P - 6 l/ha) showed some delay in growth, but it was overcome after the sixty-fifth day and on the 125th day after treatment the shoot length from that variant was already 26.2 cm greater than

that of the control. The longest shoots had the vines from V5 (Gardoprim plus Gold at a dose of 6 l/ha – 322.7 cm. The exceeding compared to the control for V5 reached 69.3 cm.

In 2015 a delay in growth was reported compared to the control again for V3 and V5 (Gardoprim plus Gold - 6 l/ha, resulting in reduction in the average length of shoots by 40.1 cm for V3 and 32.1 cm for V5. The longest shoots on the sixty-fifth day and all

Figure 2. Average number of cuttings per vine for the period 2014 – 2015

subsequent recordings were measured in V1 (Lumax 538 SC). At the end of the vegetation the average length for this variant surpassed the control by 40.3 cm.

The shoot length at the end of the vegetation period did not always determine the obtaining of a greater number of high-quality cuttings. The cuttings for scions (variants V1 and V4) were practically equal in length, however the number of cuttings obtained from them was significantly different (Figure 2). In the variant treated with Lumax 538 SC (V1) 192.8 pcs. of cuttings for scions on average were reported for the period 2014 - 2015, which undoubtedly exceeded the control and all the other variants, while with Gardoprim plus Gold at a dose of 4 l/ha (V4) their number was 153.7 pcs. The difference between the variants is statistically significant at $P < 0.01$.

Conclusions

Lumax 538 SC, Wing P and Gardoprim plus Gold, applied to the soil immediately after the hilled up soil had been pulled away, control efficiently the annual weeds found in the mother vineyard, except *Xanthium* (*Xanthium strumarium* L.). Lumax 538 SC and Gardoprim plus Gold exhibited longer action (90 days). The action of Wing P decrease after the thirtieth day. The tested herbicides did not damage the buds and did not lead to a reduction of shoots developed from them and did not affect adversely their growth. Lumax 538 SC provided optimal conditions for vine growth and obtaining the greatest number of quality cuttings.

References

- Boyer J, Carsouille J and Crozier Ph**, 2004. Vigne, alterner les herbicides de prélevée : Quatre essais de six ans chacun montrent l'intérêt de programmes pluriannuels raisonnés, 577, 35-37.
- Boychev A**, 1980. Studies on water and nutrition regimen of weeds in the vineyards. Horticultural and viticulture Science, 2, 98-103 (Bg).
- Carsouille J, Boyer J and Crozier Ph**, 2004. Vigne, réduire les doses d'herbicides de prélevée : Trois années, dix essais de désherbage mixte. Phytoma-La Défense des végétaux, 577, 30-34.
- Dimova D and Marinkov E**, 1999. Experimental Works and Biometry. Academic publishing house of HAI, Plovdiv, 263 (Bg).
- Kandeva R**, 1986. The cuttings quality – important condition for obtaining a high rate of first-class vines. Viticulture and Horticulture, 5, 30-32 (Bg).
- Magrisso Y, Yonev S, Cholakov T, Nikov M, Danailov B, Raykov E, Georgiev A, Dimchev V and Babrikov D**, 1979. Commercial Technologies in viticulture. Sofia, Zemizdat, 322 (Bg).
- Moulis I**, 1992. Etude de la concurrence herbe – vigne. Conséquences écophysologiques sur la vigne de l'enherbement de vignoble en région Méditerranéenne. Comptes rendues du IV simposium international de physiologie de la vigne, 11 – 15 mai, Torino, 93-97.
- Muganu M, Paolucci M, Gnisci D, Barnaba FE, Bellincontro A, Mencarelli F and Grosu I**, 2013. Effect of Different Soil Management Practices on Grapevine Growth and on Berry Quality Assessed by NIR-AOTF Spectroscopy Acta horticulturae, 978, 117-125.
- Nikov M**, 1972. Growth and fruit-bearing of vines with prolonged use of herbicides without tillage. Horticultural and viticulture Science, IX, 7, 137-146 (Bg).
- Patil D, Sulikeri G and Patil H**, 2008. Studies on the weed management practices in Thompson seedless grape vineyard. Acta Horticulturae, 785, 321-326.
- Smirnov K, Kalmykov F and Morozov G**, 1987. Viticulture, Agropromizdat, 877 (Ru).
- Steenwerth K, Orellana-Calderón A, Hanifin R, Storm C and McElrone A**, 2016. Effects of Various Vineyard Floor Management Techniques on Weed Community Shifts and Grapevine Water Relations. American Journal of enology and viticulture, 67, 2, 153-162.
- Todorov I**, 2005. Production of grapevine propagation material. Dionis, Sofia, 303 (Bg).

Review

- Honey bees and their products as indicators of environmental pollution: A review** 175
D. Salkova, M. Panayotova-Pencheva

Genetics and Breeding

- Characterization of the Bulgarian sunflower hybrid Valin** 183
G. Georgiev
- Evaluation of the combining ability of mutant maize lines** 189
V. Valkova, N. Petrovska
- Evaluation of small size fruit peppers *Capsicum annuum* spp. *microcarpum* with cluster and factor analysis** 193
V. Kuneva, M. Nikolova
- Sensitivity of promising cherry hybrids and new cultivars to economically important fungal diseases** 197
K. Vasileva, S. Malchev, A. Zhivondov

Nutrition and Physiology

- Lysozyme levels in haemolymph of worker bees (*Apis mellifera* L.) from bee colonies with different degree of expression of hygienic behaviour** 201
S. Lazarov, I. Zhelyazkova, D. Salkova, R. Shumkova, S. Takova

Production Systems

- Study on energy flows of renewable sources for producing hot water on dairy farms** 205
R. Georgiev, K. Peychev, D. Georgiev, R. Slavov, S. Apostolov, J. Ellingsen, J. Tønnesen
- Loose smut of barley grown in three types of farming** 209
T. Nedelcheva, V. Maneva
- Efficacy and timing of some new products against pear psylla (*Cacopsylla pyri* L.) (Hemiptera: Psyllidae): I. Spirotetramat** 213
V. Arnaudov
- Influence of year's characteristics and the different fertilization levels on the structural elements of wheat yield** 217
V. Kuneva, R. Bazitov, A. Stoyanova

Grain combines productivity according to various unloading methods – in the field and at the edge of the field **221**

N. Delchev, K. Trendafilov, G. Tihanov, Y. Stoyanov

Agriculture and Environment

Effect of some herbicides on weeds and vines in mother plantation of Cabernet sauvignon **227**

N. Prodanova – Marinova

Influence of foliar herbicides treatment on malting barley (*Hordeum vulgare* L.) productivity of Emon, Vanessa and Vicky varieties **232**

D. Atanasova, V. Maneva

Selectivity and stability of herbicides and herbicide combinations for the grain yield of maize (*Zea Mays* L.) **237**

G. Delchev, T. Barakova

Effect of some soil herbicides on vegetative habits of almond trees of 'Nonpareil' cultivar grown in a second-year nursery field **242**

Z. Rankova, M. Tityanov

Phytosanitary conditions of the organic field and boundary **245**

D. Atanasova, V. Maneva, N. Grozeva

Product Quality and Safety

Quality traits of eggs from autosexing Easter eggers **249**

H. Lukanov, A. Genchev, A. Pavlov, I. Penchev

Amino acid composition of lamb meat from the North East Bulgarian fine fleece breed and its crossbreds from internal breeding **256**

R. Slavov, G. Mihaylova, St. Ribarski, D. Panayotov, D. Pamukova, D. Dragnev

Some results of evaluation of new-introduced apricot cultivars under conditions of Plovdiv region **262**

V. Bozhkova, M. Nesheva

Instruction for authors

Preparation of papers

Papers shall be submitted at the editorial office typed on standard typing pages (A4, 30 lines per page, 62 characters per line). The editors recommend up to 15 pages for full research paper (including abstract, references, tables, figures and other appendices).

The manuscript should be structured as follows: Title, Names of authors and affiliation address, Abstract, List of keywords, Introduction, Material and methods, Results, Discussion, Conclusion, Acknowledgements (if any), References, Tables, Figures.

The title needs to be as concise and informative about the nature of research. It should be written with small letter /bold, 14/ without any abbreviations.

Names and affiliation of authors

The names of the authors should be presented from the initials of first names followed by the family names. The complete address and name of the institution should be stated next. The affiliation of authors are designated by different signs. For the author who is going to be corresponding by the editorial board and readers, an E-mail address and telephone number should be presented as footnote on the first page. Corresponding author is indicated with *.

Abstract should be not more than 350 words. It should be clearly stated what new findings have been made in the course of research. Abbreviations and references to authors are inadmissible in the summary. It should be understandable without having read the paper and should be in one paragraph.

Keywords: Up to maximum of 5 keywords should be selected not repeating the title but giving the essence of study.

The introduction must answer the following questions: What is known and what is new on the studied issue? What necessitated the research problem, described in the paper? What is your hypothesis and goal?

Material and methods: The objects of research, organization of experiments, chemical analyses, statistical and other methods and conditions applied for the experiments should be described in detail. A criterion of sufficient information is to be possible for others to repeat the experiment in order to verify results.

Results are presented in understandable

tables and figures, accompanied by the statistical parameters needed for the evaluation. Data from tables and figures should not be repeated in the text.

Tables should be as simple and as few as possible. Each table should have its own explanatory title and to be typed on a separate page. They should be outside the main body of the text and an indication should be given where it should be inserted.

Figures should be sharp with good contrast and rendition. Graphic materials should be preferred. Photographs to be appropriate for printing. Illustrations are supplied in colour as an exception after special agreement with the editorial board and possible payment of extra costs. The figures are to be each in a single file and their location should be given within the text.

Discussion: The objective of this section is to indicate the scientific significance of the study. By comparing the results and conclusions of other scientists the contribution of the study for expanding or modifying existing knowledge is pointed out clearly and convincingly to the reader.

Conclusion: The most important consequences for the science and practice resulting from the conducted research should be summarized in a few sentences. The conclusions shouldn't be numbered and no new paragraphs be used. Contributions are the core of conclusions.

References:

In the text, references should be cited as follows: single author: Sandberg (2002); two authors: Andersson and Georges (2004); more than two authors: Andersson et al. (2003). When several references are cited simultaneously, they should be ranked by chronological order e.g.: (Sandberg, 2002; Andersson et al., 2003; Andersson and Georges, 2004).

References are arranged alphabetically by the name of the first author. If an author is cited more than once, first his individual publications are given ranked by year, then come publications with one co-author, two co-authors, etc. The names of authors, article and journal titles in the Cyrillic or alphabet different from Latin, should be transliterated into Latin and article titles should be translated into English. The original language of articles and books translated into English is indicated in parenthesis after the bibliographic reference (Bulgarian = Bg, Russian = Ru, Serbian = Sr, if in the Cyrillic, Mongolian =

Mo, Greek = Gr, Georgian = Geor., Japanese = Ja, Chinese = Ch, Arabic = Ar, etc.)

The following order in the reference list is recommended:

Journal articles: Author(s) surname and initials, year. Title. Full title of the journal, volume, pages. Example:

Simm G, Lewis RM, Grundy B and Dingwall WS, 2002. Responses to selection for lean growth in sheep. *Animal Science*, 74, 39-50

Books: Author(s) surname and initials, year. Title. Edition, name of publisher, place of publication. Example:

Oldenbroek JK, 1999. Genebanks and the conservation of farm animal genetic resources, Second edition. DLO Institute for Animal Science and Health, Netherlands.

Book chapter or conference proceedings:

Author(s) surname and initials, year. Title. In: Title of the book or of the proceedings followed by the editor(s), volume, pages. Name of publisher, place of publication. Example:

Mauff G, Pulverer G, Operkuch W, Hummel K and Hidden C, 1995. C3-variants and diverse phenotypes of unconverted and converted C3. In: *Provides of the Biological Fluids* (ed. H. Peters), vol. 22, 143-165, Pergamon Press. Oxford, UK.

Todorov N and Mitev J, 1995. Effect of level of feeding during dry period, and body condition score on reproductive performance in dairy cows. IXth International Conference on Production Diseases in Farm Animals, September 11-14, Berlin, Germany.

Thesis:

Hristova D, 2013. Investigation on genetic diversity in local sheep breeds using DNA markers. Thesis for PhD, Trakia University, Stara Zagora, Bulgaria, (Bg).

The Editorial Board of the Journal is not responsible for incorrect quotes of reference sources and the relevant violations of copyrights.

Animal welfare

Studies performed on experimental animals should be carried out according to internationally recognized guidelines for animal welfare. That should be clearly described in the respective section "Material and methods".

AGRICULTURAL SCIENCE AND TECHNOLOGY

Volume 8, Number 3
September 2016

Journal web site:
www.agriscitech.eu

Publisher:
www.alfamarket.biz